

# Differentiability Properties of Lipschitzian Bilinear Operators in Real Normed Spaces<sup>1</sup>

Kazuhisa Nakasho   
Yamaguchi University  
Yamaguchi, Japan

Yasunari Shidama  
Karuizawa Hotch 244-1  
Nagano, Japan

**Summary.** This article is devoted to the Mizar formalization of various properties of differentiability of Lipschitzian bilinear operators in real normed spaces. Main results include the Lipschitz continuity of partial derivatives, the representation of the total derivative in terms of partial derivatives, and the continuous differentiability of Lipschitzian bilinear operators on open subsets of the product space.

MSC: 46G05 47A07 68V20

Keywords: bilinear operator; Lipschitz continuity; normed space

MML identifier: NDIFF12, version: 8.1.14 5.85.1476

## INTRODUCTION

In this article we continue the formalization of selected properties of differentiability of Lipschitzian bilinear operators in real normed spaces [8], using the Mizar system [1], [2]. As a natural continuation of [9] it covers topics such as partial differentiability, continuity, and total differentiability of these operators [6], [13]. The work extends results for linear operators to the bilinear case [3] and provides theorems on the behavior of differential operators up to arbitrary order. Key results include the Lipschitz continuity of partial derivatives (Section 1), the representation of the total derivative in terms of partial derivatives (Sect. 2), and the continuous differentiability of Lipschitzian bilinear operators [11], [12] on open subsets of the product space in Section 3.

---

<sup>1</sup>This work was supported by JSPS KAKENHI Grant Number 24K14897.

## 1. FUNDAMENTAL PROPERTIES AND PARTIAL DIFFERENTIABILITY

From now on  $E, F, G, S, T, W, Y$  denote real normed spaces,  $f, f_1, f_2$  denote partial functions from  $S$  to  $T$ ,  $Z$  denotes a subset of  $S$ , and  $i, n$  denote natural numbers.

Now we state the propositions:

- (1) Let us consider a bilinear operator  $f$  from  $E \times F$  into  $G$ , and a point  $z$  of  $E \times F$ . Then

- (i)  $f \cdot (\text{reproj1}(z))$  is a linear operator from  $E$  into  $G$ , and
- (ii)  $f \cdot (\text{reproj2}(z))$  is a linear operator from  $F$  into  $G$ .

PROOF: Reconsider  $L_1 = f \cdot (\text{reproj1}(z))$  as a function from  $E$  into  $G$ . For every elements  $x, y$  of  $E$ ,  $L_1(x + y) = L_1(x) + L_1(y)$ . For every vector  $x$  of  $E$  and for every real number  $a$ ,  $L_1(a \cdot x) = a \cdot L_1(x)$ . Reconsider  $L_2 = f \cdot (\text{reproj2}(z))$  as a function from  $F$  into  $G$ . For every elements  $x, y$  of  $F$ ,  $L_2(x + y) = L_2(x) + L_2(y)$ . For every vector  $x$  of  $F$  and for every real number  $a$ ,  $L_2(a \cdot x) = a \cdot L_2(x)$ .  $\square$

- (2) Let us consider a Lipschitzian bilinear operator  $f$  from  $E \times F$  into  $G$ , and a point  $z$  of  $E \times F$ . Then

- (i)  $f \cdot (\text{reproj1}(z))$  is a Lipschitzian linear operator from  $E$  into  $G$ , and
- (ii)  $f \cdot (\text{reproj2}(z))$  is a Lipschitzian linear operator from  $F$  into  $G$ , and
- (iii) there exists a point  $g$  of  $\text{NormSpaceOfBoundedBilinOps}_{\mathbb{R}}(E, F, G)$  such that  $f = g$  and for every vector  $x$  of  $E$ ,  $\|(f \cdot (\text{reproj1}(z)))(x)\| \leq \|g\| \cdot \|(z)_2\| \cdot \|x\|$  and for every vector  $y$  of  $F$ ,  $\|(f \cdot (\text{reproj2}(z)))(y)\| \leq \|g\| \cdot \|(z)_1\| \cdot \|y\|$ .

PROOF: Reconsider  $g = f$  as a point of  $\text{NormSpaceOfBoundedBilinOps}_{\mathbb{R}}(E, F, G)$ . Set  $K = \|g\|$ . Reconsider  $L_1 = f \cdot (\text{reproj1}(z))$  as a linear operator from  $E$  into  $G$ . Reconsider  $L_2 = f \cdot (\text{reproj2}(z))$  as a linear operator from  $F$  into  $G$ . Set  $K_1 = K \cdot \|(z)_2\|$ . Set  $K_2 = K \cdot \|(z)_1\|$ . For every vector  $x$  of  $E$ ,  $\|L_1(x)\| \leq K_1 \cdot \|x\|$  by [7, (16)]. For every vector  $y$  of  $F$ ,  $\|L_2(y)\| \leq K_2 \cdot \|y\|$  by [7, (16)].  $\square$

- (3) Let us consider a Lipschitzian bilinear operator  $f$  from  $E \times F$  into  $G$ . Then there exists a real number  $K$  such that

- (i)  $0 \leq K$ , and
- (ii) for every point  $z$  of  $E \times F$ , for every point  $x$  of  $E$ ,  $\|(f \cdot (\text{reproj1}(z)))(x)\| \leq K \cdot \|(z)_2\| \cdot \|x\|$  and for every point  $y$  of  $F$ ,  $\|(f \cdot (\text{reproj2}(z)))(y)\| \leq K \cdot \|(z)_1\| \cdot \|y\|$ .

PROOF: Consider  $K$  being a real number such that  $0 \leq K$  and for every vector  $x$  of  $E$  and for every vector  $y$  of  $F$ ,  $\|f(x, y)\| \leq K \cdot \|x\| \cdot \|y\|$ . Set  $L_1 = f \cdot (\text{reproj}_1(z))$ . Set  $K_1 = K \cdot \|(z)_2\|$ . For every vector  $x$  of  $E$ ,  $\|L_1(x)\| \leq K_1 \cdot \|x\|$ .  $\square$

- (4) Let us consider a Lipschitzian bilinear operator  $f$  from  $E \times F$  into  $G$ , and a point  $z$  of  $E \times F$ . Then
- (i)  $f$  is partially differentiable in  $z$  w.r.t. 1, and
  - (ii)  $\text{partdiff}(f, z) \text{ w.r.t. } 1 = f \cdot (\text{reproj}_1(z))$ , and
  - (iii)  $f$  is partially differentiable in  $z$  w.r.t. 2, and
  - (iv)  $\text{partdiff}(f, z) \text{ w.r.t. } 2 = f \cdot (\text{reproj}_2(z))$ .

The theorem is a consequence of (2).

- (5) Let us consider points  $s, t$  of  $E \times F$ , and a real number  $a$ . Then
- (i)  $s = \langle (s)_1, (s)_2 \rangle$ , and
  - (ii)  $((s + t))_1 = (s)_1 + (t)_1$ , and
  - (iii)  $((s + t))_2 = (s)_2 + (t)_2$ , and
  - (iv)  $((s - t))_1 = (s)_1 - (t)_1$ , and
  - (v)  $((s - t))_2 = (s)_2 - (t)_2$ , and
  - (vi)  $(a \cdot s)_1 = a \cdot ((s)_1)$ , and
  - (vii)  $(a \cdot s)_2 = a \cdot ((s)_2)$ .
- (6) Let us consider a Lipschitzian bilinear operator  $f$  from  $E \times F$  into  $G$ . Then there exists a real number  $K$  such that
- (i)  $0 \leq K$ , and
  - (ii) for every point  $z$  of  $E \times F$ ,  $\|\text{partdiff}(f, z) \text{ w.r.t. } 1\| \leq K \cdot \|z\|$  and  $\|\text{partdiff}(f, z) \text{ w.r.t. } 2\| \leq K \cdot \|z\|$ .

The theorem is a consequence of (3), (2), (4), and (5).

## 2. TOTAL DIFFERENTIABILITY AND CONTINUITY

Let us consider a Lipschitzian bilinear operator  $f$  from  $E \times F$  into  $G$ . Now we state the propositions:

- (7) (i) for every points  $z_1, z_2$  of  $E \times F$ ,  $\text{partdiff}(f, z_1 + z_2) \text{ w.r.t. } 1 = \text{partdiff}(f, z_1) \text{ w.r.t. } 1 + \text{partdiff}(f, z_2) \text{ w.r.t. } 1$ , and
- (ii) for every point  $z$  of  $E \times F$  and for every real number  $a$ ,  $\text{partdiff}(f, a \cdot z) \text{ w.r.t. } 1 = a \cdot (\text{partdiff}(f, z) \text{ w.r.t. } 1)$ , and

- (iii) for every points  $z_1, z_2$  of  $E \times F$ ,  $\text{partdiff}(f, z_1 - z_2) \text{ w.r.t. } 1 = \text{partdiff}(f, z_1) \text{ w.r.t. } 1 - \text{partdiff}(f, z_2) \text{ w.r.t. } 1$ .

The theorem is a consequence of (4) and (5).

- (8) (i) for every points  $z_1, z_2$  of  $E \times F$ ,  $\text{partdiff}(f, z_1 + z_2) \text{ w.r.t. } 2 = \text{partdiff}(f, z_1) \text{ w.r.t. } 2 + \text{partdiff}(f, z_2) \text{ w.r.t. } 2$ , and  
 (ii) for every point  $z$  of  $E \times F$  and for every real number  $a$ ,  $\text{partdiff}(f, a \cdot z) \text{ w.r.t. } 2 = a \cdot (\text{partdiff}(f, z) \text{ w.r.t. } 2)$ , and  
 (iii) for every points  $z_1, z_2$  of  $E \times F$ ,  $\text{partdiff}(f, z_1 - z_2) \text{ w.r.t. } 2 = \text{partdiff}(f, z_1) \text{ w.r.t. } 2 - \text{partdiff}(f, z_2) \text{ w.r.t. } 2$ .

The theorem is a consequence of (4) and (5).

- (9) Let us consider a Lipschitzian bilinear operator  $f$  from  $E \times F$  into  $G$ , and a subset  $Z$  of  $E \times F$ . Suppose  $Z$  is open. Then  
 (i)  $f$  is partially differentiable on  $Z$  w.r.t. 1, and  
 (ii)  $f$  is partially differentiable on  $Z$  w.r.t. 2, and  
 (iii)  $f \upharpoonright^1 Z$  is continuous on  $Z$ , and  
 (iv)  $f \upharpoonright^2 Z$  is continuous on  $Z$ .

PROOF: For every point  $x$  of  $E \times F$  such that  $x \in Z$  holds  $f$  is partially differentiable in  $x$  w.r.t. 1. For every point  $x$  of  $E \times F$  such that  $x \in Z$  holds  $f$  is partially differentiable in  $x$  w.r.t. 2. Set  $g_1 = f \upharpoonright^1 Z$ . Set  $g_2 = f \upharpoonright^2 Z$ . Consider  $K$  being a real number such that  $0 \leq K$  and for every point  $z$  of  $E \times F$ ,  $\|\text{partdiff}(f, z) \text{ w.r.t. } 1\| \leq K \cdot \|z\|$  and  $\|\text{partdiff}(f, z) \text{ w.r.t. } 2\| \leq K \cdot \|z\|$ . For every point  $t_0$  of  $E \times F$  and for every real number  $r$  such that  $t_0 \in Z$  and  $0 < r$  there exists a real number  $s$  such that  $0 < s$  and for every point  $t_1$  of  $E \times F$  such that  $t_1 \in Z$  and  $\|t_1 - t_0\| < s$  holds  $\|g_{1/t_1} - g_{1/t_0}\| < r$ . For every point  $t_0$  of  $E \times F$  and for every real number  $r$  such that  $t_0 \in Z$  and  $0 < r$  there exists a real number  $s$  such that  $0 < s$  and for every point  $t_1$  of  $E \times F$  such that  $t_1 \in Z$  and  $\|t_1 - t_0\| < s$  holds  $\|g_{2/t_1} - g_{2/t_0}\| < r$ .  $\square$

Let us consider a Lipschitzian bilinear operator  $f$  from  $E \times F$  into  $G$ . Now we state the propositions:

- (10) There exists a real number  $K$  such that  
 (i)  $0 \leq K$ , and  
 (ii) for every point  $z$  of  $E \times F$ , there exists a Lipschitzian linear operator  $L$  from  $E \times F$  into  $G$  such that for every point  $d_1$  of  $E$  and for every point  $d_2$  of  $F$ ,  $L(d_1, d_2) = f(d_1, (z)_2) + f((z)_1, d_2)$  and for every point  $s$  of  $E \times F$ ,  $\|L(s)\| \leq K \cdot \|z\| \cdot \|s\|$ .

PROOF: Consider  $K$  being a real number such that  $0 \leq K$  and for every vector  $x$  of  $E$  and for every vector  $y$  of  $F$ ,  $\|f(x, y)\| \leq K \cdot \|x\| \cdot \|y\|$ . Define  $\mathcal{Q}(\text{element of } E, \text{element of } F) = f(\$1, (z)_2) + f((z)_1, \$2)$ . Consider  $L_0$  being a function from  $(\text{the carrier of } E) \times (\text{the carrier of } F)$  into the carrier of  $G$  such that for every element  $x$  of the carrier of  $E$  and for every element  $y$  of the carrier of  $F$ ,  $L_0(x, y) = \mathcal{Q}(x, y)$ . Reconsider  $L = L_0$  as a function from  $E \times F$  into  $G$ . For every elements  $x, y$  of  $E \times F$ ,  $L(x+y) = L(x) + L(y)$ . For every vector  $x$  of  $E \times F$  and for every real number  $a$ ,  $L(a \cdot x) = a \cdot L(x)$ . Set  $K_1 = 2 \cdot K \cdot \|z\|$ . For every vector  $w$  of  $E \times F$ ,  $\|L(w)\| \leq K_1 \cdot \|w\|$ .  $\square$

(11) There exists a real number  $K$  such that

- (i)  $0 \leq K$ , and
- (ii) for every point  $z$  of  $E \times F$ ,  $f$  is differentiable in  $z$  and for every point  $d_1$  of  $E$  and for every point  $d_2$  of  $F$ ,  $f'(z)(d_1, d_2) = f(d_1, (z)_2) + f((z)_1, d_2)$  and  $\|f'(z)\| \leq K \cdot \|z\|$ .

PROOF: Consider  $K_0$  being a real number such that  $0 \leq K_0$  and for every vector  $x$  of  $E$  and for every vector  $y$  of  $F$ ,  $\|f(x, y)\| \leq K_0 \cdot \|x\| \cdot \|y\|$ . Consider  $K$  being a real number such that  $0 \leq K$  and for every point  $z$  of  $E \times F$ , there exists a Lipschitzian linear operator  $L$  from  $E \times F$  into  $G$  such that for every point  $d_1$  of  $E$  for every point  $d_2$  of  $F$ ,  $L(d_1, d_2) = f(d_1, (z)_2) + f((z)_1, d_2)$  and for every point  $s$  of  $E \times F$ ,  $\|L(s)\| \leq K \cdot \|z\| \cdot \|s\|$ . Consider  $L$  being a Lipschitzian linear operator from  $E \times F$  into  $G$  such that for every point  $d_1$  of  $E$  and for every point  $d_2$  of  $F$ ,  $L(d_1, d_2) = f(d_1, (z)_2) + f((z)_1, d_2)$  and for every point  $s$  of  $E \times F$ ,  $\|L(s)\| \leq K \cdot \|z\| \cdot \|s\|$ . Reconsider  $L_0 = L$  as a point of the real norm space of bounded linear operators from  $E \times F$  into  $G$ .

Define  $\mathcal{Q}(\text{element of } E, \text{element of } F) = f(\$1, \$2)$ . Consider  $R$  being a function from  $(\text{the carrier of } E) \times (\text{the carrier of } F)$  into the carrier of  $G$  such that for every element  $d_1$  of the carrier of  $E$  and for every element  $d_2$  of the carrier of  $F$ ,  $R(d_1, d_2) = \mathcal{Q}(d_1, d_2)$ . For every real number  $r$  such that  $r > 0$  there exists a real number  $d$  such that  $d > 0$  and for every point  $w$  of  $E \times F$  such that  $w \neq 0_{E \times F}$  and  $\|w\| < d$  holds  $\|w\|^{-1} \cdot \|R_{/w}\| < r$ . For every point  $w$  of  $E \times F$  such that  $w \in$  the neighbourhood of  $z$  holds  $f_{/w} - f_{/z} = L_0(w - z) + R_{/w-z}$  by [10, (12)].  $\square$

(12) Let us consider a Lipschitzian bilinear operator  $f$  from  $E \times F$  into  $G$ , a point  $z$  of  $E \times F$ , a point  $d_1$  of  $E$ , and a point  $d_2$  of  $F$ . Then  $f'(z)(d_1, d_2) = (\text{partdiff}(f, z) \text{ w.r.t. } 1)(d_1) + (\text{partdiff}(f, z) \text{ w.r.t. } 2)(d_2)$ . The theorem is a consequence of (11) and (4).

(13) Let us consider a Lipschitzian bilinear operator  $f$  from  $E \times F$  into  $G$ . Then

- (i) for every points  $z_1, z_2$  of  $E \times F$ ,  $f'(z_1 + z_2) = f'(z_1) + f'(z_2)$ , and
- (ii) for every point  $z$  of  $E \times F$  and for every real number  $a$ ,  $f'(a \cdot z) = a \cdot f'(z)$ , and
- (iii) for every points  $z_1, z_2$  of  $E \times F$ ,  $f'(z_1 - z_2) = f'(z_1) - f'(z_2)$ .

The theorem is a consequence of (12), (7), and (8).

- (14) Let us consider a Lipschitzian bilinear operator  $f$  from  $E \times F$  into  $G$ , and a subset  $Z$  of  $E \times F$ . Suppose  $Z$  is open. Then

- (i)  $f$  is differentiable on  $Z$ , and
- (ii)  $f'|_Z$  is continuous on  $Z$ .

PROOF: Consider  $K$  being a real number such that  $0 \leq K$  and for every point  $z$  of  $E \times F$ ,  $f$  is differentiable in  $z$  and for every point  $d_1$  of  $E$  and for every point  $d_2$  of  $F$ ,  $f'(z)(d_1, d_2) = f(d_1, (z)_2) + f((z)_1, d_2)$  and  $\|f'(z)\| \leq K \cdot \|z\|$ . Set  $g_1 = f'|_Z$ . For every point  $t_0$  of  $E \times F$  and for every real number  $r$  such that  $t_0 \in Z$  and  $0 < r$  there exists a real number  $s$  such that  $0 < s$  and for every point  $t_1$  of  $E \times F$  such that  $t_1 \in Z$  and  $\|t_1 - t_0\| < s$  holds  $\|g_1|_{t_1} - g_1|_{t_0}\| < r$ .  $\square$

### 3. HIGHER-ORDER DERIVATIVES AND THEIR PROPERTIES

Now we state the propositions:

- (15) Let us consider a Lipschitzian bilinear operator  $f$  from  $E \times F$  into  $G$ . Then

- (i)  $f'|_{\Omega_{E \times F}}$  is Lipschitzian linear operator from  $E \times F$  into the real norm space of bounded linear operators from  $E \times F$  into  $G$ , differentiable on  $\Omega_{E \times F}$ , and continuous on the carrier of  $E \times F$ , and
- (ii) for every point  $z$  of  $E \times F$ ,  $(f'|_{\Omega_{E \times F}})'(z) = f'|_{\Omega_{E \times F}}$ .

The theorem is a consequence of (14), (13), and (11).

- (16) Let us consider a Lipschitzian linear operator  $L$  from  $E$  into  $F$ . Then

- (i)  $L'(\Omega_E)(0) = L$ , and
- (ii)  $L'(\Omega_E)(1) = \Omega_E \mapsto L$ , and
- (iii)  $L'(\Omega_E)(2) = \Omega_E \mapsto (\Omega_E \mapsto (\Omega_E \mapsto 0_F))$ , and
- (iv)  $L'(\Omega_E)(3) = \Omega_E \mapsto (\Omega_E \mapsto (\Omega_E \mapsto (\Omega_E \mapsto 0_F)))$ .

PROOF: For every object  $z$  such that  $z \in \text{dom } L'|_{\Omega_E}$  holds  $L'|_{\Omega_E}(z) = L$  by [5, (26)]. For every object  $z$  such that  $z \in \text{dom } (L'|_{\Omega_E})'|_{\Omega_E}$  holds  $(L'|_{\Omega_E})'|_{\Omega_E}(z) = \Omega_E \mapsto (\Omega_E \mapsto 0_F)$ . Reconsider  $L_1 = L'(\Omega_E)(2)$  as

a partial function from  $E$  to the real norm space of bounded linear operators from  $E$  into the real norm space of bounded linear operators from  $E$  into  $F$ . For every object  $z$  such that  $z \in \text{dom } L_1'_{|\Omega_E}$  holds  $L_1'_{|\Omega_E}(z) = \Omega_E \mapsto (\Omega_E \mapsto (\Omega_E \mapsto 0_F))$ .  $\square$

- (17) Let us consider a natural number  $i$ . Then  $0_{\text{diff}_{\text{SP}}(E^{(i+1)}, F)} = \Omega_E \mapsto 0_{\text{diff}_{\text{SP}}(E^i, F)}$ .

Let us consider a Lipschitzian linear operator  $L$  from  $E$  into  $F$  and a natural number  $i$ . Now we state the propositions:

- (18)  $\text{diff}_{\Omega_E}(L, i+2) = \Omega_E \mapsto 0_{\text{diff}_{\text{SP}}(E^{(i+2)}, F)}$ .

PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv \text{diff}_{\Omega_E}(L, \$_1 + 2) =$

$\Omega_E \mapsto 0_{\text{diff}_{\text{SP}}(E^{(\$_1+2)}, F)}$ .  $\mathcal{P}[0]$ . For every natural number  $i$  such that  $\mathcal{P}[i]$  holds  $\mathcal{P}[i+1]$ . For every natural number  $i$ ,  $\mathcal{P}[i]$ .  $\square$

- (19) (i)  $\text{diff}_{\Omega_E}(L, i+1)$  is differentiable on  $\Omega_E$ , and

(ii)  $\text{diff}_{\Omega_E}(L, i+1)'_{|\Omega_E} = \Omega_E \mapsto 0_{\text{diff}_{\text{SP}}(E^{(i+2)}, F)}$ , and

(iii)  $\text{diff}_{\Omega_E}(L, i+1)'_{|\Omega_E}$  is continuous on  $\Omega_E$ .

PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv \text{diff}_{\Omega_E}(L, \$_1 + 1)$  is differentiable on  $\Omega_E$  and  $\text{diff}_{\Omega_E}(L, \$_1 + 1)'_{|\Omega_E} = \Omega_E \mapsto 0_{\text{diff}_{\text{SP}}(E^{(\$_1+2)}, F)}$  and  $\text{diff}_{\Omega_E}(L, \$_1 + 1)'_{|\Omega_E}$  is continuous on  $\Omega_E$ .  $\mathcal{P}[0]$ . For every natural number  $i$  such that  $\mathcal{P}[i]$  holds  $\mathcal{P}[i+1]$ . For every natural number  $i$ ,  $\mathcal{P}[i]$ .  $\square$

- (20) (i)  $\text{diff}_{\Omega_E}(L, i)$  is differentiable on  $\Omega_E$ , and

(ii)  $\text{diff}_{\Omega_E}(L, i)'_{|\Omega_E}$  is continuous on  $\Omega_E$ .

The theorem is a consequence of (16) and (19).

- (21) Let us consider a Lipschitzian bilinear operator  $B$  from  $E \times F$  into  $G$ , and a natural number  $i$ . Then

(i)  $\text{diff}_{\Omega_{E \times F}}(B, i)$  is differentiable on  $\Omega_{E \times F}$ , and

(ii)  $\text{diff}_{\Omega_{E \times F}}(B, i)'_{|\Omega_{E \times F}}$  is continuous on  $\Omega_{E \times F}$ .

PROOF: Reconsider  $L = B'_{|\Omega_{E \times F}}$  as a Lipschitzian linear operator from  $E \times F$  into the real norm space of bounded linear operators from  $E \times F$  into  $G$ . Set  $G_1 =$  the real norm space of bounded linear operators from  $E \times F$  into  $G$ . Define  $\mathcal{P}[\text{natural number}] \equiv \text{diff}_{\Omega_{E \times F}}(B, \$_1 + 1) = \text{diff}_{\Omega_{E \times F}}(L, \$_1)$  and  $\text{diff}_{\text{SP}}((E \times F)^{(\$_1+1)}, G) = \text{diff}_{\text{SP}}((E \times F)^{\$1}, G_1)$ .  $\mathcal{P}[0]$  by [4, (11), (7)]. For every natural number  $n$  such that  $\mathcal{P}[n]$  holds  $\mathcal{P}[n+1]$  by [4, (10), (13)]. For every natural number  $n$ ,  $\mathcal{P}[n]$ .  $\square$

## REFERENCES

- [1] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Matuszewski, Adam Naumowicz, Karol Pąk, and Josef Urban. Mizar: State-of-the-art and

- beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8\_17.
- [2] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Matuszewski, Adam Naumowicz, and Karol Pąk. The role of the Mizar Mathematical Library for interactive proof development in Mizar. *Journal of Automated Reasoning*, 61(1):9–32, 2018. doi:10.1007/s10817-017-9440-6.
  - [3] Nelson Dunford and Jacob T. Schwartz. *Linear operators I*. Interscience Publ., 1958.
  - [4] Noboru Endou and Yasunari Shidama. Differentiation in normed spaces. *Formalized Mathematics*, 21(2):95–102, 2013. doi:10.2478/forma-2013-0011.
  - [5] Yuichi Futa, Noboru Endou, and Yasunari Shidama. Isometric differentiable functions on real normed space. *Formalized Mathematics*, 21(4):249–260, 2013. doi:10.2478/forma-2013-0027.
  - [6] Isao Miyadera. *Functional Analysis*. Riko-Gaku-Sya, 1972.
  - [7] Kazuhisa Nakasho. Bilinear operators on normed linear spaces. *Formalized Mathematics*, 27(1):15–23, 2019. doi:10.2478/forma-2019-0002.
  - [8] Kazuhisa Nakasho and Yuichi Futa. Isomorphism between spaces of multilinear maps and nested compositions over real normed vector spaces. *Formalized Mathematics*, 30(1):67–77, 2022. doi:10.2478/forma-2022-0006.
  - [9] Kazuhisa Nakasho and Yasunari Shidama. On implicit and inverse function theorems on Euclidean spaces. *Formalized Mathematics*, 30(3):159–168, 2022. doi:10.2478/forma-2022-0012.
  - [10] Kazuhisa Nakasho, Yuichi Futa, and Yasunari Shidama. Continuity of bounded linear operators on normed linear spaces. *Formalized Mathematics*, 26(3):231–237, 2018. doi:10.2478/forma-2018-0021.
  - [11] Laurent Schwartz. *Théorie des ensembles et topologie, tome 1. Analyse*. Hermann, 1997.
  - [12] Laurent Schwartz. *Calcul différentiel, tome 2. Analyse*. Hermann, 1997.
  - [13] Kôzaku Yosida. *Functional Analysis*. Springer, 1980.

*Accepted December 9, 2024*

---