

Fundamentals of Finitary Proofs¹

Taneli Huuskonen
Tampere, Finland

Summary. An abstract, generic textbook notion of a finitary proof and some of its basic properties are presented, using the Mizar system. A general form of Lindenbaum’s lemma is included.

MSC: 03B22 68V20

Keywords: finitary proof; proof system; Lindenbaum’s lemma

MML identifier: PROOFS.1, version: 8.1.14 5.87.1483

INTRODUCTION

An abstract, generic textbook notion of a finitary proof and some of its basic properties are presented, using the Mizar system [1], [2]. The approach is analogous to that of many textbooks, such as [11] or [3]. A general form of Lindenbaum’s lemma is included.

The outline of the paper is as follows: first three sections define formulas and rules, proof steps and derivability. Section 4 describes formally the behaviour of supersets of formulas and rules. Section 5 contains the key definition in this article: the structure defining proof systems (prefixed by **1-sorted**); one can notice that due to the set-theoretic approach claimed in the Mizar Mathematical Library, a binary relation can denote either a single, or more rules (hence a type is just a rule, but the selector in the structure has the name “rules”). Closing sections contain Lindenbaum’s and Teichmüller-Tukey lemmas.

Part of the contents were taken from [9], which was written to formalize some ideas from [4], [5], and [8]. This general approach could allow either to

¹This work has been supported by the National Science Centre of Poland, grant number UMO-2017/25/B/HS1/00503.

rewrite previous articles in this uniform language in the process of revisions [7], or to develop other logics, such as Suszko's logics ([12], [13], [14]), intuitionistic logic [10], or even fuzzy logics [6].

1. PRELIMINARIES: FORMULAS AND RULES

From now on i, j, k, l, m, n denote natural numbers, a, b, c, t, u denote objects, X, Y, Z denote sets, D, D_1, D_2, H denote non empty sets, and p, q, r, s denote finite sequences.

Let R be a binary relation. We say that R is finitary if and only if

(Def. 1) for every a such that $a \in \text{dom } R$ holds a is a finite set.

Let us observe that every binary relation which is empty is also finitary and there exists a binary relation which is finitary.

We introduce the notation formula as a synonym of object.

A rule is a finitary binary relation.

A formula-finset is a finite set.

A formula-sequence is a finite sequence. Let H be a set.

A rule of H is a rule defined by

(Def. 2) $\text{dom } it \subseteq \text{Fin } H$ and $\text{rng } it \subseteq H$.

Let H be a non empty set.

A formula of H is a formula defined by

(Def. 3) $it \in H$.

Let H be a set.

A formula-finset of H is a formula-finset defined by

(Def. 4) $it \subseteq H$.

A formula-sequence of H is a formula-sequence defined by

(Def. 5) it is a finite sequence of elements of H .

In the sequel R, R_1, R_2 denote rules, A, A_1, A_2 denote non empty sets, B, B_1, B_2 denote sets, P, P_1, P_2 denote formula-sequences, and S, S_1, S_2 denote formula-finsets.

Let us consider P . Observe that the functor $\text{rng } P$ yields a formula-finset. Let us consider H . Let B_1, B_2 be subsets of H . Note that the functor $B_1 \cup B_2$ yields a subset of H . Let us consider S_1 and S_2 . One can check that the functor $S_1 \cup S_2$ yields a formula-finset. Let us consider H . Let S_1, S_2 be formula-finsets of H . Let us note that the functor $S_1 \cup S_2$ yields a formula-finset of H . Let us consider R_1 and R_2 . Note that the functor $R_1 \cup R_2$ yields a rule. Let us consider H . Let R_1, R_2 be rules of H . Let us observe that the functor $R_1 \cup R_2$ yields a rule of H .

2. PROOF STEPS

Let us consider B , R , P , and n . We say that (P, n) is a correct step w.r.t. B , R if and only if

- (Def. 6) $P(n) \in B$ or there exists a formula-finset Q such that $\langle Q, P(n) \rangle \in R$ and for every t such that $t \in Q$ there exists k such that $k \in \text{dom } P$ and $k < n$ and $P(k) = t$.

We say that P is (B, R) -correct if and only if

- (Def. 7) for every k such that $k \in \text{dom } P$ holds (P, k) is a correct step w.r.t. B , R .

Let us observe that every formula-sequence which is non (B, R) -correct is also non empty.

Let us consider H . Let us observe that there exists a formula-sequence of H which is (B, R) -correct and there exists a formula-sequence which is (B, R) -correct. Now we state the proposition:

- (1) Let us consider an element a of A . Then $\langle a \rangle$ is (A, R) -correct.

Let us consider A and R . Let us observe that there exists a formula-sequence which is non empty and (A, R) -correct.

3. DERIVABILITY

Let us consider B , R , and S . We say that S is (B, R) -derivable if and only if

- (Def. 8) there exists P such that $S = \text{rng } P$ and P is (B, R) -correct.

Now we state the propositions:

- (2) If P is (B, R) -correct and $P = P_1 \frown P_2$, then P_1 is (B, R) -correct.
 (3) If P_1 is (B, R) -correct and P_2 is (B, R) -correct, then $P_1 \frown P_2$ is (B, R) -correct.
 (4) If S_1 is (B, R) -derivable and S_2 is (B, R) -derivable, then $S_1 \cup S_2$ is (B, R) -derivable. The theorem is a consequence of (3).
 (5) If $B \subseteq B_1$ and $R \subseteq R_1$ and P is (B, R) -correct, then P is (B_1, R_1) -correct.

Let us consider B and a . We say that a is B -axiomatic if and only if

- (Def. 9) $a \in B$.

Let us consider R . We say that $B, R \vdash a$ if and only if

- (Def. 10) there exists P such that $a \in \text{rng } P$ and P is (B, R) -correct.

We say that a is (B, R) -provable if and only if

(Def. 11) $B, R \vdash a$.

4. EXTENSIONS

Let us consider B and B_1 . We say that B_1 is B -extending if and only if

(Def. 12) $B \subseteq B_1$.

Let us consider R and R_1 . We say that R_1 is R -extending if and only if

(Def. 13) $R \subseteq R_1$.

Let us consider B . Observe that there exists a set which is B -extending. Let us consider R . Let us observe that there exists a rule which is R -extending. Let us consider B .

An extension of B is a B -extending set. Let us consider H . Let B be a subset of H . Note that there exists a subset of H which is B -extending.

An extension of B is a B -extending subset of H . Let us consider R .

An extension of R is an R -extending rule. Let us consider H . Let B be a subset of H and t be a formula of H . The functor $B + t$ yielding an extension of B is defined by the term

(Def. 14) $B \cup \{t\}$.

Now we state the proposition:

(6) a is $(B \cup \{t\})$ -axiomatic if and only if a is B -axiomatic or $a = t$.

From now on C denotes an extension of B and E denotes an extension of R .

Let us consider B and C . Let us note that every set which is C -extending is also B -extending and every object which is non C -axiomatic is also non B -axiomatic.

Let us consider R and E . Let us note that every rule which is E -extending is also R -extending.

Let us consider B and R_1 . We say that R_1 is (B, R) -derivable if and only if

(Def. 15) for every S and t such that $\langle S, t \rangle \in R_1$ holds $B \cup S, R \vdash t$.

Now we state the propositions:

(7) $B, R \vdash t$ if and only if there exists S such that $t \in S$ and S is (B, R) -derivable.

(8) If $a \in B$, then $B, R \vdash a$. The theorem is a consequence of (1).

Let us consider B and R . One can verify that every object which is non (B, R) -provable is also non B -axiomatic. Now we state the propositions:

(9) If for every a such that $a \in S$ holds $B, R \vdash a$, then there exists S_1 such that $S \subseteq S_1$ and S_1 is (B, R) -derivable.

PROOF: Define $\mathcal{X}[\text{set}] \equiv$ there exists S_1 such that $\$1 \subseteq S_1$ and S_1 is (B, R) -derivable. $\mathcal{X}[\emptyset]$. For every sets x , B_1 such that $x \in S$ and $B_1 \subseteq S$ and $\mathcal{X}[B_1]$ holds $\mathcal{X}[B_1 \cup \{x\}]$. $\mathcal{X}[S]$. $\square$

- (10) If S is (B, R) -derivable and $B \cap S \subseteq B_1$, then S is (B_1, R) -derivable.

PROOF: Consider P such that $S = \text{rng } P$ and P is (B, R) -correct. P is (B_1, R) -correct. $\square$

- (11) If for every a such that $a \in S$ holds $B, R \vdash a$ and $\langle S, t \rangle \in R$, then $B, R \vdash t$. The theorem is a consequence of (9).

- (12) If $B, R \vdash a$, then $a \in B$ or there exists S such that $\langle S, a \rangle \in R$ and for every b such that $b \in S$ holds $B, R \vdash b$.

- (13) If S_1 is (B, R) -derivable and S_2 is (S_1, R) -derivable, then $S_1 \cup S_2$ is (B, R) -derivable.

PROOF: Consider P_1, P_2 such that P_1 is (B, R) -correct and $S_1 = \text{rng } P_1$ and P_2 is (S_1, R) -correct and $S_2 = \text{rng } P_2$. Set $P = P_1 \cap P_2$. For every k such that $k \in \text{dom } P_1$ holds (P, k) is a correct step w.r.t. B, R . P is (B, R) -correct. $\square$

- (14) If $B_1, R \vdash a$ and for every b such that $b \in B_1$ holds $B, R \vdash b$, then $B, R \vdash a$. The theorem is a consequence of (7), (9), (10), and (13).

- (15) If $B, R \vdash a$, then $C, E \vdash a$. The theorem is a consequence of (5).

Let us consider B, R , and a . Note that a is (B, R) -provable if and only if the condition (Def. 16) is satisfied.

(Def. 16) for every C and E , $C, E \vdash a$.

Let us consider C . Note that every object which is non (C, R) -provable is also non (B, R) -provable. Let us consider E . Observe that every object which is non (C, E) -provable is also non (B, R) -provable. Now we state the propositions:

- (16) $R_1 \cup R_2$ is (B, R) -derivable if and only if R_1 is (B, R) -derivable and R_2 is (B, R) -derivable.

- (17) Let us consider a subset B of H , a rule R of H , and a . If $B, R \vdash a$, then $a \in H$.

5. PROOF SYSTEMS

We consider proof systems which extend 1-sorted structures and are systems

$\langle \text{a carrier, axioms, rules} \rangle$

where the carrier is a set, the axioms constitute a subset of the carrier, the rules constitute a rule of the carrier.

Let P be a proof system. A formula-finset of P is a formula-finset of the carrier of P . Let a be an object. We say that $P \vdash a$ if and only if

(Def. 17) the axioms of P , the rules of $P \vdash a$.

Note that there exists a proof system which is non empty.

From now on P denotes a non empty proof system, B, B_1, B_2 denote subsets of P , and F denotes a finite subset of P . Now we state the proposition:

(18) If $P \vdash a$, then a is an element of P .

Let us consider P and B . We say that $P \vdash B$ if and only if

(Def. 18) for every a such that $a \in B$ holds $P \vdash a$.

Let us consider B_1 and B_2 . One can check that the functor $B_1 \cup B_2$ yields a subset of P .

6. CONSISTENCY

Let us consider P . We say that P is consistent if and only if

(Def. 19) there exists a such that $a \in P$ and $P \not\vdash a$.

Let us consider B . The functor $P \cup B$ yielding a non empty proof system is defined by the term

(Def. 20) $\langle \text{the carrier of } P, (\text{the axioms of } P) \cup B, \text{the rules of } P \rangle$.

Let us note that there exists a non empty proof system which is consistent and strict. Let P be a strict proof system and E be an empty subset of P . Let us observe that $P \cup E$ reduces to P . Let us consider P . We introduce the notation P is inconsistent as an antonym for P is consistent.

Let us consider B . We say that B is consistent if and only if

(Def. 21) $P \cup B$ is consistent.

Let P be a consistent, non empty proof system. Note that there exists a subset of P which is consistent.

Let us consider P and B . We introduce the notation B is inconsistent as an antonym for B is consistent.

One can check that there exists a subset of P which is inconsistent.

We say that P is paraconsistent if and only if

(Def. 22) every finite subset of P is consistent.

One can verify that every non empty proof system which is paraconsistent is also consistent and there exists a non empty proof system which is consistent and non paraconsistent.

7. CONTRADICTIONS AND LINDENBAUM'S LEMMA

Let us consider P , B , and B_1 . We say that B_1 is B -omitting if and only if
 (Def. 23) there exists a such that $a \in B$ and $P \cup B_1 \not\models a$.

Now we state the proposition:

(19) If B is inconsistent, then B_1 is consistent iff B_1 is B -omitting. The theorem is a consequence of (8) and (14).

Let us consider P . Let B be an inconsistent subset of P . One can verify that every subset of P which is B -omitting is also consistent and every subset of P which is non B -omitting is also inconsistent.

Let us consider B . One can verify that there exists a subset of P which is non B -omitting. Now we state the proposition:

(20) If B_1 is B -omitting and $B_2 \subseteq B_1$, then B_2 is B -omitting. The theorem is a consequence of (15).

Let us consider P and B . The functor $\text{Omit}(P, B)$ yielding a family of subsets of P is defined by the term

(Def. 24) $\{B_1, \text{ where } B_1 \text{ is a subset of } P : B_1 \text{ is } B\text{-omitting}\}$.

One can verify that the functor $\text{Omit}(P, B)$ is defined by

(Def. 25) for every B_1 , $B_1 \in \text{it}$ iff B_1 is B -omitting.

Let us consider B_1 . We say that B_1 is B -maximally-omitting if and only if
 (Def. 26) B_1 is B -omitting and for every B_2 such that $B_1 \subset B_2$ holds B_2 is not B -omitting.

Observe that every subset of P which is B -maximally-omitting is also B -omitting.

Let us consider X . We say that X is finite-character if and only if

(Def. 27) for every a , $a \in X$ iff there exists a set B such that $B = a$ and for every finite subset S of B , $S \in X$.

Let us observe that X is finite-character if and only if the condition (Def. 28) is satisfied.

(Def. 28) for every Y , $Y \in X$ iff for every finite subset S of Y , $S \in X$.

Let F be a family of subsets of X . Observe that F is finite-character if and only if the condition (Def. 29) is satisfied.

(Def. 29) for every subset B of X , $B \in F$ iff for every finite subset S of B , $S \in F$.

One can check that there exists a family of subsets of X which is non empty and finite-character and every set which is empty is also finite-character and there exists a set which is non empty and finite-character.

8. TEICHMÜLLER-TUKEY LEMMA

Now we state the proposition:

- (21) Let us consider a non empty, finite-character set X . Then there exists an element Y of X such that for every element Z of X , $Y \not\subseteq Z$.

PROOF: For every set C such that $C \subseteq X$ and C is $\subseteq$ -linear there exists Y such that $Y \in C$ and for every Z such that $Z \in C$ holds $Z \subseteq Y$. Consider Y such that $Y \in X$ and for every Z such that $Z \in X$ and $Z \neq Y$ holds $Y \not\subseteq Z$. $\square$

Let us consider P and F . One can check that $\text{Omit}(P, F)$ is finite-character. Now we state the proposition:

- (22) If B is F -omitting, then there exists B_1 such that $B \subseteq B_1$ and B_1 is F -maximally-omitting. The theorem is a consequence of (21).

Let us consider P and B . We say that B is maximally-consistent if and only if

- (Def. 30) B is consistent and for every B_1 such that $B \subset B_1$ holds B_1 is inconsistent.

Now we state the proposition:

- (23) If P is consistent and non paraconsistent and B is consistent, then there exists B_1 such that $B \subseteq B_1$ and B_1 is maximally-consistent. The theorem is a consequence of (22).

The scheme *UnOpCongr* deals with a non empty set $\mathcal{X}$ and a unary functor $\mathcal{F}$ yielding an element of $\mathcal{X}$ and an equivalence relation $\mathcal{E}$ of $\mathcal{X}$ and states that

- (Sch. 1) There exists a unary operation f on Classes $\mathcal{E}$ such that for every element x of $\mathcal{X}$, $f([x]_{\mathcal{E}}) = [\mathcal{F}(x)]_{\mathcal{E}}$
provided

- for every elements x, y of $\mathcal{X}$ such that $\langle x, y \rangle \in \mathcal{E}$ holds $\langle \mathcal{F}(x), \mathcal{F}(y) \rangle \in \mathcal{E}$.

The scheme *BinOpCongr* deals with a non empty set $\mathcal{X}$ and a binary functor $\mathcal{F}$ yielding an element of $\mathcal{X}$ and an equivalence relation $\mathcal{E}$ of $\mathcal{X}$ and states that

- (Sch. 2) There exists a binary operation f on Classes $\mathcal{E}$ such that for every elements x, y of $\mathcal{X}$, $f([x]_{\mathcal{E}}, [y]_{\mathcal{E}}) = [\mathcal{F}(x, y)]_{\mathcal{E}}$
provided

- for every elements x_1, x_2, y_1, y_2 of $\mathcal{X}$ such that $\langle x_1, x_2 \rangle, \langle y_1, y_2 \rangle \in \mathcal{E}$ holds $\langle \mathcal{F}(x_1, y_1), \mathcal{F}(x_2, y_2) \rangle \in \mathcal{E}$.

The scheme *ProofInduction* deals with a set $\mathcal{B}$ and a rule $\mathcal{R}$ and a unary predicate $\mathcal{S}$ and states that

(Sch. 3) For every a such that $\mathcal{B}, \mathcal{R} \vdash a$ holds $\mathcal{S}[a]$
provided

- for every a such that $a \in \mathcal{B}$ holds $\mathcal{S}[a]$ and
- for every X and a such that $\langle X, a \rangle \in \mathcal{R}$ and for every b such that $b \in X$ holds $\mathcal{S}[b]$ holds $\mathcal{S}[a]$.

Let us consider R and X . We say that X is R -closed if and only if

(Def. 31) for every Y and a such that $\langle Y, a \rangle \in R$ and $Y \subseteq X$ holds $a \in X$.

9. THEOREMS

Let us consider D and R . A theorem of D, R is an object defined by

(Def. 32) $D, R \vdash it$.

Note that the type possesses the sethood property. Let us consider X . The functor $\text{Theorems}(X, R)$ yielding a set is defined by the term

(Def. 33) $\{t, \text{ where } t \text{ is an element of } X \cup \text{rng } R : X, R \vdash t\}$.

Note that the functor $\text{Theorems}(X, R)$ is defined by

(Def. 34) for every a , $a \in it$ iff $X, R \vdash a$.

Note that $\text{Theorems}(X, R)$ is X -extending and R -closed and there exists a set which is X -extending and R -closed. Now we state the proposition:

(24) $X, R \vdash a$ if and only if for every R -closed, X -extending set Y , $a \in Y$.

Let us consider P . The functor $\text{Theorems}(P)$ yielding a subset of P is defined by the term

(Def. 35) $\text{Theorems}((\text{the axioms of } P), (\text{the rules of } P))$.

Let us consider X . We say that X is P -closed if and only if

(Def. 36) X is $(\text{the rules of } P)$ -closed and $(\text{the axioms of } P)$ -extending.

Let us note that $\text{Theorems}(P)$ is P -closed and every set which is P -closed is also $(\text{the rules of } P)$ -closed and $(\text{the axioms of } P)$ -extending and every set which is $(\text{the rules of } P)$ -closed and $(\text{the axioms of } P)$ -extending is also P -closed and there exists a subset of P which is P -closed and there exists a set which is P -closed. Now we state the proposition:

(25) $P \vdash a$ if and only if for every P -closed set X , $a \in X$. The theorem is a consequence of (24).

REFERENCES

- [1] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Matuszewski, Adam Naumowicz, Karol Pąk, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8_17.
- [2] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Matuszewski, Adam Naumowicz, and Karol Pąk. The role of the Mizar Mathematical Library for interactive proof development in Mizar. *Journal of Automated Reasoning*, 61(1):9–32, 2018. doi:10.1007/s10817-017-9440-6.
- [3] Heinz-Dieter Ebbinghaus, Jörg Flum, and Wolfgang Thomas. *Mathematical Logic*. Springer, 1994.
- [4] Joanna Golińska-Pilarek and Taneli Huuskonen. Logic of descriptions. A new approach to the foundations of mathematics and science. *Studies in Logic, Grammar and Rhetoric*, 40(27), 2012.
- [5] Joanna Golińska-Pilarek and Taneli Huuskonen. Grzegorzczyk’s non-Fregean logics. In Rafał Urbaniak and Gillman Payette, editors, *Applications of Formal Philosophy: The Road Less Travelled*, Logic, Reasoning and Argumentation. Springer, 2015.
- [6] Adam Grabowski. On fuzzy negations and laws of contraposition. Lattice of fuzzy negations. *Formalized Mathematics*, 31(1):151–159, 2023. doi:10.2478/forma-2023-0014.
- [7] Adam Grabowski and Christoph Schwarzweller. Revisions as an essential tool to maintain mathematical repositories. In M. Kauers, M. Kerber, R. Miner, and W. Windsteiger, editors, *Towards Mechanized Mathematical Assistants. Lecture Notes in Computer Science*, volume 4573, pages 235–249. Springer: Berlin, Heidelberg, 2007.
- [8] Andrzej Grzegorzczak. Filozofia logiki i formalna LOGIKA NIESYMPLIKACYJNA. *Zagadnienia Naukoznawstwa*, XLVII(4), 2012. In Polish.
- [9] Taneli Huuskonen. Grzegorzczyk’s logics. Part I. *Formalized Mathematics*, 23(3):177–187, 2015. doi:10.1515/forma-2015-0015.
- [10] Takao Inoué and Riku Hanaoka. Intuitionistic Propositional Calculus in the extended framework with modal operator. Part II. *Formalized Mathematics*, 30(1):1–12, 2022. doi:10.2478/forma-2022-0001.
- [11] Elliott Mendelson. *Introduction to Mathematical Logic*. Chapman Hall/CRC, 1997. <http://books.google.pl/books?id=Z01p4QGspoYC>.
- [12] Roman Suszko. Non-Fregean logic and theories. *Analele Universitatii Bucuresti. Acta Logica*, 9:105–125, 1968.
- [13] Roman Suszko. Semantics for the sentential calculus with identity. *Studia Logica*, 28: 77–81, 1971.
- [14] Roman Suszko. Abolition of the Fregean axiom. In R. Parikh, editor, *Logic Colloquium: Symposium on Logic held at Boston, 1972–73*, volume 453 of *Lecture Notes in Mathematics*, pages 169–239, Heidelberg, 1975. Springer.

Accepted December 24, 2024
