

REVIEW

Serum Albumin Concentration and the Risk of Cardiovascular Disease and Acute Coronary Syndrome — a Narrative Review

Maria Czinege¹, Florina Ruța^{2*}, Victoria Nyulas³, Vasile-Bogdan Halațiu^{4,5}, Tiberiu Nyulas^{4,5}, Theodora Benedek⁶

¹ Department of Doctoral School, George Emil Palade University of Medicine, Pharmacy, Science and Technology, Târgu Mureș, Romania

² Department of Community Nutrition and Food Hygiene, George Emil Palade University of Medicine, Pharmacy, Science and Technology, Târgu Mureș, Romania

³ Department of Informatics and Medical Biostatistics, George Emil Palade University of Medicine, Pharmacy, Science and Technology, Târgu Mureș, Romania

⁴ Department of Physiology, George Emil Palade University of Medicine, Pharmacy, Science and Technology, Târgu Mureș, Romania

⁵ Clinic of Cardiology, County Emergency Clinical Hospital, Târgu Mureș, Romania

⁶ Department of Cardiology, George Emil Palade University of Medicine, Pharmacy, Science and Technology, Târgu Mureș, Romania

ABSTRACT

Cardiovascular diseases are the primary cause of global health decline. Conditions that cause inflammation and excessive oxidation are present in the early development of the majority of cardiovascular diseases, including coronary artery disease and heart failure. Among the many physiological functions of albumin, its antioxidant, anti-inflammatory, anticoagulant, and anti-aggregant properties are the most important. Low serum albumin levels are associated with the occurrence of various cardiovascular diseases such as coronary artery disease, stroke, heart failure, and atrial fibrillation. Several recent meta-analyses support the idea that hypoalbuminemia may act as a modifiable risk factor that increases oxidative stress and inflammation, contributing to the development and progression of cardiovascular diseases. However, it is unclear whether correcting hypoalbuminemia can improve outcomes for patients with cardiovascular diseases and reduce mortality rates. In this regard, further research is crucial.

Keywords: acute coronary syndrome, serum albumin, prognosis, inflammation

ARTICLE HISTORY

Received: August 27, 2024

Accepted: September 20, 2024

CORRESPONDENCE

Florina Ruța

Str. Gheorghe Marinescu nr. 50
540142 Târgu Mureș, Romania
Email: florina.ruta@umfst.ro

INTRODUCTION

Cardiovascular disease is the primary cause of death in industrialized nations.^{1,2} Strong evidence from longitudinal research suggests that lower serum albumin levels are associated with an increased risk of heart failure and other cardiovascular diseases.³ Albumin, a protein that consti-

tutes over half of the body's serum proteins, has a crucial role in various physiological functions. Multiple clinical studies have shown that hypoalbuminemia is a strong predictor in a wide range of diseases.^{4,5}

Low serum albumin levels are associated with the development of cardiovascular conditions such as coronary heart disease, stroke, heart failure, and atrial fibrilla-

tion.⁵ Moreover, hypoalbuminemia has become a highly significant predictor in these diseases, independent of other prognostic markers.⁶ Its predictive value reflects both the severity of comorbidities and the likelihood of malnutrition–inflammation.^{5,7} In cases of acute coronary syndrome, hypoalbuminemia has been associated with increased disease severity, long-term mortality, and progression toward heart failure.⁸

The studies included in this review were identified through a comprehensive search of PubMed, Medline, Scopus, Cochrane Library, Google Academics and Web of Science databases. To refine the selection of relevant articles, we used the following keywords: “hypopalbuminemia”, “inflammation”, “cardiovascular disease”, “prognosis”, “heart failure”, “stroke”, “albumin”, “arrhythmias”, and “atrial fibrillation”.

After a thorough review of the literature, we identified 285 articles, of which 60 were removed due to duplication. A further 189 studies were excluded based on insufficient data, variations in interventions, outcomes, or an inadequate sample size. Of the 36 remaining studies, only 6 met all inclusion criteria and were included in the qualitative analysis.

PHYSIOLOGICAL AND BIOCHEMICAL PROPERTIES OF SERUM ALBUMIN

Albumin is synthesized in the liver in approximately 17 days.⁵ There are continual schisms between intravascular and interstitial compartments, with more than one-third of the total albumin found in the intravascular compartment.^{5,9} Several factors promote albumin production, including reduced colloid pressure osmosis, insulin, and amino acid intake. Conversely, high colloid osmotic pressure, starvation, inflammation, diabetes, liver disease, and sepsis can reduce its synthesis.⁵ The normal serum albumin level ranges from 3.5 to 5.0 g/dl, although the threshold for concerning levels may vary depending on the analytical method used.⁵ Women typically have slightly lower levels than men, and these levels gradually decrease with age. Albumin has several important physiological roles, including particle and drug transport, anti-inflammatory, antioxidant, and antiplatelet activity, as well as regulating fluid exchange between intravascular and interstitial spaces.^{5,10}

In addition to transporting a wide range of endogenous and exogenous substances, such as ions, bilirubin, vitamin B12, hormones, steroids, and drugs,⁵ albumin also has anti-inflammatory properties. Although not fully understood, one key mechanism involves its ability to prevent

monocytes from adhering to endothelial cells, thereby reducing inflammation and atherosclerosis.^{10–12}

In an experimental rat model of endotoxemia, albumin reduced both the messenger RNA and protein expression of cardiac nitric oxide synthase II, which regulates nitric oxide production and myocardial inflammation.¹⁰ Serum albumin is considered the most important antioxidant in the circulatory system.^{11,12} At normal concentrations, albumin prevents the accumulation of histones in arterial plaques; however, a decrease in serum albumin levels reduces its ability to prevent plaque formation. Additionally, biological data indicate that serum albumin has a key role in anticoagulation.^{5,13}

HYPOALBUMINEMIA IN ACUTE CORONARY SYNDROME

The prevalence of hypoalbuminemia in patients with acute coronary syndrome remains poorly defined, with reported rates varying across studies, being influenced by age, disease severity, and the acuity of the underlying condition.⁵ The most common parameters associated with hypoalbuminemia in acute coronary syndrome include advanced age, anemia, inflammatory mediators, and low cholesterol.^{5,14,15}

Hypoalbuminemia occurs in approximately 14% of patients with congenital cardiac disease and is independently linked to functional class, liver failure, and renal impairment.⁵ In addition, both the Framingham Offspring Study and the Multiple Risk Factor Intervention Trial have identified a significant inverse relationship between serum albumin levels and the risk of coronary heart disease.¹⁶

A recent retrospective study by Polat *et al.* involving 403 patients with unstable angina pectoris or non-ST elevation myocardial infarction (NSTEMI) found that those with hypoalbuminemia were at a higher risk of long-term mortality than those without hypoalbuminemia. Based on these findings, the authors considered albumin to be a simple and easily measurable biomarker for predicting mortality risk in patients with NSTEMI.¹⁷

THE PREDICTIVE VALUE OF HYPOALBUMINEMIA IN CARDIOVASCULAR DISEASES

Hypoalbuminemia is a strong independent predictor for all cardiovascular diseases and their causes.^{18,19} In a study by Chien *et al.*, patients with low albumin levels had significantly higher risk of all-cause mortality than those with normal albumin levels (10.2% vs. 0.5%; $p <$

0.001).¹⁹ In a study involving 546 elderly patients admitted for acute non-ischemic heart failure, Ancion *et al.* found that a serum albumin level of ≤ 34 g/L was an independent predictor of hospital mortality.²⁰ Another study examining the altered nutritional status, inflammation, and systemic vulnerability of patients with myocardial infarction after percutaneous coronary revascularization revealed that patients with malnutrition had significantly higher serum urea levels and significantly lower total cholesterol and albumin levels than patients with normal nutritional status, calculated in accordance with the CONUT guidelines.^{21,22}

The predictive value of hypoalbuminemia has been well-established in patients with heart failure, and it has emerged as a time-independent predictor of worsening outcomes in both acute and chronic heart failure.^{23,24} This suggests that hypoalbuminemia may play a role in the progression of these conditions. According to Starling's law, severe hypoalbuminemia is associated with pulmonary congestion, which can destabilize both latent and active heart failure. Albuminuric acid maintains the hydrostatic balance of the heart through its osmotic action.²⁵ Furthermore, because serum albumin has antioxidant properties, hypoalbuminemia may aggravate oxidative stress and accelerate the progression of heart failure.^{26,27}

In the Atherosclerosis Risk in Communities (ARIC) study, which followed 14,506 individuals, a significant correlation was found between serum albumin levels and the incidence of ischemic cardiopathy in smokers, independent of other known risk factors.^{5,28} This suggests an interdependent relationship between albumin levels and smoking. The median albumin level, adjusted for age and sex, was significantly lower in participants who experienced a cardiovascular event compared to those who did not ($p < 0.0001$).²⁸

Another prospective study, which followed 2,986 participants over a period of 12 years, identified low albumin levels as a predictor of first stroke, alongside factors such as body mass index and inflammatory status.^{29,30} In a prospective study by Arques *et al.*, involving 52 elderly patients with acute diastolic heart failure, hypoalbuminemia was found to be the strongest predictor of in-hospital mortality (odds ratio, 0.57; $p < 0.005$) after adjustments for age and type B natriuretic peptide levels.³¹

Serum albumin plays a major role in the prognosis of cardiovascular disease, influencing inflammation, malnutrition, and the severity of comorbidities. This raises the question of whether it could be an unrecognized, potentially modifiable risk factor. Recent meta-analyses examin-

ing the role of low serum albumin levels in the onset of cardiovascular disease, heart failure, and acute coronary syndrome have provided compelling evidence in favor of this theory.⁵ The potential impact of low serum albumin levels on the development and progression of cardiovascular diseases is likely related to the anti-inflammatory, antioxidant, anticoagulant, and antiaggregative properties of serum albumin, particularly in the case of coronary artery disease, heart failure, atrial fibrillation, stroke, and venous thromboembolism.^{30,32–36} Additionally, there is evidence that hypoalbuminemia contributes to pulmonary edema and sodium retention, both of which are key factors in the progression of heart failure.³⁷

In a study by González-Pacheco *et al.* involving 8,848 patients with acute coronary syndrome, mortality rates were highest in the group with the lowest serum albumin levels and lowest in the group with the highest levels, with a significant difference ($p < 0.0001$) observed across all groups. The main finding of the study was that a serum albumin level below 3.5 g/dl at admission was an independent predictor of new-onset heart failure and in-hospital mortality. The authors also concluded that, in this particular clinical situation, the inflammatory state may be a mechanism underlying hypoalbuminemia.³⁹

Another study, involving 2,886 patients with congenital heart disease, showed that the prevalence of hypoalbuminemia was significantly higher in patients with congenital heart disease of high complexity (18.2%) compared to patients with mild disease (11.3 %) or uncomplicated disease (12.1 %; $p < 0.001$). In the univariable regression analysis, hypoalbuminemia was a strong predictor of the outcome (hazard ratio, 3.37; $p < 0.0001$) and also remained a significant predictor of decline in the multivariable regression analysis.⁴⁰

In a study involving 8,750 patients with acute myocardial infarction, Plakht *et al.* found that serum albumin levels were an independent marker for all-cause mortality in hospitalized survivors.¹⁴ In a study that analyzed 2,794 patients who underwent isolated coronary artery bypass grafting, Bhamidipati *et al.* found that the patients with the highest adjusted mortality rate had serum albumin levels of 2–3 g/dl ($p = 0.02$), the lowest body mass index ($p < 0.05$), lower LVEF, and higher intra-aortic balloon pump use ($p < 0.001$). In addition, an increase in albuminemia was associated with a lower rate of severe complications ($p = 0.001$) and a lower risk of mortality (adjusted odds ratio, 0.61; 95% CI 0.42–0.90). The study found that albumin is more closely linked to mortality and morbidity than the body mass index in patients undergoing coronary artery bypass grafting.⁴¹

ALBUMIN AND INFLAMMATION

Albumin is a major mediator of coronary ischemia-reperfusion injury and a significant inhibitor of plaque formation and aggregation. Fibrinogen, a plasma protein involved in inflammatory and thrombotic processes, can also serve as a marker for thrombotic states⁴² and predict cardiovascular events in the general population.⁴³ Additionally, a recent study has shown a correlation between elevated fibrinogen levels and the presence and severity of coronary heart disease.⁴⁴

Albumin is not just a basal protein responsible for maintaining the oncotic pressure of plasma; it is also a transporter of various substances and is involved in both acute and chronic inflammatory processes. Hypoalbuminemia can result from malnutrition, malabsorption, increased renal loss, decreased hepatic function, acute or chronic inflammation processes, and increased plasma volume. Inflammatory processes have been linked to an increase in albumin catabolization.⁴⁵ In a recent study, Çetin *et al.* found that patients with slow coronary flow had lower albumin levels than those with normal coronary arteries.⁴⁶

Total albumin reserves are 4–5 g/kg body weight, of which 40–45% are found in the intravascular space and the remaining 55–60% in the interstitial space.⁴⁷ Due to its relatively low molecular weight compared to other major intravascular proteins, such as immunoglobulins, albumin contributes to 75–80% of the body's colloid osmotic pressure. Another important function of albumin is its ability to bind to various ligands, including free fatty acids, calcium, steroid hormone analogues, bilirubin, thyroxine, and tryptophan.⁴⁷

Given that albumin is a negative acute-phase protein, its serum concentration is influenced by both inflammation and nutritional status.⁴⁷ Inflammation alters the structure of lipoproteins, contributing to the morbidity and mortality associated with hypoalbuminemia.⁴⁷

ALBUMIN AND HEART FAILURE

Heart failure is becoming one of the most important cardiovascular diseases owing to its increasing prevalence, high mortality, and rising costs of care. The inflammatory process that accompanies heart failure is an important prothrombotic factor.⁴⁸

Hypoalbuminemia is common in patients with heart failure and appears to be a reliable indicator of mortality and morbidity among individuals with low ejection fraction (<40%).^{49,50} In addition, it has been observed that

hypoalbuminemia is significantly associated with the incidence of heart failure in elderly patients.⁵⁰

Studies have shown that hypoalbuminemia occurs both in heart failure with impaired and preserved function.^{51–53} In acute heart failure, it is a reliable predictor of mortality.⁵³ The Cardiovascular Health Study, which included 5,450 healthy individuals aged 65 and older, found that hypoalbuminemia was significantly associated with the risk of developing heart failure within 10 years.⁵⁰ The causes of hypoalbuminemia have not been specifically studied in patients with heart failure; nevertheless, it may result from malnutrition and hepatic dysfunction.⁵² It has to be noted that systemic inflammation can lower serum albumin concentrations without causing malnutrition.⁵⁴

The inflammatory response results in significant changes to plasmatic, cytokine, and complement proteins. Serum albumin levels decrease in response to inflammation, being common in clinical scenarios linked to oxidative stress and chronic inflammation.^{55–57} According to Ancion *et al.*, hypoalbuminemia is common in hospitalized patients with acute heart failure and is linked to a higher risk of in-hospital mortality.⁵⁸ In addition to traditional risk markers such as age, hemodynamic and biological parameters (leukocyte count), low serum albumin levels can also predict an in-hospital outcome.^{58,59} Studies suggest that older patients with heart failure are more likely to have hypoalbuminemia, and the prevalence of this condition rises with increasing frailty,⁶⁰ ranging from 20–25% among patients with chronic heart failure to about 90% among elderly individuals with severe acute heart failure.⁶¹

ALBUMIN AND STROKE

Stroke is an acute neurological injury resulting from cerebral ischemia, especially acute ischemic stroke, and cerebral hemorrhage secondary to intracerebral hemorrhage. Acute stroke, internal cerebral hemorrhage, and severe cerebral hemorrhage account for 62%, 28%, and 10%, respectively, of strokes worldwide.⁶² One of the most abundant proteins in the blood, albumin, goes through chemical changes to its N-terminal strand in ischemia, which results in the production of ischemia-modified albumin.⁶³

Studies have shown a favorable relationship between albumin levels and the outcome of stroke.^{64,65} Additionally, a low serum albumin level is one of the predictors of a first non-embolic stroke in the elderly.^{65,66} Experimental studies also suggest that albumin has a neuroprotective effect, either by reducing cerebral edema or due to its antioxidant and antiapoptotic properties.^{65,67}

Albumin increases microcirculatory flow, plasmin velocity, and oxygen transport capacity.⁶⁸ Additionally, serum albumin has a positive effect on the immune system.^{69,70} After an ischemic episode, albumin treatment increases collateral fiber density, which has a significant therapeutic effect.⁶⁹

Several studies have shown that hypoalbuminemia is associated with a poor prognosis for patients with heart failure,³⁶ stroke,⁶⁹ renal failure needing hemodialysis, and infections.⁷¹ Further research has revealed that hypoalbuminemia increases the risk of a new myocardial infarction in patients with coronary artery disease.⁷²

ALBUMIN AND ARRHYTHMIAS

Although albumin itself does not directly cause arrhythmias, alterations in its levels may reflect underlying physiological changes that can contribute to the development of arrhythmias. Monitoring albumin levels in clinical practice can provide valuable information about a patient's overall health status and help identify individuals who are more susceptible to cardiovascular complications, including diabetes.

Atrial fibrillation, the most common arrhythmia, is accompanied by the infiltration of immune cells and inflammatory proteins into the heart valves. Inflammatory mediators can alter atrial electrophysiology and structural substrates, increasing the heart's vulnerability to atrial fibrillation. Inflammation in patients with atrial fibrillation may originate from various sources, including underlying inflammatory mechanisms. Numerous systemic conditions, such as coronary heart disease, hypertension, and obesity, are associated with decreased anti-inflammatory markers and elevated levels of pro-inflammatory cytokines.^{73,74}

Guo *et al.* examined multiple inflammatory markers associated with atrial fibrillation, such as C-reactive protein, tumor necrosis factor, interleukin (IL)-2, and IL-6. They proposed several mechanisms to explain the link between inflammation and AF, such as increased platelet activation, endothelial activation or damage, monocyte factor production, and elevated fibrinogen expression.⁷⁵

Zhang *et al.* reported a direct correlation between atrial fibrillation and inflammation, validating the role of myocardial infarction and inflammatory biomarkers in predicting the recurrence of atrial fibrillation.⁷⁶ Fibrinogen and albumin levels are particularly important biomarkers that can help predict the risk of atrial fibrillation.^{77,78}

In a study by Mukamal *et al.*, higher fibrinogen and lower albumin levels were associated with an increased

risk of atrial fibrillation and a higher risk of cardiovascular disease.^{77,78}

IMPLICATIONS FOR CLINICAL PRACTICE

The measurement of serum albumin levels is an affordable, easy-to-use laboratory test that provides vital information for the prognosis of cardiovascular diseases.^{5,30} Consequently, it should be considered in the evaluation of any patient who has a cardiovascular risk. The first sign of a subclinical infection could be hypoalbuminemia, malnourishment, associated inflammatory bowel disease, enteral or urinary loss, and hypervolemia.⁵ Additionally, a dietetic study should be conducted to identify inappropriate behavior and improve eating habits. A clinical study revealed an improvement in the functional status of patients with suspected heart surgery during parenteral nutrition.^{79,80}

The management of malnutrition was reported to be favorable in a recent study. In a randomized clinical trial involving 120 patients with malnutrition and heart failure, the management of malnutrition was associated with a significant reduction in all-cause mortality during a 12-month follow-up ($p < 0.01$).^{5,51} However, it is unclear whether correcting hypoalbuminemia can improve survival among patients with heart failure. The Saline versus Albumin Fluid Evaluation (SAFE) study, which examined the use of albumine in critically ill patients, showed that the mortality rate of patients receiving albumin was equal to that of patients receiving salinized solutions.⁸⁰ Another meta-analysis concluded that a serum albumin level of >3 g/dl may slow down the rate at which deleterious effects manifest in critically ill patients with severe hypoalbuminemia.⁸¹

Inflammation is associated with reduced albumin levels, increased levels of acute-phase proteins such as hs-CRP, and increased expression of mRNA, IL-6, IL-1, and tumor necrosis factor.⁸² Studies indicate that inflammatory activity, as reflected by elevated hs-CRP, is a key factor contributing to lower serum albumin levels in acute coronary syndrome.⁸²

Patients with hypoalbuminemia should be considered at higher risk of mortality, and the underlying causes should be investigated. In this context, hypoalbuminemia is as a crucial marker with a strong physiological basis, and should be regarded as an essential component in the clinical decision-making process for managing this complex patient population.⁸³

The characteristics and results of the studies included in this analysis are presented in Table 1.

TABLE 1. General characteristics and findings of the studies included in the analysis

Study	Disease	Year	No. of patients	Mean age (years)	Male sex (%)	Outcome
Nihat <i>et al.</i> ¹⁷	Unstable angina pectoris/ NSTEMI	2020	403	64 ± 12	60.0	Patients with unstable angina pectoris or NSTEMI and hypoalbuminemia are more likely to die over the long term than those without hypoalbuminemia.
Chien <i>et al.</i> ¹⁹	Coronary heart disease	2017	734	62.1 ± 12.4	84.1	Reduced serum albumin concentration (<3.5 g/dl) impairs patients' chances of having stable congestive heart failure.
Ancion <i>et al.</i> ²⁰	Acute non-ischemic heart failure	2017	546	72 ± 12	59.0	Serum albumin level is a simple predictive factor in non-ischemic heart failure.
Northern Manhattan Study ²⁹	Incident stroke	2014	2,986	69 ± 10	37.2	There is a relationship between low serum albumin levels and stroke.
González-Pacheco <i>et al.</i> ³⁹	Acute coronary syndrome	2017	8,848	62 ± 13	74.9	A serum albumin level <3.5 g/dl is an independent predictor of newly-diagnosed heart failure in patients with acute coronary syndrome.
Kempny <i>et al.</i> ⁴⁰	Adult congenital heart disease	2015	2,886	33.3 ± 12.9	50.1	Hypoalbuminemia is a robust, independent predictor of mortality in patients with adult congenital heart disease.
Bhamidipati <i>et al.</i> ⁴¹	Coronary artery bypass grafting	2011	2,794	64.6 ± 12.9	30.1	In isolated coronary artery bypass grafting recipients, albumin is more closely linked to mortality and morbidity than the body mass index.

CONCLUSIONS

Serum albumin level is a significant predictive indicator in coronary heart disease and its measurement should be part of the routine evaluation of patients with coronary artery disease even after they are admitted to the hospital. This should lead to more attentive monitoring and measures that improve a potentially unfavorable prognosis. Hypoalbuminemia is an independent predictor of in-hospital mortality, particularly in older patients with acute myocardial infarction. Further well-designed studies evaluating functional capacity, particularly over the long term, and the causes of hypoalbuminemia in coronary artery disease are still needed.

CONFLICT OF INTEREST

Nothing to declare.

ACKNOWLEDGEMENT

This work was supported by the George Emil Palade University of Medicine, Pharmacy, Science and Technology of Târgu Mureș, Romania, research grant no. 294/2/14.01.2020.

REFERENCES

1. Van Dronkelaar C, Tieland M, Cederholm T, Reijnierse EM, Weijs PJM, Kruizenga H. Malnutrition Screening Tools Are Not Sensitive Enough to Identify Older Hospital Patients with Malnutrition. *Nutrients*. 2023;15(24):5126. doi: 10.3390/nu15245126
2. Schalk BW, Visser M, Bremmer MA, Penninx BW, Bouter LM, Deeg DJ. Change of serum albumin and risk of cardiovascular disease and all-cause mortality: Longitudinal Aging Study Amsterdam. *Am J Epidemiol*. 2006;164(10):969–977. doi: 10.1093/aje/kwj312
3. Gopal DM, Kalogeropoulos AP, Georgiopoulos VV, et al. Health ABC Study. Serum albumin concentration and heart failure risk the Health, Aging, and Body Composition Study. *Am Heart J*. 2010;160(2):279–285. doi: 10.1016/j.ahj.2010.05.022
4. Lyons O, Whelan B, Bennett K, O'Riordan D, Silke B. Serum albumin as an outcome predictor in hospital emergency medical admissions. *Eur J Intern Med*. 2010;21(1):17–20. doi: 10.1016/j.ejim.2009.10.010
5. Arques S. Serum albumin and cardiovascular disease: State-of-the-art review. *Annales de Cardiologie et d'Angéiologie*. 2020;69(4):192–200. doi: 10.1016/j.ancard.2020.07.012
6. Dorobanțu M, Halațiu VB, Gheorghe-Fronea O, et al. The Association between Apolipoprotein B, Cardiovascular Risk Factors and Subclinical Atherosclerosis-Findings from the SEPHAR National Registry on Hypertension in Romania. *Int J Mol Sci*. 2023;24(3):2813. doi: 10.3390/ijms24032813
7. Kurtul A, Murat SN, Yarlioglues M, et al. Usefulness of serum albumin concentration to predict high coronary SYNTAX score and in-hospital mortality in patients with acute coronary syndrome. *Angiology*. 2016;67(1):34–40. doi: 10.1177/0003319715575220

8. Oduncu V, Erkol A, Karabay CY, et al. The prognostic value of serum albumin levels on admission in patients with acute ST segment elevation myocardial infarction undergoing a primary percutaneous coronary intervention. *Coron Artery Dis.* 2013;24(2):88–94. doi: 10.1097/MCA.0b013e32835c46fd
9. Zhang WJ, Frei B. Albumin selectively inhibits TNF alpha-induced expression of vascular cell adhesion molecule-1 in human aortic endothelial cells. *Cardiovasc Res.* 2002;55(4):820–829. doi: 10.1016/S0008-6363(02)00492-3
10. Walley KR, McDonald TE, Wang Y, Dai S, Russell JA. Albumin resuscitation increases cardiomyocyte contractility and decreases nitric oxide synthase II expression in rat endotoxemia. *Crit Care Med.* 2003;31(1):187–194. doi: 10.1097/00003246-200301000-00029
11. Roche M, Rondeau P, Singh NR, Tarnus E, Bourdon E. The antioxidant properties of serum albumin. *FEBS Lett.* 2008;582(13):1783–1787. doi: 10.1016/j.febslet.2008.04.057
12. Anraku M, Chuang VT, Maruyama T, Otagiri M. Redox properties of serum albumin. *Biochim Biophys Acta.* 2013;1830(12):5465–5472. doi: 10.1016/j.bbagen.2013.04.036
13. Paar M, Rossmann C, Nussbold C, et al. Anticoagulant action of low, physiologic, and high albumin levels in whole blood. *PLoS One.* 2017;12(8):e0182997. doi: 10.1371/journal.pone.0182997
14. Plakht Y, Gilutz H, Shiyovich A. Decreased admission serum albumin level is an independent predictor of long-term mortality in hospital survivors of acute myocardial infarction. Soroka Acute Myocardial Infarction II (SAMI-II) project. *Int J Cardiol.* 2016;219:20–24. doi: 10.1016/j.ijcard.2016.05.067
15. Czinege M, Nyulas V, Halaşiu VB, Ţolescu C, Cojocariu LO, et al. Interrelationship between Altered Left Ventricular Ejection Fraction and Nutritional Status in the Post-Acute Myocardial Infarction Patient. *Nutrients.* 2024;16(13):2142. doi: 10.3390/nu16132142
16. Djousse L, Rothman KJ, Cupples LA, Levy D, Ellison RC. Serum albumin and risk of myocardial infarction and all-cause mortality in the Framingham Offspring Study. *Circulation.* 2002;106(23):2919–2924. doi: 10.1161/01.cir.0000042673.07632.76
17. Nihat P, Mustafa O, Mehmet AI, Bayram A. Prognostic Significance of Serum Albumin in Patients With Acute Coronary Syndrome. *Angiology.* 2020;71(10):903–908. doi: 10.1177/0003319720941747
18. Famakin B, Weiss P, Hertzberg V, et al. Hypoalbuminemia predicts acute stroke mortality: Paul Coverdell Georgia Stroke Registry. *J Stroke Cerebrovasc Dis.* 2010;19(1):17–22. doi: 10.1016/j.jstrokecerebrovasdis.2009.01.015
19. Chien SC, Chen CY, Leu HB, et al. Association of low serum albumin concentration and adverse cardiovascular events in stable coronary heart disease. *Int J Cardiol.* 2017;241(15):1–5. doi: 10.1016/j.ijcard.2017.04.003
20. Ancion A, Allepaerts S, Oury C, Gori AS, Piérard LA, Lancellotti P. Serum albumin level and hospital mortality in acute non-ischemic heart failure. *ESC Heart Fail.* 2017;4(2):138–145. doi: 10.1002/ehf2.12128
21. Rus V, Chitu M, Cernea S, et al. Altered nutritional status, inflammation and systemic vulnerability in patients with acute myocardial infarction undergoing percutaneous coronary revascularisation: A prospective study in a level 3 cardiac critical care unit. *Nutr Diet.* 2020;77(2):212–222. doi: 10.1111/1747-0080.12536
22. Czinege M, Halaşiu VB, Nyulas V, et al. Nutritional Status and Recurrent Major Cardiovascular Events Following Acute Myocardial Infarction—A Follow-Up Study in a Primary Percutaneous Coronary Intervention Center. *Nutrients.* 2024;16(7):1088. doi: 10.3390/nu16071088
23. Novack V, Pencina M, Zahger D, et al. Routine laboratory results and thirty day and one-year mortality risk following hospitalization with acute decompensated heart failure. *PLoS One.* 2010;5(8):e12184. doi: 10.1371/journal.pone.0012184
24. Yu Q, Wang Z, Ding X. Serum prealbumin as a predictor of adverse outcomes in patients with heart failure: asystematic review and metaanalysis. *Biomarkers in Medicine.* 2022;16(7):569–575. doi: 10.2217/bmm-2021-1065
25. Jacob M, Paul O, Mehringer L, et al. Albumin augmentation improves the condition of guinea pig hearts after 4 hr of cold ischemia. *Transplantation.* 2009;87(7):956–965. doi: 10.1097/TP.0b013e31819c83b5
26. Avram L, Avram C, Olah P, et al. Knowledge about food additives among adults – pilot study. *Progr Nutr.* 2021;23(1). doi: 10.23751/pn.v23i1.10631
27. Fanali G, di Masi A, Trezza V, et al. Human serum albumin: from bench to bedside. *Mol Aspects Med.* 2012;33(3):209–290. doi: 10.1016/j.mam.2011.12.002
28. Nelson JJ, Liao D, Sharrett AR, et al. Serum albumin level as a predictor of incident coronary heart disease: the Atherosclerosis Risk in Communities (ARIC) study. *Am J Epidemiol.* 2000;151(5):468–477. doi: 10.1093/oxfordjournals.aje.a010232
29. Xu WH, Dong C, Rundek T, Elkind MS, Sacco RL. Serum albumin levels are associated with cardioembolic and cryptogenic ischemic strokes: Northern Manhattan Study. *Stroke.* 2014;45(4):973–978. doi: 10.1161/STROKEAHA.113.003835
30. Arques S. Human serum albumin in cardiovascular diseases. *Eur J Int Med.* 2018;52:8–12. doi: 10.1016/j.ejim.2018.04.014
31. Arques S, Pieri B, Biegle G, Roux E, Gelisse R, Jauffret B. Comparative value of B-type natriuretic peptide and serum albumin concentration in the prediction of in-hospital mortality in elderly patients admitted for acute severe heart failure. *Ann Cardiol Angeiol (Paris).* 2009;58(5):279–283. doi: 10.1016/j.ancard.2009.09.003
32. Ronit A, Kirkegaard-Klitbo DM, Dohlmann TL, et al. Plasma albumin and incident cardiovascular disease: Results from the CGPS and an updated meta-analysis. *Arterioscler Thromb Vasc Biol.* 2020;40(2):473–482. doi: 10.1161/ATVBAHA.119.313681
33. Liu G, Hu X, Long M, Du ZM, Li Y, Hu CH. Meta-analysis of the impact of pre-procedural serum albumin on mortality in patients undergoing transcatheter aortic valve replacement. *Int Heart J.* 2020;61(1):67–76. doi: 10.1536/ihj.19-395
34. Pignatelli P, Farcomeni A, Menichelli D, Pastori D, Violi F. Serum albumin and risk of cardiovascular events in primary and secondary prevention: a systematic review of observational studies and bayesian meta-regression analysis. *Intern Emerg Med.* 2020;15(1):135–143. doi: 10.1007/s11739-019-02204-2
35. Peng W, Zhang C, Wang Z, Yang W. Prediction of all-cause mortality with hypoalbuminemia in patients with heart failure: a meta-analysis. *Biomarkers.* 2019;24(7):631–637. doi: 10.1080/1354750X.2019.1652686
36. Zhu L, Chen M, Lin X. Serum albumin level for the prediction of all-cause mortality in acute coronary syndrome patients: a meta-analysis. *Biosci Rep.* 2020;40(1):BSR20190881. doi: 10.1042/BSR20190881

37. Melenovsky V, Andersen MJ, Andress K, Reddy YN, Borlaug BA. Lung congestion in chronic heart failure: haemodynamic, clinical, and prognostic implications. *Eur J Heart Fail.* 2015;17(11):1161-1171. doi: 10.1002/ejhf.417
38. González P, Héctor Luis M, Amezcua G, et al. Levels of High-Density Lipoprotein Cholesterol are Associated With Biomarkers of Inflammation in Patients With Acute Coronary Syndrome. *Am J Cardiol.* 2015;116(11):1651-1657. doi: 10.1016/j.amjcard.2015.09.009
39. González-Pacheco H, Amezcua-Guerra LM, Sandoval J, et al. Prognostic implications of serum albumin levels in patients with acute coronary syndromes. *Am J Cardiol.* 2017;119(7):951-958. doi: 10.1016/j.amjcard.2016.11.054
40. Aleksander K, Gerhard-Paul D, Rafael G, et al. Hypoalbuminaemia predicts outcomes in adult patients with congenital heart disease. *Heart.* 2015;101(9):699-705. doi: 10.1136/heartjnl-2014-306970
41. Bhamidipati CM, LaPar DJ, Mehta GS, et al. Albumin is a better predictor of outcomes than body mass index following coronary artery bypass grafting. *Surgery.* 2011;150(4):626-634. doi: 10.1016/j.surg.2011.07.056
42. Reinhart WH. Fibrinogen-marker or mediator of vascular disease? *Vasc Med.* 2003;8(3):211-216. doi: 10.1191/1358863x03vm494ra
43. Kaptoge S, Di Angelantonio E, Pennells L, et al. C-reactive protein, fibrinogen, and cardiovascular disease prediction. *N Engl J Med.* 2012;367(14):1310-1320. doi: 10.1056/NEJMoa1107477
44. Tabakcı MM, Gerin F, Sunbul M, et al. Relation of plasma fibrinogen level with the presence, severity, and complexity of coronary artery disease. *Clin Appl Thromb Hemost.* 2017;23(6):638-644. doi: 10.1177/1076029616629210
45. Don BR, Kaysen GA. Serum albumin: relationship to inflammation and nutrition. *Semin Dial.* 2004;17(6):432-437. doi: 10.1111/j.0894-0959.2004.17603.x
46. Cetin M, Zencir C, Tasolar H, Baysal E, Balli M, Akturk E. The association of serum albumin with coronary slow flow. *Wien Klin Wochenschr.* 2014;126(15-16):468-473. doi: 10.1007/s00508-014-0559-8
47. Xia Y, Xia C, Wu L, Li Z, Li H, Zhang J. Systemic Immune Inflammation Index (SII), System Inflammation Response Index (SIRI) and Risk of All-Cause Mortality and Cardiovascular Mortality: A 20-Year Follow-Up Cohort Study of 42,875 US Adults. *J Clin Med.* 2023;12(3):1128. doi: 10.3390/jcm12031128
48. Ryszard T, Janusz S, Magdalena Starek-S, Radosław T, Andrzej R. Prognostic significance of red cell distribution width and its relation to increased pulmonary pressure and inflammation in acute heart failure. *Cardiology Journal.* 2020;27(4):394-403. doi: 10.5603/CJ.a2018.0103
49. Horwich TB, Kalantar-Zadeh K, MacLellan RW, Fonarow GC. Albumin levels predict survival in patients with systolic heart failure. *Am Heart J.* 2008;155(5):883-889. doi: 10.1016/j.ahj.2007.11.043
50. Filippatos GS, Desai RV, Ahmed MI, et al. Hypoalbuminaemia and incident heart failure in older adults. *Eur J Heart Fail.* 2011;13(10):1078-1086. doi: 10.1093/eurjhf/hfr088
51. Bonilla-Palomas JL, Gamez-Lopez AL, Moreno-Conde M, et al. Hypoalbuminemia in acute heart failure patients: Causes and its impact on hospital and long-term mortality. *J Card Fail.* 2014;20(5):350-358. doi: 10.1016/j.cardfail.2014.01.016
52. Arques S, Ambrosi P. Human serum albumin in the clinical syndrome of heart failure. *J Card Fail.* 2011;17(6):451-458. doi: 10.1016/j.cardfail.2011.02.010
53. Liu M, Chan CP, Yan BP, et al. Albumin levels predict survival in patients with heart failure and preserved ejection fraction. *Eur J Heart Fail.* 2012;14(1):39-44. doi: 10.1093/eurjhf/hfr154
54. Proboteanu P, Caloian B, Pop D. Heart Failure as the First Clinical Manifestation of Basedow's Disease. *Journal of Cardiovascular Emergencies.* 2024;10(2):78-82. doi: 10.2478/jce-2024-0012
55. Marfella R, Siniscalchi M, Esposito K, et al. Effects of stress hyperglycemia on acute myocardial infarction: role of inflammatory immune process in functional cardiac outcome. *Diabetes Care.* 2003;26(11):3129-3135. doi: 10.2337/diacare.26.11.3129
56. Danielski M, Ikizler TA, McMonagle E, et al. Linkage of hypoalbuminemia, inflammation, and oxidative stress in patients receiving maintenance hemodialysis therapy. *Am J Kidney Dis Off J Natl Kidney Found.* 2003;42(2):286-294. doi: 10.1016/S0272-6386(03)00653-X
57. Van der Pol A, Van Gilst WH, Voors AA, Van der Meer P. Treating oxidative stress in heart failure: Past, present and future. *Eur J Heart Fail.* 2019;21(4):425-435. doi: 10.1002/ejhf.1320
58. Ancion A, Allepaerts S, Oury C, Gori AS, Piérard LA, Lancellotti P. Serum albumin level and hospital mortality in acute non-ischemic heart failure. *ESC Heart Fail.* 2017;4(2):138-145. doi: 10.1002/ehf2.12128
59. Sajad RM, Bharathi, Lakshmi VS. High-Sensitivity C-Reactive Protein-to-Albumin Ratio in Predicting the Major Adverse Cardiovascular Event in Acute Coronary Syndrome at Presentation. *Indian Journal of Cardiovascular Disease in Women.* 2022;7(3):130-136. doi: 10.25259/mm_ijcdw_441
60. Arques S, Roux E, Sbragia P, Gelisse R, Pieri B, Ambrosi P. Usefulness of serum albumin concentration for in-hospital risk stratification in frail, elderly patients with acute heart failure. Insights from a prospective, monocenter study. *Int J Cardiol.* 2008;125(2):265-267. doi: 10.1016/j.ijcard.2007.07.094
61. Grimm G, Haslacher H, Kampitsch T, et al. Sex differences in the association between albumin and all-cause and vascular mortality. *Eur J Clin Invest.* 2009;39(10):860-865. doi: 10.1111/j.1365-2362.2009.02189.x
62. Krishnamurthi RV, Feigin VL, Forouzanfar MH, et al. Global and regional burden of first-ever ischaemic and haemorrhagic stroke during. Findings from the Global Burden of Disease Study 2010. *Lancet Glob Health.* 2013;1(5):e259-81. doi: 10.1016/S2214-109X(13)70089-5
63. Shevtsova A, Gordienko I, Tkachenko V, Ushakova G. Ischemia-Modified Albumin: Origins and Clinical Implications. *Dis Markers.* 2021;19:9945424. doi: 10.1155/2021/9945424
64. Babu MS, Kaul S, Dadheech S, Rajeshwar K, Jyothy A, Munshi A. Serum albumin levels in ischemic stroke and its subtypes: correlation with clinical outcome. *Nutrition.* 2013;29(6):872-875. doi: 10.1016/j.nut.2012.12.015
65. Joanna B, Jacek K, Elżbieta CC, et al. Worse Neurological State During Acute Ischemic Stroke is Associated with a Decrease in Serum Albumin Levels. *J Mol Neurosci.* 2016;58(4):493-496. doi: 10.1007/s12031-015-0705-4
66. Boselli M, Aquilani R, Baiardi P, et al. Supplementation of essential amino acids may reduce the occurrence of infections in rehabilitation patients with brain injury. *Nutr Clin Pract.* 2012;27(1):99-113. doi: 10.1177/0884533611431068

67. Chen J, Fredrickson V, Ding Y, et al. Enhanced neuroprotection by local intra-arterial infusion of human albumin solution and local hypothermia. *Stroke*. 2013;44(1):260-262. doi: 10.1161/STROKEAHA.112.675462
68. Defazio RA, Zhao W, Deng X, Obenaus A, Ginsberg MD. Albumin therapy enhances collateral perfusion after laser-induced middle cerebral artery branch occlusion: a laser speckle contrast flow study. *J Cereb Blood Flow Metab*. 2012;32(11):2012-2022. doi: 10.1038/jcbfm.2012.102
69. Dziedzic T, Pera J, Wnuk M, Szczudlik A, Słowik A. Serum albumin level as a determinant of cortisol release in patients with acute ischemic stroke. *Atherosclerosis*. 2012;221(1):212-214. doi: 10.1016/j.atherosclerosis.2011.12.017
70. Belayev L, Liu Y, Zhao W, Busto R, Ginsberg MD. Human albumin therapy of acute ischemic stroke. Marked neuroprotective efficacy at moderate doses and with a broad therapeutic window. *Stroke*. 2001;32(2):553-560. doi: 10.1161/01.str.32.2.553
71. Wallace SM, Walton BI, Kharbanda RK, Hardy R, Wilson AP, Swanton RH. Mortality from infective endocarditis: clinical predictors of outcome. *Heart*. 2002;88(1):53-60. doi: 10.1136/heart.88.1.53
72. Ballmer PE. Causes and mechanisms of hypoalbuminaemia. *Clin Nutr*. 2001;20(3):271-273. doi: 10.1054/clnu.2001.0439
73. Gregor MF, Hotamisligil GS. Inflammatory mechanisms in obesity. *Annu Rev Immunol*. 2011;29:415-445. doi: 10.1146/annurev-immunol-031210-101322
74. Marzilli M, Merz CN, Boden WE, et al. Obstructive coronary atherosclerosis and ischemic heart disease: an elusive link! *J Am Coll Cardiol*. 2012;60(11):951-956. doi: 10.1016/j.jacc.2012.02.082
75. Guo Y, Lip GYH, Apostolakis S. Inflammation in atrial fibrillation. *J Am Coll Cardiol*. 2012;60(22):2263-2270. doi: 10.1016/j.jacc.2012.04.063
76. Zhang H, Li J, Chen X, et al. Association of systemic inflammation score with atrial fibrillation: a case-control study with propensity score matching. *Heart Lung Circ*. 2018;27(4):489-496. doi: 10.1016/j.hlc.2017.04.007
77. Avram C, Gligor A, Roman D, Soylu A, Nyulas V, Avram L. Machine learning based assessment of preclinical health questionnaires. *International Journal of Medical Informatics*. 2023;180:105248. doi: 10.1016/j.ijmedinf.2023.105248
78. Kenneth JM, Janne ST, Jens F, Morten G, Gorm J. Fibrinogen and Albumin Levels and Risk of Atrial Fibrillation in Men and Women (the Copenhagen City Heart Study). *Am J Cardiol*. 2006;98(1):75-81. doi: 10.1016/j.amjcard.2006.01.067
79. Aquilani R, Opasich C, Verri M, et al. Is nutritional intake adequate in chronic heart failure patients? *J Am Coll Cardiol*. 2003;42(7):1218-1223. doi: 10.1016/s0735-1097(03)00946-x
80. Finfer S, Bellomo R, Boyce N, French J, Myburgh J, Norton R. SAFE Study Investigators. A comparison of albumin and saline for fluid resuscitation in the intensive care unit. *New Engl J Med*. 2004;350(22):2247-2256. doi: 10.1056/NEJMoa040232
81. Vincent JL, Dubois MJ, Navickis RJ, Wilkes MM. Hypoalbuminemia in acute illness: is there a rationale for intervention? A meta-analysis of cohort studies and controlled trials. *Ann Surg*. 2003;237(3):319-334. doi: 10.1097/01.SLA.0000055547.93484.87
82. Nicholson JP, Wolmarans MR, Park GR. The role of albumin in critical illness. *Br J Anaesth*. 2000;85(4):599-610. doi: 10.1093/bja/85.4.599
83. Ha CE, Bhagavan NV. Novel insights into the pleiotropic effects of human serum albumin in health and disease. *Biochim Biophys Acta*. 2013;1830(12):5486-5493. doi: 10.1016/j.bbagen.2013.04.012