

# The Role of Systemic Inflammatory Markers in the Progression and Risk of Abdominal Aortic Aneurysm Rupture – a Systematic Review

Cezar Pântea<sup>1</sup>, Adrian Vasile Mureșan<sup>2,3</sup>

<sup>1</sup> University of General Medicine, George Emil Palade University of Medicine, Pharmacy, Science and Technology, Târgu Mureș, Romania

<sup>2</sup> Clinic of Vascular Surgery, Emergency County Hospital, Târgu Mureș, Romania

<sup>3</sup> Department of Vascular Surgery, George Emil Palade University of Medicine, Pharmacy, Science and Technology, Târgu Mureș, Romania

## CORRESPONDENCE

**Cezar Pântea**

Email: [cezarpantea7@gmail.com](mailto:cezarpantea7@gmail.com)

## ARTICLE HISTORY

Received: November 15, 2023

Accepted: February 2, 2024

## ABSTRACT

Abdominal aortic aneurysm (AAA) is a serious condition where the aorta becomes abnormally dilated, which can lead to serious health risks. Inflammation plays a crucial role in AAA. The neutrophil-to-lymphocyte ratio (NLR) is a known prognostic marker for various medical conditions, including AAA. This review aims to assess the role of NLR in predicting adverse outcomes in patients with AAA. A systematic review of articles from PubMed covering the years 2013 to 2023 was conducted using the PRISMA guidelines. Out of 1,701 articles identified, 24 full-text publications were screened, and only 8 were included. The studies used receiver operating characteristic analysis to determine the best NLR cut-offs, which ranged from 4 to 9.7. The studies provide evidence that links NLR to mortality and rupture in patients with AAA. The utility of NLR utility extends beyond AAA, demonstrating significance in conditions like acute limb ischemia, chronic kidney disease, and coronary artery disease. This review underscores the potential of NLR as a valuable prognostic tool in AAA, offering insights for risk stratification and patient management.

**Keywords:** abdominal aortic aneurysm, inflammatory biomarkers, mortality, outcome

## INTRODUCTION

Abdominal aortic aneurysm (AAA) represents a potentially lethal medical condition characterized by the abnormal dilation of the aorta, the largest blood vessel in the abdomen. In AAA the maximum diameter of the abdominal aorta exceeds 30 mm, or there is an enlargement of >50% compared to the healthy aorta. The most frequent risk factors of AAA include smoking, aging, and male sex,<sup>1,2</sup> but the latest studies suggest a more significant role of inflammation responses,

mediated by matrix metalloproteinase activation, oxidative stress, intraluminal thrombus, smooth muscle apoptosis, and extracellular matrix degeneration.<sup>3–5</sup>

The neutrophil-to-lymphocyte ratio (NLR) was recently suggested as a prognostic marker for various diseases, such as end-stage kidney disease,<sup>6</sup> acute limb ischemia,<sup>7</sup> or in the assessment of outcome and patency after lower limb revascularization<sup>8</sup> and arteriovenous fistulae maturation.<sup>9</sup>

Despite numerous attempts at the medical management of AAA, the surgical approach remains the most suitable, especially for those at risk of rupture.<sup>10,11</sup>

The primary aim of this systematic review was to assess the prognostic significance of NLR in predicting adverse outcomes in patients with AAA. Our secondary aim was to establish a threshold biomarker value associated with the risks of rupture, mortality, and complications following surgical or endovascular interventions.

## METHODS

We performed a systematic review of the literature, based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines, using papers from the PubMed database and the search terms ‘abdominal aortic aneurysm’ and ‘NLR’.

Two independent reviewers (C.P. and A.V.M.) conducted the literature search. We included all papers with relevant titles that were published in English between January 1, 2013, and January 1, 2023. Studies in languages other than English, studies unrelated to this topic, inadequate data presented, non-confirmatory instances, and duplicate publications were excluded.

A total of 1,701 articles fit the search parameters, and an additional 5 articles were added from the reference lists of these articles. After removing duplicates and applying the inclusion and exclusion criteria, only 24 full-text publica-




FIGURE 1. PRISMA flow chart

tions were eligible. We eliminated 16 papers due to unclear methodology and bias, and included a total of 8 articles in our systematic review. The PRISMA flow chart is shown in Figure 1.

## RESULTS

In this comprehensive systematic review, we included eight studies published in the past decade, involving a total of 3,549 patients diagnosed with AAA. The average age of the patients was 75.8 years, with an average maximum diam-

TABLE 1. The demographic and clinical data of the participants

| Study                                 | Year | Country   | No. of patients | Mean age (years) | Male sex, n (%) | Hypertension, n (%) | Ischemic heart disease, n (%) | Diabetes, n (%) | Peripheral arterial disease, n (%) | Active smoking, n (%) |
|---------------------------------------|------|-----------|-----------------|------------------|-----------------|---------------------|-------------------------------|-----------------|------------------------------------|-----------------------|
| Appleton <i>et al.</i> <sup>12</sup>  | 2014 | UK        | 350             | 72.9             | 275 (78.57%)    | 291 (83%)           | 154 (44%)                     | —               | 46 (13%)                           | 123 (35%)             |
| Kordzadeh <i>et al.</i> <sup>13</sup> | 2015 | UK        | 80              | 75               | 66 (82.5%)      | 43 (64.2%)          | —                             | —               | —                                  | 19 (25%)              |
| Aurelian <i>et al.</i> <sup>14</sup>  | 2019 | Romania   | 255             | 68.3             | 230 (90.4%)     | 189 (74.11%)        | —                             | 36 (14.11%)     | —                                  | 66 (25.88%)           |
| Bath <i>et al.</i> <sup>15</sup>      | 2019 | USA       | 1,908           | 72.2             | 1,457 (76.4%)   | —                   | 885 (46.4%)                   | 386 (20.2%)     | —                                  | —                     |
| King <i>et al.</i> <sup>16</sup>      | 2020 | USA       | 108             | 75.5             | 78 (72.2%)      | 88 (81.5%)          | 52 (48.1%)                    | 27 (25%)        | 25 (23.1%)                         | 32 (19.6%)            |
| Tuncay <i>et al.</i> <sup>17</sup>    | 2021 | Turkey    | 42              | —                | —               | 35 (83.3%)          | 11 (26.19%)                   | 10 (23.8%)      | —                                  | 17 (40.47%)           |
| Oceau <i>et al.</i> <sup>18</sup>     | 2021 | USA       | 777             | 75.8             | 652 (84%)       | 686 (88%)           | 523 (67%)                     | 332 (43%)       | —                                  | 111 (14%)             |
| Garagoli <i>et al.</i> <sup>19</sup>  | 2022 | Argentina | 29              | —                | 20 (68.96%)     | 20 (68.96%)         | —                             | 4 (13.79%)      | —                                  | 18 (62.06%)           |

TABLE 2. AAA characteristics and outcomes, and cut-off values of NLR

| Study                                 | AAA diameter (cm) | Study group value | Control group value | Cut-off value | AUC/ROC analysis | Sensitivity (%) | Specificity (%) | Outcome                                                                             | Follow-up period |
|---------------------------------------|-------------------|-------------------|---------------------|---------------|------------------|-----------------|-----------------|-------------------------------------------------------------------------------------|------------------|
| Appleton <i>et al.</i> <sup>12</sup>  | –                 | 4.2               | 2.8                 | 5             | –                | –               | –               | Mortality                                                                           | 30 days          |
| Kordzadeh <i>et al.</i> <sup>13</sup> | 7.55              | –                 | –                   | 5             | –                | –               | –               | Mortality                                                                           | 30 days          |
| Aurelian <i>et al.</i> <sup>14</sup>  | 6.84              | 9.3               | 3.39                | 5             | –                | –               | –               | Rupture                                                                             | –                |
| Bath <i>et al.</i> <sup>15</sup>      | –                 | 13.8              | 5.93                | 9.7           | 0.662            | –               | –               | Composite outcome: in-hospital mortality, cardiac problem, or myocardial infarction | –                |
| King <i>et al.</i> <sup>16</sup>      | 5.55              | –                 | –                   | 4             | 0.780            | 62.2%           | 87.3%           | Mortality                                                                           | 5 years          |
| Tuncay <i>et al.</i> <sup>17</sup>    | –                 | 8.9               | 4.4                 | 6.6           | –                | 80%             | 89%             | Rupture                                                                             | 2.5 years        |
| Octeau <i>et al.</i> <sup>18</sup>    | 5.6               | –                 | –                   | 3.6           | –                | –               | –               | Mortality                                                                           | 5 years          |
| Garagoli <i>et al.</i> <sup>19</sup>  | –                 | –                 | –                   | 6.4           | 0.88             | 100%            | 77%             | Mortality                                                                           | In-hospital      |

TABLE 3. The association of OR and HR values with outcomes in AAA

| Study                                 | OR/HR | 95% CI |       | p value | Outcome              | Kaplan–Meier                                 | logrank p value |
|---------------------------------------|-------|--------|-------|---------|----------------------|----------------------------------------------|-----------------|
|                                       |       | Lower  | Upper |         |                      |                                              |                 |
| Kordzadeh <i>et al.</i> <sup>13</sup> | 4.28  | 1.27   | 14.42 | 0.02    | Mortality            | –                                            | –               |
| Aurelian <i>et al.</i> <sup>14</sup>  | 5.085 | 3.002  | 8.61  | <0.0001 | Rupture              | –                                            | –               |
| Bath <i>et al.</i> <sup>15</sup>      | 2.19  | 1.45   | 3.31  | 0.002   | Renal failure        | –                                            | –               |
|                                       | 2.41  | 1.21   | 4.77  | 0.01    | Cardiac complication |                                              |                 |
|                                       | 1.73  | 1.22   | 2.45  | 0.003   | Respiratory problems |                                              |                 |
|                                       | 2.59  | 1.65   | 4.07  | <0.001  | Infection            |                                              |                 |
| King <i>et al.</i> <sup>16</sup>      | 1.19  | 1.12   | 1.27  | <0.001  | Mortality            | Mortality based on NLR optimal cut-off value | <0.001          |
| Octeau <i>et al.</i> <sup>18</sup>    | 1.046 | 1.016  | 1.078 | 0.003   | Mortality            | –                                            | –               |

eter of the abdominal aorta of 6.385 cm. The study population consisted predominantly of male patients (79.21%). Hypertension emerged as the most prevalent comorbidity, affecting 82.38% of patients, followed by ischemic heart disease (51.02%) and diabetes (25.48%). In addition, 23.52% of the patients were active smokers. Other details regarding various variables are presented in Table 1.

From the eight studies reviewed, only three conducted receiver operating characteristic (ROC) analyses, revealing optimal NLR cut-off values. Specifically, Bath *et al.*<sup>15</sup> identified a value of 9.7, while Garagoli *et al.*<sup>19</sup> determined a cut-off value of 6.4, with a sensitivity of 100% and specificity of 77%. In the study by King *et al.*<sup>16</sup> the optimal NLR cut-off was found to be 4, with a sensitivity of 62.2% and specificity of 87.3%. Notably, three studies opted for a threshold value of 5, as detailed in Table 2.

Kordzadeh *et al.*,<sup>13</sup> King *et al.*,<sup>16</sup> and Octeau *et al.*<sup>18</sup> provided evidence supporting the association between NLR and the risk of mortality within 30 days and over a 5-year period following surgical or endovascular interventions for

AAA. In the investigation conducted by Aurelain *et al.*,<sup>14</sup> outlined in Table 3, elevated baseline NLR values were identified as being positively correlated with the risk of AAA rupture (odds ratio (OR) 5.085,  $p < 0.0001$ ). Additionally, Bath *et al.*<sup>15</sup> showcased that heightened levels of this biomarker are linked to increased risks of renal failure, cardiac complications, respiratory problems, and infection.

DISCUSSION

This type of analysis has been shown to be useful not only in assessing the impact of AAAs, but also in other conditions, such as acute limb ischemia. In this regard, Fajardo *et al.* showed that mortality was lower in the NLR <5 group (33%) than in the NLR >5 group (49%) ( $p \leq 0.001$ ), and amputation-free survival (AFS) was significantly higher in the NLR <5 group (50%) than in the NLR >5 group (26%) ( $p \leq 0.001$ ). In a multivariate analysis, preoperative NLR >5 was independently associated with 5-year AFS (hazard ratio (HR) 2.325, 95% confidence interval (CI) 1.732–

3.121).<sup>20</sup> Furthermore, a more recent study by Pasqui *et al.*,<sup>21</sup> which also included the platelet-to-lymphocyte ratio (PLR), showed that freedom from all amputations was significantly higher when the NLR and PLR were below the identified cut-off values ( $p < 0.0001$ ). In the same study, the NLR and PLR were found to be independent risk factors.<sup>21</sup>

Another field in which the NLR has shown its usefulness is nephrology, more specifically chronic kidney disease (CKD), a global public health problem with a high mortality rate and a rapid progression to end-stage kidney disease (ESKD). Recently, the role of inflammation and the correlation between inflammatory markers and the progression of CKD have been studied. Malhotra *et al.*<sup>22</sup> have shown that NLR could be a potential substitute marker for hs-CRP in patients undergoing hemodialysis, as it is a useful systemic inflammation test, especially in places with limited resources. Its superiority has also been proven in other studies in which the NLR reached higher levels than other markers, such as hs-CRP and interleukins, or served as an indicator of acute kidney injury in patients with sepsis or of cardiovascular events in patients with ESKD.<sup>23,24</sup> Patients with ESKD have to undergo hemodialysis, some of them using the arteriovenous fistula performed by the vascular surgeon. Some studies found a link between high inflammatory status and the failure of the arteriovenous fistula,<sup>25,26</sup> making them a useful tool for preoperative risk group stratification, better patient management, and developing predictive patterns.

The association between inflammation and atherosclerosis is well known. This process affects all arteries, particularly the cerebral, the peripheral, and the coronary arteries. Regarding the latter, several studies demonstrated that these hematological parameters are associated with the severity and prognosis of coronary artery disease. Nunez *et al.* showed that the highest values of NLR measured within the first 96 h after the onset of ST-elevation myocardial infarction symptoms carried a significant prognostic value for subsequent mortality. Moreover, this association remained significant after adjusting for key predictors such as age, reperfusion criteria, renal function, and indicators of myocardial extensions (left ventricular dysfunction, systolic blood pressure, and Killip's classification).<sup>27</sup>

Similar results were obtained in China, where a cohort study involving 2,618 patients with acute myocardial infarction presented high NLR values ( $>5.509$ ) and a high risk of mortality during hospitalization. The predictive role of high NLR values regarding negative outcomes, especially major adverse cardiovascular events (MACE), was confirmed by another study, which compared the neutrophil-to-monocyte ratio (NMR), PLR, the lymphocyte-

to-monocyte ratio (LMR), and NLR and found that these markers had the best ability to predict in-hospital death after non-ST-elevation myocardial infarction.<sup>28,29</sup>

## Study limitations

The main limitation of this study is the absence of ROC analysis in five of the eight enrolled studies. Hence, the threshold values of the NLR were not proposed and calculated according to the Youden index but were implemented from other studies in which the role of the NLR in other cardiovascular pathologies was analyzed. Therefore, it is necessary to carry out prospective multicenter studies to identify a threshold value of the NLR that can be implemented in current practice for a better stratification of risk groups. Furthermore, the heterogeneity of the groups and of the surgical or endovascular treatment between the studies, as well as the presence of AAA rupture influence the evolution of the patients and the laboratory data.

## CONCLUSIONS

AAA is a potentially fatal disorder characterized by abnormal aortic dilation, with inflammation having a critical role in its pathogenesis. The NLR has emerged as a possible predictive marker, with implications in a variety of medical settings. This systematic review showed that the NLR is useful in predicting the development, progression, and rupture of AAA, and the mortality risk of patients with this condition.

## ACKNOWLEDGEMENT

This work was supported by the George Emil Palade University of Medicine, Pharmacy, Science and Technology of Târgu Mureș, Research Grant number 165/6/10.01.2023.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## REFERENCES

1. Umehayashi R, Uchida HA, Wada J. Abdominal aortic aneurysm in aged population. *Aging (Albany NY)*. 2018;10:3650–3651.
2. Lindquist Liljeqvist M, Hultgren R, Bergman O, et al. Tunica-Specific Transcriptome of Abdominal Aortic Aneurysm and the Effect of Intraluminal Thrombus, Smoking, and Diameter Growth Rate. *Arterioscler Thromb Vasc Biol*. 2020;40:2700–2713.
3. Sakalihasan N, Limet R, Defawe OD. Abdominal aortic aneurysm. *Lancet*. 2005;365:1577–1589.
4. Sakalihasan N, Michel JB, Katsargyris A, et al. Abdominal aortic aneurysms. *Nat Rev Dis Primers*. 2018;4:34.
5. Piacentini L, Werba JP, Bono E, et al. Genome-Wide Expression Profiling Unveils Autoimmune Response Signatures in the Perivascular Adipose

- Tissue of Abdominal Aortic Aneurysm. *Arterioscler Thromb Vasc Biol.* 2019;39:237–249.
6. Mureșan AV, Russu E, Arbănași EM, et al. The Predictive Value of NLR, MLR, and PLR in the Outcome of End-Stage Kidney Disease Patients. *Biomedicines.* 2022;10:1272.
  7. Arbănași EM, Mureșan AV, Coșarcă CM, et al. Neutrophil-to-Lymphocyte Ratio and Platelet-to-Lymphocyte Ratio Impact on Predicting Outcomes in Patients with Acute Limb Ischemia. *Life.* 2022;12:822.
  8. Russu E, Mureșan AV, Arbănași EM, et al. The Predictive Role of NLR and PLR in Outcome and Patency of Lower Limb Revascularization in Patients with Femoropopliteal Disease. *Journal of Clinical Medicine.* 2022;11:2620.
  9. Kaller R, Arbănași EM, Mureșan AV, et al. The Predictive Value of Systemic Inflammatory Markers, the Prognostic Nutritional Index, and Measured Vessels' Diameters in Arteriovenous Fistula Maturation Failure. *Life.* 2022;12:1447.
  10. Bruijn LE, van Stroe Gómez CG, Curci JA, et al. A histopathological classification scheme for abdominal aortic aneurysm disease. *JVS Vasc Sci.* 2021;2:260–273.
  11. Golledge J. Abdominal aortic aneurysm: update on pathogenesis and medical treatments. *Nat Rev Cardiol.* 2019;16:225–242.
  12. Appleton ND, Bailey DM, Morris-Stiff G, Lewis MH. Neutrophil to Lymphocyte Ratio Predicts Perioperative Mortality Following Open Elective Repair of Abdominal Aortic Aneurysms. *Vasc Endovascular Surg.* 2014;48:311–316.
  13. Kordzadeh A, Malietzis G, Browne T, Prionidis I, Panayiotopoulos YP. Neutrophil to lymphocyte ratio (NLR) of five predicts 30-day morbidity in ruptured abdominal aortic aneurysms (rAAA): A retrospective cohort study. *Int J Surg.* 2015;15:45–48.
  14. Aurelian SV, Adrian M, Andercou O, et al. Neutrophil-to-Lymphocyte Ratio: A Comparative Study of Rupture to Nonruptured Infrarenal Abdominal Aortic Aneurysm. *Ann Vasc Surg.* 2019;58:270–275.
  15. Bath J, Smith JB, Kruse RL, Vogel TR. Association of neutrophil-to-lymphocyte ratio with outcomes after elective abdominal aortic aneurysm repair. *J Cardiovasc Nurs.* 2019;37:213–220.
  16. King AH, Kim AH, Kwan S, et al. Elevated Neutrophil to Lymphocyte Ratio is Associated with Worse Outcomes after Carotid Endarterectomy in Asymptomatic Patients. *J Stroke Cerebrovasc.* 2021;30:106120.
  17. Tuncay A. The effect of neutrophil/lymphocyte ratio and tomography measurements at the time of diagnosis in predicting aneurysm rupture in patients with abdominal aortic aneurysms. *Turkish Journal of Vascular Surgery.* 2021;30:1–6.
  18. Oceau D, Faries C, Barnes H, et al. Neutrophil-to-Lymphocyte Ratio Associated With Adverse Events After Endovascular Aneurysm Repair (EVAR). *Ann Vasc Surg.* 2021;75:45–54.
  19. Garagoli F, Fiorini N, Pérez MN, et al. Neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio predict in-hospital mortality in symptomatic but unruptured abdominal aortic aneurysm patients. *Int Angiol.* 2022;41:188–195.
  20. González-Fajardo JA, Brizuela-Sanz JA, Aguirre-Gervás B, et al. Prognostic Significance of an Elevated Neutrophil–Lymphocyte Ratio in the Amputation-free Survival of Patients with Chronic Critical Limb Ischemia. *Ann Vasc Surg.* 2014;28:999–1004.
  21. Pasqui E, de Donato G, Giannace G, et al. The relation between neutrophil/lymphocyte and platelet/lymphocyte ratios with mortality and limb amputation after acute limb ischaemia. *Vascular.* 2022;30:267–275.
  22. Malhotra R, Marcelli D, von Gersdorff G, et al. Relationship of Neutrophil-to-Lymphocyte Ratio and Serum Albumin Levels with C-Reactive Protein in Hemodialysis Patients: Results from 2 International Cohort Studies. *Nephron.* 2015;130:263–270.
  23. Abe T, Kato S, Tsuruta Y, et al. Neutrophil/lymphocyte ratio as a predictor of cardiovascular events in incident dialysis patients: a Japanese prospective cohort study. *Clin Exp Nephrol.* 2015;19:718–724.
  24. Yilmaz H, Cakmak M, Inan O, Darcin T, Akcay A. Can neutrophil–lymphocyte ratio be independent risk factor for predicting acute kidney injury in patients with severe sepsis? *Renal Failure.* 2015;37:225–229.
  25. Brahmabhatt A, Remuzzi A, Franzoni M, Misra S. The molecular mechanisms of hemodialysis vascular access failure. *Kidney Int.* 2016;89:303–316.
  26. Hu H, Patel S, Hanisch JJ, et al. Future research directions to improve fistula maturation and reduce access failure. *Semin Vasc Surg.* 2016;29:153–171.
  27. Núñez J, Núñez E, Bodí V, et al. Usefulness of the Neutrophil to Lymphocyte Ratio in Predicting Long-Term Mortality in ST Segment Elevation Myocardial Infarction. *Am J Cardiol.* 2008;101:747–752.
  28. Chen B, Yuan L, Chen X, et al. Correlations and Prognostic Roles of the Nutritional Status and Neutrophil-to-lymphocyte Ratio in Elderly Patients with Acute Myocardial Infarction Undergoing Primary Coronary Intervention. *Int Heart J.* 2020;61:1114–1120.
  29. Ji Z, Liu G, Guo J, et al. The Neutrophil-to-Lymphocyte Ratio Is an Important Indicator Predicting In-Hospital Death in AMI Patients. *Front Cardiovasc Med.* 2021;8:706852.