

Geological peculiarities from the Konin Lignite Mine, central Poland: An overview

Robert Wachocki^{1*}, Lilianna Chomiak², Jakub Kłesk³, Piotr Maciaszek⁴,
 Paweł Urbański⁵, Marek Widera³, Tomasz Zieliński³

¹ Konin Lignite Mine, 600-lecia 9, 62-540 Kleczew, Poland;
 e-mail: robert.wachocki@vp.pl

² District Authority Office in Poznań, Jackowskiego 18, 60-509 Poznań, Poland;
 e-mail: chomiak@amu.edu.pl

³ Institute of Geology, Adam Mickiewicz University, Krygowskiego 12, 61-680 Poznań, Poland;
 e-mail: jakub.klesk@amu.edu.pl (J.K.); widera@amu.edu.pl (M.W.); zielu@amu.edu.pl (T.Z.)

⁴ Polish Geological Institute – National Research Institute, Marine Geology Branch, Kościerska 5, 80-328 Gdańsk, Poland; e-mail: pmac@pgi.gov.pl

⁵ Polish Geological Institute – National Research Institute, Rakowiecka 4, 00-975 Warszawa, Poland;
 e-mail: purb@pgi.gov.pl

* corresponding author

Abstract

The 80th anniversary (1945–2025) of the Konin Lignite Mine (KLM) invites some summaries of the mine's characteristics. Therefore, the current study is devoted to rocks/sediments and tectonic or sedimentary structures that were observed and examined in lignite opencasts in the vicinity of the town of Konin. Some of them can be considered wonders and/or curiosities of nature, some are unique, and others are quite common. Hence, they were generally defined as geological peculiarities in this article. In stratigraphic order they are sandstones, cleats, crevasse splays, palaeochannels and palaeosols. They represent various lithostratigraphic units (formations and members) of the Neogene of central Poland, while their age ranges from the Early Miocene to the earliest Pliocene. Among the listed objects, quartzite sandstones (situated below and between the lignite beds) and palaeosols in the Poznań Clays are very common, known from other lignite opencasts in Poland. In the case of cleats and crevasse splays occurring within the lignite seam exploited by the KLM, they are among the most numerous and best developed of all lignite-bearing formations in the world. On the other hand, the presence of palaeochannels in fine-grained sediments, constituting the overburden of the exploited lignite seam, provides additional and convincing evidence for the fluvial origin of the Poznań Clays.

Key words: sandstones, cleats, crevasse splays, palaeochannels, palaeosols, Neogene

1. Introduction

The current study includes the following four opencasts belonging to the Konin Lignite Mine (KLM): Lubstów, Kazimierz N, Józwin IIB and Tomisławice. From mid-2023, only in the Tomisławice opencast, is lignite still being exploited (Fig. 1). However, in the last 2–3 years, its extraction there has been only about 0.5–0.9 mill. tonnes. Based on data from the KLM, it is estimated that lignite min-

ing in the Konin region will ultimately end in 2026–2027 (Frydrychowicz et al., 2024; Kasztelewicz et al., 2025).

Over the years, lignite opencast mining has provided great opportunities for field research. Available for direct observation were primarily exploited lignite seams and overburden deposits several dozen metres thick. This is very important because most of the geological peculiarities characterised in this paper have not been identified by exploratory



Fig. 1. Location of the open-casts belonging to the Konin Lignite Mine (KLM) mentioned in the text. Note the years of lignite mining.

drilling. In other words, they would not have been discovered and explored if it were not for open-cast mining of lignite.

The main aim of this study is to present some of the most interesting geological objects and structures that have been discovered in the open-casts of the KLM during its mining operations. The review nature of this paper means that the geological peculiarities studied are briefly described and interpreted. Detailed information about them can be found in the original studies cited in the text.

2. Geology of the study area

The lignite open-casts described in this article belong to the middle portion of the Szczecin-Miechów Synclinorium, namely to the Mogilno-Łódź Segment (Żelaźniewicz et al., 2011). To be more precise, they include the northern parts of the Konin Elevation (Widera, 1998, 2021). Three of them (i.e. Kazi-


mierz N, Józwin IIB, Tomisławice open-casts) cover fault-bounded and relatively shallow graben-like structures that are up dozens of metres deep (Widera, 2007, 2022). Lignite is found in them and on their flanks; hence, the lignite deposits represent both epirogenetic and tectonic genetic types. The exception is the Lubstów open-cast, which is located in the area of the third deepest (up to >220–240 m) tectonic graben in the Polish Lowlands territory filled with productive lignite seams (Widera, 2021; Widera et al., 2024a). In this case, lignite occurs only in the graben; therefore, this deposit should be classified exclusively as a tectonic genetic type (Widera, 2016a).

In the area of all the open-casts, rocks of the Mesozoic were drilled. These are mainly marls and carbonate sandstones of Late Cretaceous age (Pożaryski, 1952; Dadlez et al., 2000). Then, in the deepest parts of the grabens, Paleogene sediments occur. They consist mainly of marine glauconitic sands of early Oligocene (Rupelian) age in the area of the Lubstów and Tomisławice open-casts. On the contrary, in the case of the neighbouring Kazimierz N and Józwin IIB open-casts, the Neogene rests directly on the Mesozoic top (Fig. 2).

The Neogene lithostratigraphy in the Lubstów open-cast area differs significantly from that in the territories of the other characterised open-casts. This is due to the fact that in Lubstów, there is a very thick (up to 86.2 m) second Lusatian lignite seam (LLS-2), belonging to the Ścinawa Formation, which is missing in other open-casts in the Konin region (Fig. 2). The age of the LLS-2 ranges from the latest Early Miocene to the early Middle Miocene (Piwocki & Ziemińska-Tworzydło, 1997). However, in the Kazimierz N, Józwin IIB and Tomisławice open-casts, only the first Mid-Polish lignite seam (MPLS-1) is present. Lithostratigraphically, it belongs to the lower part of the Poznań Formation, called the Grey Clay Member. The MPLS-1 is up to several metres thick, on average about 6.6–6.9 m (Widera, 2021; Chomiak et al., 2024). Furthermore, in the study area, its age has been determined quite precisely at ~15.1–14.3 Ma (Widera et al., 2021a, b), i.e. the middle part of the Middle Miocene (Fig. 2).

The Wielkopolska Member (the upper part of the Poznań Formation) is also important for the described geological peculiarities. It was most fully developed (up to >30 m thick) in the Kazimierz N and Józwin IIB open-casts, where it consisted mainly of clays, silts and muds. However, among these fine-grained and multi-coloured sediments, deposits of palaeochannels and palaeosols were also found, which will be discussed below. The Neogene in each open-cast is covered by 40–60 m of glaciogenic Quaternary sediments (Fig. 2).

Fig. 2. Simplified stratigraphy with approximate location of the studied peculiarities in the opencasts of the Konin Lignite Mine (KLM).



3. Material and methods

This paper is based mainly on materials from the former Lubstów, Kazimierz N and Józwin IIB opencasts, as well as from the still-operating Tomisławice opencast. In general, fieldwork covered the years in which lignite was mined in individual opencasts (see Fig. 1), but after 1995, i.e. for the last three decades. Then, all the investigated rocks/sediments and structures were identified, described and measured, and samples were taken for laboratory analyses. Other information, including the borehole data needed to prepare lithostratigraphic profiles (see Fig. 2), was obtained from the KLM archive.

The research methods used in the current study have been described in detail in original papers. Thus, the methodology for analysing quartzite sandstones is given in the article of Górniak et al. (1996). In the case of cleats, the methods of their investigation and the presentation of results were described, for example, by Laubach et al. (1998) and

Widera (2014a, b). In turn, the crevasse splays and palaeochannels were subjected to basic sedimentological studies, including facies analysis (Miall, 2006; Boggs, 2012; Zieliński, 2014; Maciaszek et al., 2019; Widera et al., 2019; 2024b; Zieliński & Widera, 2020; Kędzior et al., 2021; and other references therein). Finally, for most of the Poznań Clays, interpreted in this study as palaeosols, in addition to sedimentological methods, mineralogical and nuclear ones were used (Klęsk et al., 2022, 2023; Klęsk, 2023).

4. Results

4.1. Lower to Mid-Miocene sandstones

Description. Quartzite sandstones were observed in all opencasts belonging to the KLM. They were located below exploited seams (LLS-2, MPLS-



Fig. 3. Quartzite sandstones in the Konin region. **A** – Large sandstone bodies in the internal dump of the Lubstów lignite opencast, 2005; **B, C** – The Konin Pillar made of sandstone in 2008 and 2011, respectively.

1), their benches or layers (see Fig. 2), which were not always preserved in the rock record. In the latter case, this is indirectly proven by the presence of fossilised roots of peat-forming woody vegetation in the upper parts of the sandstones. These rocks were most common in the Lubstów opencast in the years 1990–2005, where they occurred on several levels and were characterised by the greatest thickness and lateral spread (see Górniak et al., 1996).

The largest sandstone bodies were available for observation in the mine walls of the Lubstów opencast, but mainly in its internal dump. Most of them had a volume of more than 5 m³, while the largest sandstones had volumes even >20 m³ (Fig. 3A). They cannot be compared to those from the Bełchatów Lignite Mine, where the volume of the largest lens of such sandstones was estimated at approx. 200,000 m³ (Hycnar et al., 2022; Ratajczak et al., 2022). However, it is not the size but the signifi-

cance of quartzite sandstones from the Konin vicinity for the history of lignite mining and geology in Poland that makes it worth devoting some attention to them once again.

Interpretation. Quartzite sandstones are known from all lignite deposits, including lignite opencasts in Poland. For example, those from the nearby Adamów Lignite Mine were definitely larger (>100 m³) and better developed (Widera, 2007; Widera et al., 2022a). An interesting fact for the reader may be that the so-called Konin Pillar was made of quartzite sandstone (Fig. 3B, C), exploited in the current eastern suburbs of Konin, in the village of Brzeźno. It is the most famous symbol of Konin and the oldest road sign (milepost) in Europe, outside the borders of the former Roman Empire. The date of 1151 engraved on it is considered to be the beginning of lignite mining in Poland (e.g., Ciuk, 1994; Widera, 2021; Frydrychowicz et al., 2024; Kasztelewicz et al., 2025). Simply, to extract the sandstone, the lignite layer had to be removed (Skoczylas, 1994; Kowalczykiewicz, 1995).

For decades, there have been at least two hypotheses about the origin of the silica that holds quartz grains together. Firstly, it was associated with the dissolution of diatom shells, and secondly, with the dissolution of siliceous components common in the carbonate rocks of the Mesozoic top (e.g., Widera, 2022; Widera et al., 2022a). Therefore, studies of these freshwater sandstones from the aforementioned Lubstów opencast allowed a convincing explanation of their genesis. Using a scanning electron microscope (SEM), it was found that they were formed in the environment of Miocene palaeosols – a pedological hypothesis (Górniak et al., 1996). The sub-lignite sands were bonded by silica, as evidenced by the presence of organo-quartz aggregates in the examined samples and crystalline




Fig. 4. Sketch of the formation of freshwater quartzite sandstones in a palaeosol environment (modified from Górniak et al., 1996). Note where silica dissolves and precipitates.

silica in the form of regenerative rims. Thus, the process of silica activation was linked to the decomposition of organic matter found in the palaeosol. In more alkaline places, the detrital quartz grain was dissolved, while in places with a relative increase in acidity, silica precipitated on the same quartz grain as regenerative rims (Fig. 4; Górniak et al., 1996).

4.2. Mid-Miocene cleats

Description. The term cleats is used exclusively for coals with different ranks (lignite, hard coal, etc.) and corresponds to the vertical fractures or joints in other sedimentary rocks (e.g., McCulloch et al., 1974; Laubach et al., 1998; Widera 2014a). In the described case, cleats are natural opening-mode fractures that are almost perpendicular to the bottom and roof of the MPLS-1. These cleats, based on geometric relations (abutting and crossing patterns), are arranged in two sets that are perpendicular to both each other and the stratification. The more prominent set (longer) is called face cleats,

while the secondary set (shorter) is called butt cleats (Fig. 5).

Cleats were found in all opencasts belonging to the KLM, although they are best developed in the Kazimierz N and Józwin IIB opencasts. Of course, the face cleats are longer and more regularly oriented than the butt cleats, which are abutting the face cleats (Fig. 5). In general, the face cleats run NW–SE, while the butt cleats are oriented NE–SW (Fig. 6A). The described cleats dip very steeply, i.e. their angle of dip is in the range of 75–90° (Fig. 6B). The face and butt cleat spacing ranges approx. from 10 to 17 cm. However, their aperture is in the range of 0.2–1.9 cm (see Fig. 5). Other geometric attributes of these cleats (spacing and aperture) are characterised in detail by Widera (2014b).

Interpretation. The examined face and butt cleats are nearly perpendicular to each other, i.e. close to 90°. Simply put, they form an almost perfect, orthogonal (NW–SE and NE–SW) cleat system (compare Figs. 5, 6). On the other hand, fractures in the MPLS-1 in the vicinity of Konin can be classified as macrocleats, because their maximum ap-




Fig. 5. Cleats within the lignite seam (MPLS-1) from the Konin region. **A** – The Kazimierz N lignite opencast, 2011; **B** – The Józwin IIB lignite opencast, 2014. Note the face and butt cleats.




Fig. 6. Characteristics of cleat surfaces influencing the lignite seam (MPLS-1) from the Kazimierz N and Józwin IIB lignite opencasts. **A** – Rose diagram for face and butt cleat strikes; **B** – Rose diagram for face and butt cleat dips. Note the average values.

erture width is greater than 0.1 mm (Hooker et al., 2013).

The origin of these cleats also requires a short interpretation. First of all, they are parallel (NW-SE) and perpendicular (NE-SW) to the elongation of the main tectonic structures in central Poland. This phenomenon can be explained in at least two ways. The first hypothesis links the creation of the cleats (mainly the face cleats) with the uplift of anticlinal structures with NW-SE elongation in the sub-Cenozoic basement. The second hypothesis connects the formation of the cleats directly with the activity of the salt structure (e.g. the deep-seated salt pillow), occurring in the sub-Mesozoic substrate of the Konin area. Most likely, the NW-SE orientation of the face cleats is mainly associated with the older, regional extension, while the NE-SW butt cleat orientation is the result of local extension taking place in the cover of the salt structure (Widera, 2014b).

4.3. Mid-Miocene crevasse splays

Description. Among the numerous crevasse-splay sand bodies occurring in the KLM opencasts (e.g., Widera, 2016b, 2017, 2020; Chomiak, 2020; Widera et al., 2022b, 2023, 2024b; Dziamara et al., 2022, 2023; Chomiak et al., 2024; Wachocki et al., 2024), two of the most spectacu-

lar ones were selected for characterisation in this study. The first one is the crevasse-splay microdelta from the Józwin IIB opencast, and the second one represents the crevasse-splay complex from the Tomisławice opencast (Figs. 7, 8). All these crevasse-splay deposits were present within the exploited MPLS-1.

The crevasse-splay microdelta sediments were available for study over a length exceeding 550 m. They formed a lens-shaped and up to 2-m-thick body, extending in the N-S direction. Its middle part, up to 90 cm thick, is the most characteristic. It is composed of sands and coaly sands, steeply dipping at an angle of up to 25°. Moreover, these sediments are strongly deformed post-depositionally in both ductile and brittle manners (Fig. 7B).

In the case of the crevasse-splay complex, its sediments were perfectly available for direct observations over a distance of 400 m. The thickness of the sandy complex (with lignite intercalations) reached up to 4.8 m in the field. In this case, the sand beds are separated by 3–4 layers of lignite, with a maximum total thickness of 1–1.2 m (Fig. 8A, B). Among the mineral-organic facies, planar cross-stratified sands and coaly sands deserve mention. Their stratification is dipping at angles of up to 25° (Fig. 8C). These facies continue for over 200–300 m and occur within the lowest and highest portion of the described crevasse-splay complex.





Fig. 7. Crevasse-splay microdelta in the Józwin IIB lignite opencast in 2018. A - Broad view of the sandy crevasse splay located between two benches of the MPLS-1; B - Close-up view of the microdelta foreset; note clearly inclined layers and their deformations both brittle and ductile.

Fig. 8. Crevasse-splay complex from the Tomisławice lignite opencast in 2022. **A** – Broad view of the crevasse-splay complex; **B** – Close-up view of the crevasse-splay complex showing sandy and coaly layers; **C** – Close-up view of sand with a planar cross-stratification typical of a crevasse-splay microdelta.



Interpretation. Crevasse splays form in over-bank (extra-channel) areas by breaching natural levees during the initial stages of a flood. Therefore, they are located in close proximity to river channels (e.g., Miall, 2006; Boggs, 2012; Zieliński, 2014). Ancient crevasse-splay deposits in global coal/lignite-bearing successions have been documented relatively often. On the contrary, exposures of such sediments were not known until 2015 from the lignite-bearing Miocene of Poland (see Widera et al., 2024b). In addition, the examined crevasse splays from the Tomisławice and Józwin IIB lignite opencasts are among the most numerous and genetically most diverse in the world.

The first case described above concerns mainly, and the second case only partially, sediments deposited in standing water as a crevasse-splay microdelta. This is proven by well-developed ‘prograding splay deposits’, dipping at an angle of up to 25° and with a lateral extension of tens to hundreds of metres (compare Figs. 7, 8; Chomiak et al., 2019; Działamara et al., 2023). Other structures, not described so far among the crevasse-splay sediments from the world’s coal/lignite-bearing areas, are brittle and plastic deformations. Since the area of the Józwin IIB opencast during the Mid-Miocene peat accumulation covered a tectonically active area (Klecze Graben), the formation of the aforementioned deformations can be associated with tectonics. Most likely, plastic deformations caused by a weak seis-

mic shock occurred first. Then, brittle deformations were superimposed on them and tectonic breccia was formed as a result of a strong seismic shock (see Fig. 7; Chomiak et al., 2019, 2024).

The crevasse-splay complex from the Tomisławice opencast must be interpreted slightly differently than the one from the Józwin IIB opencast. Among the four (locally five) crevasse splays of this complex, the lowest and the highest were formed in standing water as a microdelta. The remaining crevasse splays were created in flowing water on the ‘dry’ surface of the Mid-Miocene mire, i.e. backswamp (Działamara et al., 2022, 2023; Chomiak et al., 2024). Finally, based on the age of peat accumulation, which was then transformed into lignite interlayers (up to 1.2 m thick in total; see Fig. 8A, B), it was indirectly estimated that the interpreted complex was created for approx. 50,000 years (Widera et al., 2024b).

4.4. Mid-Miocene to Upper Miocene palaeochannels

Description. In the last two decades, numerous palaeochannels have been discovered in the Poznań Clays of the Wielkopolska Member (see Fig. 2). Most of them occurred in the Kazimierz N opencast (e.g., Widera, 2012, 2013), although the largest and best sedimentologically developed ones were




Fig. 9. Broad view of the fragments of the biggest sandy and sandy-muddy palaeochannels in the opencasts of the Konin Lignite Mine (KLM). **A** - The Kazimierz N lignite opencast, 2007; **B**, **C** - The Józwin IIB lignite opencast in 2016 and 2018, respectively; note the deeply incised palaeochannel into the lignite seam and secondary palaeochannels inserted into the primary palaeochannel.

those from the Józwin IIB opencast (Fig. 9; Widera et al., 2017, 2019, 2021b; Maciaszek et al., 2019, 2020; Zieliński & Widera, 2020; Kędzior et al., 2021). In the KLM opencasts, a total of >30 palaeochannels have been documented so far, mostly surrounded by varicoloured Poznań Clays (i.e. muds, silts and clays) with palaeosol horizons, which are characterised below.

The filling of the described palaeochannels was most often a mixture of fine-grained sands to sandy silts and sandy muds (Fig. 9). However, there were also exclusively sandy or muddy ones (Widera 2012, 2013; Kędzior et al., 2021). It is also worth mentioning the occurrence of secondary channels (2–3 m deep, up to 30 m wide) within the primary channels (up to 12 m deep, up to 150 m wide), filled with muds and sandy muds (Fig. 9C). In the cross-sectional view, the majority of the examined palaeochannels were relatively deep and narrow, for which the average width/thickness ratio (w/t ratio) was <15 . The sediments filling them, especially in the Józwin IIB opencast, were characterised by numerous sedimentary structures, including heterolithic bedding, such as flaser, wavy and lenticular structures (Maciaszek

et al., 2019; Widera et al., 2019; Zieliński & Widera, 2020).

Interpretation. The cross-sectional geometry of the palaeochannels, with the w/t ratio <15 , allows them to be classified as broad ribbons (Gibling, 2006). On the other hand, the channel-fill sediments, containing the above-mentioned heterolithic bedding, demonstrate an extremely low-energy fluvial environment (e.g., Maciaszek et al., 2019). It must be added that this specific type of lamination is unique both in contemporary global fluvial sedimentary environments and in the rock record (Martin, 2000). However, the widespread presence of large-scale stratification within the palaeochannels (see Widera et al., 2019; Zieliński & Widera, 2020) indicates short-term increases in water flow energy (e.g., Miall, 2006; Boggs, 2012; Zieliński, 2014).

Taking into account the geometry of the palaeochannels and the results of the facies analysis of the sediments filling them, the morphological type of the late Neogene rivers was determined. In fact, this only applies to the area of the Konin lignite opencasts. Thus, the mentioned geometry and characteristic facies indicate an anastomosing (anabranching) type of palaeochannels. Such a pattern of these

palaeochannels was also mapped in a small area of the Józwin IIB opencast (Widera et al., 2019, their fig. 14). On the other hand, some secondary palaeochannels show features of lateral movement (i.e. meandering), while the primary ones were rather stable. All this information means that the studied palaeochannels from the Konin area have recently been interpreted as typical for the late Neogene anastomosing-to-meandering transitional fluvial system (Zieliński & Widera, 2020).

4.5. Mid-Miocene to lowermost Pliocene palaeosols

Description. In contrast to the above-described and interpreted palaeochannel deposits, now the attention will be focused on the overbank (extra-channel) deposits also occurring within the Poznań Clays. In this case, the colour variability of these very fine-grained sediments should be considered as a geological peculiarity (Fig. 10). Locally, directly on the lignite seam (MPLS-1) roof, they are made of grey clays with dispersed organic matter and fossilised wood fragments (xylites) – the uppermost Grey Clay Member. However, the most

important clays here are the so-called green and flame-like – the Wielkopolska Member (see Fig. 2).

In general, the Poznań Clays of the Wielkopolska Member consist of alternating, multi-coloured muds, silts and clays. However, what distinguishes them in the KLM opencasts, masking other textural and structural features of the sediment, are horizontally (if not disturbed post-depositionally) arranged coloured bands. Some of them are several to several hundred metres long, up to 0.3 m thick and black, and they contain >60 wt% of organic matter. They are just thin layers of lignite (Fig. 10A, B).

On the other hand, there are also sedimentary sequences which, at the top, are enriched with organic matter in amounts up to >1% wt% or do not contain it at all. In the latter case, the top layers are characterised by ‘warm’ colours (red, cherry-red, orange, yellow), which become increasingly pale towards the base (Fig. 10C, D). Of course, sediments with ‘cold’ colours (grey to black, yellow-green, blue-grey, etc.) can also be observed in the field (see Kłesk et al., 2023).

Interpretation. The above-described sediments have recently been interpreted as palaeosols formed on the floodplains of a late Neogene river system (e.g., Widera et al., 2017, 2021b; Kędzior et al., 2021).




Fig. 10. Examples of palaeosol horizons from the opencasts of the Konin Lignite Mine (KLM). **A** – The Kazimierz N lignite opencast, 2004; **B–D** – The Józwin IIB lignite opencast in 2013, 2016 and 2021, respectively. Note the colour variation of the clayey deposits and the presence of thin lignite layers.

However, their colour variability depends mainly on the presence/absence of hematite and goethite, as well as on the presence/absence of organic matter in the interpreted sediments. It should be noted that the colour of the sediments depends on the hydrological conditions (indirectly reducing-oxidising) prevailing post-depositionally (i.e. in the inter-flood periods) on the mentioned floodplains. Thus, 'warm'-coloured layers were formed at low groundwater level as Vertisols, while 'cold'-coloured layers formed at high groundwater levels as Histosols (Klęsk et al., 2023).

The above statement was confirmed by detailed studies of the Poznań Clays from the Józwin IIB opencast (Fig. 10D; Klęsk et al., 2023, 2024). For this purpose, in addition to sedimentological analysis, the following methods were used: powder X-ray diffraction (PXRD), Mössbauer spectroscopy (^{57}Fe -MS) and scanning electron microscopy with energy-dispersive spectroscopy (SEM-EDS). It turned out that sediments with 'warm' colours are relatively rich in hematite and/or goethite. On the contrary, sediments with 'cold' colours are poor in these main pigments but may contain pyrite, anatase, 'green illite' and traces of organic matter. Another strong yellow pigment commonly known from the Poznań Clays, e.g. jarosite (Duczmal-Czernikiewicz, 2013; Klęsk et al., 2022; Klęsk, 2023), may be masked mainly by organic matter. This was the situation that occurred in the grey clays lying directly on the exploited MPLS-1 in the Józwin IIB opencast (Klęsk et al., 2023).

5. Discussion

The quartzite sandstones characterised in this paper, and especially the cleats and crevasse splays from the KLM opencasts, are among the best known in Poland and in the world, respectively. Their cognitive value is great, but their origin has already been explained in the global geological literature. It can even be said that their discoveries complement the existing geological knowledge on this topic. On the contrary, the genesis of the Poznań Clays, containing the palaeochannels and palaeosols within them, looks different. Although it is mainly important for Polish geology, it requires a short discussion. This is due to the fact that the origin of the Poznań Clays has been arousing interest and controversy since the second half of the 19th century.

For nearly a century, until the 1960s, the lacustrine hypothesis of the origin of the Poznań Clays proposed by German researchers was widely accepted. There was even talk of a 'Pliocene Lake'

(e.g., Areń, 1964; Piwocki et al., 2004). Then, a lacustrine-fluvial-marine hypothesis appeared (e.g., Dyjor, 1970). In turn, two decades ago, mainly based on borehole data, Piwocki et al. (2004) proposed a fluvial hypothesis for the formation of the Poznań Clays. These researchers attributed the sandy bodies to the presumed palaeochannels and the vari-coloured fine-grained sediments (with lignite beds and even seams) to the floodplains. In addition, they associated the colour of these sediments with the degree of iron oxidation and the content of organic matter (Piwocki et al., 2004). We strongly believe that our research in the KLM opencasts, covering palaeochannel and extra-channel sediments with palaeosols and thin layers of lignite (compare Figs. 9, 10), has provided additional and convincing evidence for the fluvial genesis of the Poznań Clays. Thus, the hypothesis proposed by Piwocki et al. (2004) was positively verified.

6. Conclusions

Lignite mining in the Konin region contributed to a better understanding of the geology of this area. In the opencasts of the Konin Lignite Mine (KLM), many rocks and sediments were uncovered that can certainly be called geological peculiarities. Most of them constitute difficulties in mining activities, but they have great cognitive and aesthetic value.

Studies of quartzite sandstones from the Lubstów opencast allowed, for the first time in Poland, the determination of their pedological genesis. In turn, the cleats and crevasse splays occurring in the first Mid-Polish lignite seam (MPLS-1) are among the most numerous and best developed, not only in Polish coal/lignite-bearing formations. They complement the general knowledge about their stratigraphic architecture and genetic diversity.

The multi-coloured Poznań Clays are particularly aesthetically attractive. However, the well-documented palaeochannels and palaeosols within them confirm their terrestrial origin. Therefore, it can be stated that the Poznań Clays were formed in the environment of the late Neogene anastomosing or anastomosing-to-meandering transitional fluvial system.

Acknowledgements

The authors are very grateful to two anonymous reviewers for their efforts in evaluating our manuscript.

References

- Areń B., 1964. *Atlas geologiczny Polski. Zagadnienia stratygraficzno-facjalne. Trzeciorzęd, 11/a* [Geological atlas of Poland: stratigraphic and facies problems. Tertiary, 11/a]. Wydawnictwa Geologiczne, Warszawa.
- Boggs S., 2012. *Principles of Sedimentation and Stratigraphy*. New Jersey, Prentice Hall, 218 pp.
- Ciuk E., 1994. Z dziejów rozwoju poszukiwań, badań i górnictwa węgla brunatnego na ziemiach polskich [From the history of the development of lignite exploration, research and mining in Poland]. *Węgiel Brunatny* 9, 33–37.
- Chomiak L., 2020. Crevasse splays within a lignite seam at the Tomisławice opencast mine near Konin, central Poland: architecture, sedimentology and depositional model. *Geologos* 26, 25–37.
- Chomiak L., Maciaszek P., Wachocki R., Widera M. & Zieliński T., 2019. Seismically-induced soft-sediment deformation in crevasse-splay microdelta deposits (Middle Miocene, central Poland). *Geological Quarterly* 63, 162–177.
- Chomiak L., Działara M., Klęsk J., Wachocki R. & Widera M., 2024. „Rój” mioceńskich stożków krewasowych w Konińskim Zagłębiu Węgla Brunatnego, Polska środkowa – stan wiedzy [A “swarm” of Miocene crevasse splays in the Konin Lignite Basin, central Poland – state of knowledge]. *Przegląd Geologiczny* 72, 215–224.
- Dadlez R., Marek S. & Pokorski J. (Eds), 2000. *Mapa geologiczna Polski bez utworów kenozoiku, 1:1000000* [Geological map of Poland without Cenozoic deposits, scale 1:1000000]. Polish Geological Institute, Warsaw (in Polish).
- Duczmal-Czernikiewicz A., 2013. Evidence of soils and palaeosols in the Poznań Formation (Neogene, Polish Lowlands). *Geological Quarterly* 57, 189–204.
- Dyjur S., 1970. Seria poznańska w Polsce zachodniej [The Poznań series in West Poland]. *Kwartalnik Geologiczny* 14, 819–835.
- Działara M., Glacová V., Klęsk J., Marszałko M., Wachocki R. & Widera M., 2022. Wpływ przerostów piaszczystych w węglu brunatnym na działalność górniczą – przykład ze złoża „Pątnów IV” (okolice Konina) w środkowej Polsce [The impact of sandy interbeddings in lignite on mining activities – an example from the “Pątnów IV” deposit (Konin vicinity) in central Poland]. *Przegląd Geologiczny* 70, 376–383.
- Działara M., Kaczmarek P., Klęsk J., Wachocki R. & Widera M., 2023. Facies and statistical analyses of a crevasse-splay complex at the Tomisławice opencast lignite mine in central Poland. *Geologos* 29, 173–181.
- Frydrychowicz D., Galantkiewicz E., Kasztelewicz Z. & Widera M., 2024. 80 lat Kopalni Węgla Brunatnego Konin. Część I – historia rozpoznania złóż i ich eksploatacja [80 years of the Konin Lignite Mine. Part I – a history of deposits’ exploration and mining]. *Przegląd Geologiczny* 72, 532–545.
- Gibling M.R., 2006. Width and thickness of channel bodies and valley fills in the geological record: a literature compilation and classification. *Journal of Sedimentary Research* 76, 731–770.
- Górniak K., Bahranowski K., Ratajczak T. & Szydlak T., 1996. Regeneracja ziarn kwarcu w piaszczystych glebach korzeniowych w złożu węgla brunatnego Lubstów k. Konina [Formation of quartz overgrowth in arenaceous seatearths from the Lubstów brown-coal deposit (central Poland)]. *Przegląd Geologiczny* 44, 626–630.
- Hooker J.N., Laubach S.E. & Marrett R., 2013. Fracture-aperture size-frequency, spatial distribution, and growth processes in strata-bounded and non-strata-bounded fractures, Cambrian Mesón Group, NW Argentina. *Journal of Structural Geology* 54, 54–71.
- Hycnar E., Ratajczak T. & Uberman R., 2022. *Studium litologiczno-surowcowe skał trudnourabialnych ze złoża węgla brunatnego Betchatów* [A lithological and raw material study of difficult-to-process rocks from the Betchatów lignite deposit]. Wydawnictwo IGS MiE PAN, Kraków, 85 pp. (in Polish).
- Kasztelewicz Z., Frydrychowicz D., Galantkiewicz E. & Widera M., 2025. The past, present and future of the Konin Lignite Mine in central Poland. *Geologos* 31, 45–59.
- Kędzior A., Widera M. & Zieliński T., 2021. Ancient and modern anastomosing rivers: insights from sedimentological and geomorphological case studies of the Triassic, Neogene and Holocene of Poland. *Geological Quarterly* 65, 54.
- Klęsk J., 2023. Siarka i jej główne minerały w ilach poznańskich (górny neogen) oraz ich wpływ na barwę osadu [Sulphur and its main minerals in the Poznań Clays (upper Neogene) – their influence on the sediment colour]. *Przegląd Geologiczny* 71, 249–254.
- Klęsk J., Błachowski A., Diduszko R., Kruszewski Ł. & Widera M., 2022. Iron-bearing phases affecting the colour of upper Neogene clayey sediments from Dymaczewo Stare, west-central Poland. *Geologos* 28, 129–139.
- Klęsk J., Błachowski A., Kruszewski Ł., Michalska D., Mrozek-Wysocka M. & Widera M., 2023. Colours of the upper Neogene “Poznań Clays” in the light of sedimentological, mineralogical and nuclear methods. *Geological Quarterly* 67, 49.
- Klęsk J., Błachowski A., Kruszewski Ł., Kubiak M. & Widera M., 2024. Main colouring minerals in the ‘Poznań Clays’: Case studies from the upper Neogene in the Polish Lowlands. *Inżynieria Mineralna – Journal of the Polish Mineral Engineering Society* 53, 145–152.
- Kowalczykiewicz Z., 1995. *Zaczęło się w Brzeźnie* [It started in Brzeźno]. Wydawnictwo KWB Konin, Konin, 138 pp. (in Polish).
- Laubach S.E., Marrett R.A., Olson J.E. & Scott A.R., 1998. Characteristics and origins of coal cleat: a review. *International Journal of Coal Geology* 35, 175–207.
- Maciaszek P., Chomiak L., Wachocki R. & Widera M., 2019. The interpretive significance of ripple-derived sedimentary structures within the late Neogene fluvial succession, central Poland. *Geologos* 25, 1–13.
- Maciaszek P., Chomiak L., Urbański P. & Widera M., 2020. New insights into the genesis of the ‘Poznań

- Clays' – upper Neogene of Poland. *Civil and Environmental Engineering Reports* 30, 18–32.
- Martin A.J., 2000. Flaser and wavy bedding in ephemeral streams: a modern and an ancient example. *Sedimentary Geology* 136, 1–5.
- McCulloch C.M., Lambert S.W. & White J.R., 1976. *Determining cleat orientations of deeper coalbeds from overlying coals*. U.S. Bureau of Mines Report of Investigations, 8116, 24 pp.
- Miall A.D., 2006. *The Geology of Fluvial Deposits*. 4th ed. Springer-Verlag, Berlin, 582 pp.
- Piwocki M. & Ziemińska-Tworzydło M., 1997. Neogene of the Polish Lowlands – lithostratigraphy and pollen-spore zones. *Geological Quarterly* 41, 21–40.
- Piwocki M., Badura J. & Przybylski B., 2004. *Neogen*. [In:] Peryt T.M. & Piwocki M. (Eds): *Budowa geologiczna Polski*, t. 1, Stratygrafia część 3a, Kenozoik – paleogen, neogen (in Polish): 71–133. Polish Geological Institute, Warszawa.
- Požaryski W., 1952. Podłoże mezozoiczne Kujaw [Mesozoic basement of the Kuiavia]. *Biuletyn Państwowego Instytutu Geologicznego* 55, 26–31.
- Ratajczak T., Słomka T., Wagner M., Hycnar E., Mucha J. & Wasilewska-Błaszczak M., 2022. *Kopać albo nie kopać? węgiel brunatny ze złoża "Bełchatów" (Pole "Szczerców")*: charakter litologiczno-petrograficzny i geochemiczny oraz jakość technologiczna [To dig or not to dig? the lignite from the "Bełchatów" deposit ("Szczerców" field) lithological and petrographic character and geochemical technological quality]. Wydawnictwa AGH, Kraków, 206 pp. (in Polish).
- Skoczylas J., 1994. *Użytkowanie surowców skalnych w początkach państwa polskiego w Wielkopolsce* [The use of rock raw materials in the early Polish state in Wielkopolska]. Polskie Towarzystwo Geologiczne, Instytut Geologii UAM, Poznań, 63–74.
- Wachocki R., Dziamara M., Klęsk J. & Widera M., 2024. Comparison of the geology presented in a cross-section and that observed in the field – a lesson from the Tomislawice lignite opencast mine, Konin Basin in central Poland. *Inżynieria Mineralna – Journal of the Polish Mineral Engineering Society* 53, 417–425.
- Widera M., 1998. Ewolucja paleomorfologiczna i paleotektoniczna elewacji konińskiej [Palaeomorphological and palaeotectonical evolution of the Konin Elevation]. *Geologos* 3, 55–103.
- Widera M., 2007. *Litostratygrafia i paleotektonika kenozoiku podplejstocenijskiego Wielkopolski* [Lithostratigraphy and Palaeotectonics of the Sub-Pleistocene Cenozoic of Wielkopolska]. Adam Mickiewicz University Press, Poznań, 224 pp. (in Polish with English summary).
- Widera M., 2012. Fluwialna geneza ogniwa wielkopolskiego na podstawie danych z obszaru środkowej Polski [Fluvial origin of the Wielkopolska Member based on data from central Poland]. *Górnictwo Odkrywkowe* 53, 109–118.
- Widera M., 2013. Sand- and mud-filled fluvial palaeochannels in the Wielkopolska Member of the Neogene Poznań Formation, central Poland. *Annales Societatis Geologorum Poloniae* 83, 19–28.
- Widera M., 2014a. What are cleats? Preliminary studies from the Konin lignite mine, Miocene of central Poland. *Geologos* 20, 3–12.
- Widera M., 2014b. Lignite cleat studies from the first Middle-Polish (first Lusatian) lignite seam in central Poland. *International Journal of Coal Geology* 131, 227–238.
- Widera M., 2016a. Genetic classification of Polish lignite deposits: A review. *International Journal of Coal Geology* 158, 107–118.
- Widera M., 2016b. Depositional environments of overbank sedimentation in the lignite-bearing Grey Clays Member: New evidence from Middle Miocene deposits of central Poland. *Sedimentary Geology* 335, 150–165.
- Widera M., 2017. Sedimentary breccia formed atop a Miocene crevasse-splay succession in central Poland. *Sedimentary Geology* 360, 96–104.
- Widera M., 2020. Slump folds within mid-Miocene crevasse-splay deposits: a unique example from the Tomislawice lignite opencast mine in central Poland. *Geological Quarterly* 64, 711–722.
- Widera M., 2021. *Geologia polskich złóż węgla brunatnego* [Geology of Polish lignite deposits]. Bogucki Wydawnictwo Naukowe, Poznań, 180 pp. (in Polish).
- Widera M., 2022. *Zarys geologii okolic Poznania, Turku i Konina* [Outline of geology in the vicinity of Poznań, Turek and Konin]. Bogucki Wydawnictwo Naukowe, Poznań, 97 pp. (in Polish).
- Widera M., Kowalska E. & Fortuna M., 2017. A Miocene anastomosing river system in the area of Konin Lignite Mine, central Poland. *Annales Societatis Geologorum Poloniae* 87, 157–168.
- Widera M., Chomiak L. & Zieliński T., 2019. Sedimentary facies, processes and paleochannel pattern of an anastomosing river system: an example from the Upper Neogene of Central Poland. *Journal of Sedimentary Research* 89, 487–507.
- Widera M., Bechtel A., Chomiak L., Maciaszek P., Słodkowska B., Wachocki R., Worobiec E., Worobiec G. & Zieliński T., 2021a. Palaeoenvironmental reconstruction of the Konin Basin (central Poland) during lignite accumulation linked to the Mid-Miocene Climate Optimum. *Palaeoclimatology, Palaeoecology, Palaeogeography* 568, 110307.
- Widera M., Zieliński T., Chomiak L., Maciaszek P., Wachocki R., Bechtel A., Słodkowska B., Worobiec E. & Worobiec G., 2021b. Tectonic-climatic interactions during changes of depositional environments in the Carpathian foreland: An example from the Neogene of central Poland. *Acta Geologica Polonica* 71, 519–542.
- Widera M., Dzieduszyńska D. & Petera-Zganiacz J., 2022a. Geological and palaeogeographical peculiarities of the Adamów Graben area, central Poland. *Geologos* 28, 1–17.
- Widera M., Glacová V. & Marschalko M., 2022b. Origin of clastic partings and their impact on ash yield in mined lignite: A case study from Middle Miocene of central Poland. *Journal of Cleaner Production* 378, 134401.
- Widera M., Chomiak L. & Wachocki R., 2023. Distinct types of crevasse splays formed in the area of Middle Miocene mires, central Poland: Insights from geolog-

- ical mapping and facies analysis. *Sedimentary Geology* 443, 106300.
- Widera M., Klęsk J. & Urbański P., 2024a. Geology of the deepest Cenozoic lignite-rich grabens in Poland with particular reference to their lithostratigraphy: a comparative study. *Geology, Geophysics and Environment* 50, 131–143.
- Widera M., Działara M., Klęsk J. & Wachocki R., 2024b. Four in one: a new crevasse-splay complex in the middle Miocene of central Poland. *Annales Societatis Geologorum Poloniae* 94, 1–18.
- Zieliński T., 2014. *Sedymentologia. Osady rzek i jezior* [Sedimentology. River nad lake deposits]. Adam Mickiewicz University Press, Poznań, 594 pp. (in Polish).
- Zieliński T. & Widera M., 2020. Anastomosing-to-meandering transitional river in sedimentary record: A case study from the Neogene of central Poland. *Sedimentary Geology* 404, 105677.
- Żelaźniewicz A., Aleksandrowski P., Buła Z., Karnkowski P.H., Konon A., Ślaczka A., Żaba J. & Żytka K., 2011. *Regionalizacja tektoniczna Polski* [Tectonic division of Poland]. Komitet Nauk Geologicznych PAN, Wrocław, 60 pp. (in Polish).

Manuscript submitted: 29 January 2025

Revision accepted: 8 March 2025