

ACCEPTED AUTHOR VERSION OF THE MANUSCRIPT:

Meat quality of indigenous pig breeds in selected countries of central and southwestern Europe – a review

DOI: 10.2478/aoas-2025-0122

Dagmara Łodyga^{1,2}, Małgorzata Kasprowicz-Potocka¹, Anita Zaworska-Zakrzewska¹, Joanna Składanowska-Baryza³, Vicente Rodriguez-Estevez⁴, Santos Sanz-Fernandez⁴, Carolina Reyes-Palomo⁴, Paolo Ferrari⁵, Ewa Sell-Kubiak², Karolina Szulc³, Agnieszka Ludwiczak^{3♦}

¹Department of Animal Nutrition, Poznań University of Life Sciences, Wołyńska 33, 60-637 Poznań, Poland

²Department of Genetics and Animal Breeding, Poznań University of Life Sciences, Wołyńska 33, 60-637 Poznań, Poland

³Department of Animal Breeding and Product Quality Assessment, Poznań University of Life Sciences, Słoneczna 1, 62-002 Suchy Las, Poland

⁴Department of Animal Production, International Agricultural Campus of Excellence (ceiA3), University of Córdoba, Campus de Rabanales, 14014 Córdoba, Spain

⁵Research Centre for Animal Production (CRPA), Viale Timavo, 43/2, 42121, Reggio Emilia, Italy

♦Corresponding author: agnieszka.ludwiczak@up.poznan.pl

Received date: 21 June 2025

Accepted date: 21 October 2025

To cite this article: (2026). Łodyga D., Kasprowicz-Potocka M., Zaworska-Zakrzewska A., Składanowska-Baryza J., Rodriguez-Estevez V., Sanz-Fernandez S., Reyes-Palomo C., Ferrari P., Sell-Kubiak E., Szulc K., Ludwiczak A. (2026). Meat quality of indigenous pig breeds in selected countries of central and southwestern Europe – a review, *Annals of Animal Science*, DOI: 10.2478/aoas-2025-0122

This is unedited PDF of peer-reviewed and accepted manuscript. Copyediting, typesetting, and review of the manuscript may affect the content, so this provisional version can differ from the final version.

Meat quality of indigenous pig breeds in selected countries of central and southwestern Europe – a review

Dagmara Łodyga^{1,2}, Małgorzata Kasprończ-Potocka¹, Anita Zaworska-Zakrzewska¹, Joanna Składanowska-Baryza³, Vicente Rodriguez-Estevez⁴, Santos Sanz-Fernandez⁴, Carolina Reyes-Palomo⁴, Paolo Ferrari⁵, Ewa Sell-Kubiak², Karolina Szulc³, Agnieszka Ludwiczak^{3*}

¹Department of Animal Nutrition, Poznań University of Life Sciences, Wołyńska 33, 60-637 Poznań, Poland

²Department of Genetics and Animal Breeding, Poznań University of Life Sciences, Wołyńska 33, 60-637 Poznań, Poland

³Department of Animal Breeding and Product Quality Assessment, Poznań University of Life Sciences, Słoneczna 1, 62-002 Suchy Las, Poland

⁴Department of Animal Production, International Agricultural Campus of Excellence (ceiA3), University of Córdoba, Campus de Rabanales, 14014 Córdoba, Spain

⁵Research Centre for Animal Production (CRPA), Viale Timavo, 43/2, 42121, Reggio Emilia, Italy

*Corresponding author: agnieszka.ludwiczak@up.poznan.pl

DOI: 10.2478/aoas-2025-0122

Abstract

The conservation and promotion of local pig breeds through targeted breeding initiatives, support for niche markets, and appropriate policy measures are crucial for halting the loss of genetic diversity and ensuring a resilient, adaptable pig production industry. This fact prompted a review of the literature on native pig breeds in Europe, which are valued for their adaptability to extensive systems, their contribution to biodiversity, and the high-quality meat characteristics they produce. This review aims to examine the contribution of non-improved pig breeds to meat quality and identify the sources of the underlying differences. This paper synthesizes performance and meat quality data from major Western, Southern, Central, and Eastern European breeds, highlighting substantial diversity in growth rates, carcass yields, and physicochemical meat parameters. Despite slower growth and variable yields, many native breeds produce meat with favourable water-holding capacity, rich colour profiles, and distinct textural properties, characteristics linked to traditional feeding regimes and longer fattening cycles. The results underscore the potential of native pig breeds, not only in preserving regional heritage, but also in delivering premium-quality pork for niche markets. Understanding

their growth dynamics, carcass composition, and meat characteristics is essential for developing sustainable, high-value pork chains across Europe.

Key words: pork, native breeds, genetic diversity, extensive farming, autochthonous breeds, organic farming

The global pork industry is increasingly dominated by a narrow set of high-yielding breeds (e.g., Large White, Landrace, Duroc, Pietrain, and Hampshire and their crosses), which reflects a troubling drift toward genetic uniformity (Sponenberg et al., 2019). Many local and heritage breeds have seen drastic population declines, with some now classified as threatened or extinct (fao.org/dad-is/risk-status-of-animal-genetic-resources/en/). Loss of such breeds not only represents a decline in biodiversity but also the disappearance of unique characteristics that may be critical for future challenges (e.g., climate change, new diseases). Conservation and valorisation of local pig breeds—through breeding programmes, niche markets, and policy support—are essential to reverse the erosion of genetic diversity and maintain a robust, adaptable pig farming sector (Čandek-Potokar et al., 2019).

This review focuses on native breeds from Central and South-Western Europe, which are still used in traditional pig production systems in this region due to their excellent adaptation and meat quality. Conventional breeds may be less suitable in regions where free-range environments predominate, as they are not conducive to large litters and pose a risk of piglet mortality. Climatic conditions in these regions may also be unfavourable for conventional breeds, and native breeds cope well with food shortages and make good use of feed (Leenhouwers & Merks, 2013; Wilkinson, 2011). Moreover, in the era of industrial food production, preserving biodiversity among pigs is essential globally. In 1992, the International Convention on Biological Diversity was established, which contributed to the creation of a global system for the protection of genetic resources of farm animals (Babicz et al., 2020). More than 70% of the factors responsible for biodiversity loss are related to food production and agriculture (Kalinowska et al., 2016). To reduce the adverse effects resulting from intensive production, it is crucial to gradually implement the principles of sustainable agriculture which are associated with the extensification of animal husbandry or organic production, and thus with the promotion and maintenance of native breeds (Eisler et al., 2014), especially in those areas where traditional pig production systems using native breeds still exist. These breeds represent a reservoir of genetic diversity within the pig species. They are associated with traditional, old-fashioned production methods and pork products, sometimes related to specific agro-ecosystems, such as the Iberian pig in the Dehesa (Rodríguez-Estévez et al., 2012). They have been neglected mainly due to their lack of

competitiveness in industrial pig farming, with their lack of popularity being the main threat for their extinction. Moreover, African swine fever (ASF) can also be mentioned as a danger for native pig breeds in Europe. The disease spreads among wild boar populations, and despite the implemented biosecurity measures, outbreaks in domestic pigs were also noted (Szymańska and Dziwulaki, 2022; Škorput, et al., 2025).

Although the popularity of European local pig breeds has grown in recent years, as evidenced by a growing number of scientific publications on the subject, comprehensive syntheses remain limited. This explains the need for the present review, which aims to consolidate current knowledge and provide a clearer understanding of these breeds' value in terms of meat quality.

Local autochthonous breeds of pigs are generally more robust in response to changing environmental factors (e.g., heat waves) and more resilient to diseases and parasites than crossbred and hybrid animals (Wojtysiak & Połtowicz, 2014). This means that local breeds' productivity, fertility and meat quality are less affected by changes in environmental factors, including feed. In addition, these breeds can be fed diets based on natural resources, forage, and by-products without negative effects on health and welfare (Proorocu, 2021); thus, because these breeds are adapted to local and natural resources, they can efficiently utilise low-quality feed that does not compete with human food resources such as grains or legumes, and their breeding supports more sustainable food systems. Hence, they are a key pillar for smallholder farming and promote food security. There is, however, a drawback considering the profitability of rearing local breeds, as they are characterised by an unfavourable feed conversion ratio, increased carcass fatness, and lower carcass meatiness than commercial breeds (Kapelański et al., 1999). Varying carcass and meat quality is often the result of interactions between breed, feed management, and natural environmental factors (Rosenvold & Andersen, 2003). The breeding programmes for local breeds usually focus on preserving genetic diversity and limiting inbreeding levels rather than improving animal meat production, or robustness and foraging efficiency to make the best use of natural resources and by-products. Nevertheless, aiming to enhance the quality of pork may increasingly consider local breeds for commercial crossbreeding (Milczarek et al., 2012; Martins et al., 2020).

This review examines the carcass characteristics and meat quality of selected native pig breeds from Central and Southwestern Europe, with a focus on understanding the value of native breeds in terms of their unique traits, such as higher intramuscular fat content enhancing tenderness and juiciness, a favourable fatty acid profile rich in monounsaturated fats like oleic acid (notably in Iberian pigs), and superior sensory properties that reflect traditional, artisanal meat quality.

Material and methods

The European breeds studied here were selected from the FAO database (<https://www.fao.org/dad-is/browse-by-country-and-species/en/>) in May 2024, based on

the criterion of having a census of at least 1,000 individuals (Table 1). even though these censuses may not have been exact. Subsequently, a literature search was immediately conducted using Google Scholar. The keywords used were names of the selected breeds and different slaughter and physicochemical parameters of pork meat (e.g., “pork”, “native breeds”, “genetic diversity”, “extensive farming”, “autochthonous breeds”, “organic farming”, “pH value”, “colour”, “cooking loss”, “shear force”, “drip loss”, “fatty acid profile”, “protein”, and “fat”) (Figure 1). For the final description in this review, only breeds with at least one research paper written in English on carcass or pork parameters were included. The most important information about the slaughter procedures and meat quality is included in Tables 1 – 4. Additional information has been added as supplementary tables.

Results and discussion

Background and carcass characteristics of the breeds

Some general considerations regarding the performance of native pig breeds involve their use in extensive farming systems, which may include varying degrees of grazing. These systems are characterised by increased physical activity, dietary variability, lack of uniformity, longer production cycles, and the need to adapt fattening periods to the availability of natural resources or by-products. Additionally, management practices may aim to capitalize on compensatory growth during the fattening phase that follows a period of slower growth. Native breeds are also typically slaughtered at older ages and have higher weights than commercially improved breeds; Ludwiczak et al. (2023) have studied some of these factors in a recent review. The variety of breeds of each country are discussed in the following section.

Western and southern Europe breeds

Among the native Italian breeds, literature provides complete data on Cinta Senese, Apulo-Calabrese, Nero Siciliano, and Mora Romagnola. The fattening performance and slaughter procedures are presented in Table 2. Nero Siciliano is a breed from the island of Sicily in southern Italy that is traditionally reared under extensive conditions (Pugliese et al., 2004). It is characterised by the highest average daily gain (ADG) among Italian breeds, reaching up to 600g per day, which suggests better feed efficiency or adaptation for faster finishing (Bozzi et al., 2019). The lowest ADG was found in Cinta Senese (~ 400 g) and Apulo-Calabrese (ADG of ~300-400 g) (Caparra et al., 2023; Sirtori et al., 2014). Considering the slaughter procedures, Mora Romangola has the thickest backfat layer. Mora Romagnola comes from the Emilia–Romagna region in northern Italy. It is a slow-growing, heavy-weight type of pig (Bozzi et al., 2019), with the fat thickness ranging from 57 mm (at the last rib) to 62 mm (over the *gluteus medius*) (Fortina et al., 2005). Apulo-Calabrese and Mora Romagnola have the highest yields (~80–83%), making them productive in terms of meat output (Caparra et al., 2023; Fortina et al., 2005).

A similar balance between growth potential and product quality is also evident in Spain. The Iberian pig, the country's most iconic native breed from the southwestern Iberian Peninsula (Tejeda et al., 2020), thrives in the extensive *dehesa* system, which enhances carcass quality while preserving the sensory characteristics of Iberian meat (Robina et al., 2013; Rodríguez-Estévez et al., 2009). To adapt to modern market needs, crossbreeding with Duroc has been introduced, improving economic efficiency while maintaining the high quality of dry-cured products such as Jamón Ibérico and Lomo Ibérico (Carrapiso et al., 2015). Quality depends on a combination of the genetics, the management system, and the final fattening diet (Ortiz et al., 2020). These factors are formalised in the Iberian pig quality standard (Real Decreto 4/2014, 2014), which defines fattening periods exceeding 12 months and slaughter weights up to 200 kg (Ayuso et al., 2018). In contrast, the Porc Negre Mollorquí reaches lower slaughter weights (~120 kilograms) and has a lower ADG (410 g vs. Iberian 563 g) (Ayuso et al., 2018; Serrano et al., 2009; Tibau et al., 2019). Another Spanish breed, the Porco Celta, differs in carcass characteristics, displaying the thinnest backfat (43–48 mm) (Franco et al., 2014; 2016).

In France, fewer breeds are documented, with the Gasconne pig being the main example (Table 1). It is a native breed of pig that survives in the foothills of the Pyrenees in southwestern France. This breed is characterised by a slow growth rate and a variable daily weight gain, with the adult body weight reaching 300 kg (Mercat et al., 2019). The slaughter yield for these pigs is 80%, and the backfat thickness is 44–50 mm (Lebret et al., 2021).

Portugal also contributes valuable genetic diversity with two traditional breeds: the Alentejano and the Bisaro. The Bisaro breed is maintained in northern Portugal, with the highest concentration of farms and animals observed in the Trás-os-Montes and Minho regions (Santos Silva et al., 2019). The meat from these animals is often smoked, which is typical for this region. The breed is characterised by slow growth (ADG ~390 g) and kept to a body weight of up to 180 kg (Santos Silva et al., 2019). Compared to the Bisaro pig, the Alentejano can reach greater ADG, from 330 to 580 g.

Central and Eastern Europe

Most Central European countries have only a few native pig breeds, often adapted to traditional fat-based production (Table 1). In Slovenia, the Krškopolje pig – the only autochthonous breed – originates from Lower Carniola (Kastelic & Čandek-Potokar, 2013). Similarly, in neighbouring Croatia, one of the three native breeds, the Black Slavonian pig, is known for producing lard, bacon, and traditional high-quality meat products (Jelić et al., 2023). Both breeds share a common trait of developing a thick backfat layer, underlining their role in regional fat-rich culinary traditions. A comparable focus on fat production is also evident in the Mangalica, a Hungarian breed that traces its roots to the Serbian Šumadija. The Sumadija was brought to Hungary in the 19th century

and crossed with local breeds such as the Alfoldi, Bakony, and Szalonta (Hoha et al., 2016; Addo & Jung, 2022). The Mangalica fatteners are slaughtered at an average body weight of 130 kg and are characterised by 65–70% fat in the carcass (Egerszegi et al., 2003). Poland, by contrast, has preserved three native breeds with a more balanced profile: Puławska, Złotnicka Spotted, and Złotnicka White. The Puławska, the country's oldest native pig, combines heritage with productivity, reaching ADG values over 800 g/day (Babicz et al., 2020), making it one of the fastest-growing native breeds in Europe. Currently, this breed is a transitional form between the fat-meat type and the meat type. Among all Polish native breeds, Złotnicka Spotted is characterised by the thickest backfat layer of 41.3 mm, and at the same time, the lowest meat yield, of 41.83 % (Bocian et al., 2012; Szulc et al., 2012). As for Eastern Europe, only the Ukrainian Mirgorod pig is described in the literature. It was created in the Poltava region in Ukraine due to crossing the Ukrainian domestic pigs of variegated colours with boars of large white, Berkshire, and Tamworth breeds (Transpov, 2016). Mirgorod pigs are characterised by an average daily gain of 630 g, which places them among the fastest-growing native pigs in Europe (Tsereniuk et al., 2023). The Preštice Black-Pied Pig is the only indigenous Czech pig breed, developed in the 19th century through crossing local animals with English and German breeds. It is classified as a fat-meat type, traditionally used in the production of regional pork products due to its higher backfat thickness and moderate intramuscular fat (2.3–2.8%). Carcasses of pigs slaughtered at 94–115 kg show lean meat yields of around 48.8% and fat depths exceeding 36 mm. This breed is valued for its robustness, adaptability, and contribution to preserving Czech culinary heritage (Nevrkla et al. 2021). To summarise, most of the reviewed native breeds have a similar dressing yield between 70% and over 80%, and the lean meat content in the carcass is on average 40%. The highest daily gains were reported for Mirgorod (Ukraine), Puławska (Poland), Nero Siciliano (Italy), while the thickest backfat layer was in Mora Romagnola (Italy), and Mangalica (Hungary).

Meat quality

The most crucial physicochemical parameters of pork from selected native pig breeds are presented in Table 3. The parameters described in this chapter (i.e., pH, colour, water loss, and Warner–Bratzler shear force) were measured on the *longissimus thoracis et lumborum* (referred to as the LTL).

pH

The pH value is most commonly recorded in three time periods postmortem: 45 minutes (pH_{45 min}), 24 hours (pH_{24 h}), and 48 hours (pH_{48 h}). The pH value indicates the extent of changes occurring in the meat and indicates its quality and culinary suitability. Meat obtained directly after slaughter is characterised by a pH of 6.8–7.3. In this state, such pork has the highest capacity to bind water but has an unfavourable palatability and

tenderness. During the progressive post-slaughter changes resulting from glycolytic processes, the pH value of the meat decreases until it reaches the ultimate level. This depends on the species, preslaughter stress, degree of bleeding, cooling, or content of fat and connective tissue in the meat. All the breeds had very similar pH values measured between 45 minutes and 24 hours postmortem. The pH_{24h} ranged from 5.44 in the Bisaro breed (Martins et al., 2020) to 6.15 units in the Roma Romagnola (Fortina et al., 2005). Their highest values of pH ultimate (6.57 units) were reported in pork from the Mora Romagnola pigs (Fortina et al., 2005). For Prestice Black-Pied pigs, Nevrkla et al. (2018) report pH_{45min} is 6.29 and pH_{24h} is 5.65, values which are somewhat higher than in many commercial hybrids and indicate good meat stability. The lowest pH_{24h} values (< 5.5) were found in the meat of Porto Celta (Spain), Bisaro (Portugal), and Mangalitsa (Hungary).

Colour

Another parameter that has been examined in most publications is meat colour (CIE - Commission Internationale de l'Éclairage), which includes three main coordinates: L* (lightness), a* (redness), and b* (yellowness), and allows for quality classification. It is also no less important in the context of consumer preference studies. The primary muscle's pigment, myoglobin, converts into different chemical forms, depending on the storage conditions, including oxygen availability, temperature, and the final pH value in meat (Suman & Joseph, 2013). Consumers usually prefer meat with an intense, bright red colour. However, various factors such as breed, housing system, and nutrition, affect meat colour, thus the values reported in studies often show a wide range. Studies show that an L* value in the range of 49–60 for the LTL is perceived as desirable by consumers (Warriss & Brown, 1995). The L* (lightness) was the lowest (~40.0 units) in pork of the Iberian pigs (Spain), Mangalitsa (Hungary), and Mora Romagnola (Italy), meaning that those pigs had the darkest meat. Another essential colour coordinate is redness, marked with a*. The reddest meat (a* > 10.0 units) was found in Nero Siciliano (Italy), Cinta Senese (Italy), Porto Celta (Spain), and Alentejano (Portugal). The highest value was recorded for the Black Slavonian pig (a* = 20.02) (Karolyi et al., 2004). These findings suggest a clear interaction between pH decline and colour, reinforcing their joint role in determining consumer perception.

Cooking loss and drip loss

The ability of meat to retain water—both during storage (drip loss) and heat treatment (cooking loss)—is also influenced by pH and protein structure, creating another layer of association with the parameters above. Meat has the property of retaining water under the influence of force, and it is the water capacity (WHC) (Kilic et al., 2023; Torres Filho et al., 2017). On average, 90% of the total water is found in muscle proteins, and the remaining 10% is retained by connective tissue. A critical parameter of processed meat during heat treatment is cooking loss. During this process, the moisture content in the heat-treated product is significantly reduced (Vujadinović et al., 2014). The LTL of

native pigs is characterised by a cooking loss of 14.52-35.92%. The lowest value for this parameter was recorded for the Gasconne pig - 14.5% (Lebret et al., 2021). In contrast, just over 17% was obtained for the Alentejano breed (17.7%) and the Cinta Senese pig breed (17.94%) (Martins et al., 2020; Sirtori et al., 2014). The lowest drip loss value was recorded for the Cinta Senese breed, at 0.7% (Pugliese et al., 2005), whereas the highest value, 8.8%, was obtained for the Mangalica (Debrecéni et al., 2018). The methodology may differ slightly between publications, which could constitute an additional source of variation. The type of heat treatment influences water loss, and it has been shown that meat cooked at temperatures between 60°C and 80°C exhibits lower water loss, resulting in more balanced and beneficial cooking loss compared to heat treatment by baking (Vujadinović et al., 2014). It has also been shown that the pH during meat storage affects water loss, with higher pH levels resulting in lower losses (Drummond & Sun, 2006). Cooking loss has a wide range among the native pig breeds (14.52–35.92%, Table 3), with the Gasconne pig showing the lowest value at 14.5%, indicating well-developed WHC. The Alentejano and Cinta Senese also have relatively low cooking losses (~17.7% and 17.94% respectively). The Cinta Senese again shows good WHC with the lowest drip loss (0.7%) (Pugliese et al., 2005; Sirtori et al., 2014; Aquilani et al., 2019), while the Puławska breed shows a much higher value, reaching up to 24.89 %, suggesting poorer ability to hold moisture (Wojtysiak & Połtowicz, 2014; Bogucka & Kapelański, 2016; Debrecéni et al., 2018; Milczarek et al., 2019; Milczarek, 2021).

Texture

Finally, texture, commonly measured as Warner–Bratzler shear force (WBSF), integrates the above parameters into the eating quality perceived by consumers. Tenderness depends not only on muscle structure but also on WHC and pH, which affect juiciness and fibre resistance. The measured parameters are typically shear force (WBSF), shear energy, and Young's modulus. The shear force was chosen because it was the most prominent type of analysis in the cited literature. Table 3 presents the values for this parameter obtained using meat samples tested under various conditions. Notably, some analyses were performed on raw meat, while others were performed on thermally processed samples, resulting in a wide range of values. The values for WBSF were given in several units, as reported in the source publications. Pugliese et al. (2004) and Sirtori et al. (2014) examined the WBSF of fresh meat, which was an average of 10 kg (Nero Siciliano and Cinta Senese). Some of the publications in Table 3 reported values for this parameter in kg/cm². In this case, the highest value was recorded for the Iberian breed (4.33 kg/cm²), and the lowest value was recorded for the Porco Celta breed (3.6 kg/cm²) (Franco et al., 2014; Fernández-Barroso et al., 2020). In turn, the highest values reported by the authors were shown for the Cinta Senese breed (94-99 N), and the lowest values were reported for the Gasconne breed (24.3 N) (Pugliese et al., 2005; Lebret et al., 2021).

Chemical composition of the meat.

The comparison in Table 4 shows that the protein content in the LTL is relatively

uniform, generally ranging from 20% to 25%. The lowest protein content was observed in Mangalica pigs (19–21%), as reported in the studies of Parunović et al. (2012) and Nchama et al. (2023). While protein provides structural and nutritional value, the balance between protein and intramuscular fat (IMF) ultimately shapes the sensory and technological properties in pork. The IMF content, which determines marbling, has been repeatedly identified as a key driver of meat palatability and juiciness (Teixeira et al., 2021). It is generally recommended that its level in the LTL muscle averages around 2.5% (Sládek & Dračková, 2020). In the cited publications in Table 4, the IMF does not exceed 9%, but values as low as 1.3% have been reported, for instance, in the Mangalica breed (Imrich et al., 2020). Such variation may stem from differences in breed genetics and methodological inconsistencies across studies. Beyond genetics, environmental and management factors—such as slaughter age, body weight, and feeding regime—play a central role in IMF deposition (Lebret, 2008; Wood et al., 2008). Energy-rich diets with high starch-to-protein ratios, lipid supplementation, and extended finishing periods typically increase IMF levels, while restricted feeding and high-fibre diets can reduce them. Thus, IMF represents a complex trait shaped by the interaction between breed and management, directly influencing not only marbling but also the nutritional quality of pork (De Tonnac et al., 2023).

This link between the IMF and nutritional value becomes even more evident when considering the fatty acid profile of native breeds. The composition of fatty acids—particularly the balance of saturated, monounsaturated (MUFA), and polyunsaturated fatty acids (PUFA)—is increasingly recognised as a determinant of the contribution of pork to human health (Li et al., 2015). The contents of n-3 and n-6 polyunsaturated fatty acids (PUFAs) and their proportions are associated with human diseases, including obesity and cardiovascular diseases, and a high content of n-3 polyunsaturated fatty acids and a low ratio of n-6/n-3 polyunsaturated fatty acids are more beneficial (Simopoulos 2016). The recommended n-6/n-3 PUFA ratio ranges from 1:1 to 4:1 (Simopoulos, 2002, 2006). Still, nutritional standards state that the proper ratio of n-6 to n-3 PUFA in a daily diet should be in the range of 6:1 to 4:1 (Grześkowiak et al., 2010). The n-6/n-3 ratio in pork can be influenced by the animal's diet, breed, and husbandry system. In Table 4, the ratio of n-6 to n-3 PUFAs varies between breeds and significantly deviates from the recommended value. The values closest to the ratio of 6:1 were found for three native breeds: Apulo-Calabrian (n-6/n-3= 1.99) (Scerra et al., 2024), Gascon (n-6/n-3= 6.1–6.2) (Mercat et al., 2019) and Prestice black pied breed, which had 6.7:1 (Nevrkla et al. 2021). A less favorable n-6/n-3 PUFA ratio was noted in Cinta Senese kept indoors (24.4), and Pulawska (up to 29.4). The highest MUFA content was found in the meat of the Nero Siciliano breed: 53% (Pugliese et al., 2004) and in pork from native breeds from Spain and France, for which the value of this parameter ranged from 49.5 (Porco Celta) to 52–57.1% (Porco Celta and Gasconne) (Tejeda et al., 2020; Lebret et al., 2021; Pardo et al., 2021). Lower MUFA levels were observed in some Cinta Senese (49.9%) and Porco Celta (49.6%) groups, though still within a favorable range. The last parameter included in

Table 4 is the total PUFA content. The highest value, 15.7%, was recorded for the Apulo-Calabrese breed, whereas the lowest value, 3.39%, was recorded for the meat of the Pulawska pigs (Milczarek, 2021). PUFA content was low in Iberian (6.5–6.8%) and Mangalica (5.9–15.6%) – indicating more stable fat with less susceptibility to oxidation. These findings confirm that IMF quantity and fatty acid quality are inseparably linked characteristics—both influenced by breed genetics, diet, and management—and together they determine not only consumer satisfaction but also the nutritional value of pork. MUFA levels are consistently high in Mediterranean and Central European fatty-type breeds (Iberian, Mangalica, Pulawska), reflecting their traditional selection for meat quality and flavour. Native Northern and Eastern European breeds tend to have higher PUFA than Iberian or Mangalica, which may impact fat firmness and shelf life negatively but contribute to a healthier lipid profile. Breeds raised in outdoor/free-range systems often show a better (lower) n-6/n-3 ratio, highlighting both genetic and environmental influences.

Conclusions

This comprehensive review of native pig breeds across Southern, Central, and Eastern Europe highlights the diversity of genetic traits, regional production systems, and resulting meat qualities that define traditional pig farming. Native breeds such as the Mirgorod (Ukraine), Puławska (Poland), and Nero Siciliano (Italy) show the highest average daily gains (ADG), indicating high performance under appropriate husbandry conditions. On the other hand, breeds like the Mora Romagnola (Italy) and Mangalica (Hungary) stand out for their exceptionally thick backfat, contributing to their suitability for specific traditional meat products. Breed strongly affects the fatty acid composition of pork: Mediterranean fatty-type breeds (Iberian, Alentejano, Cinta Senese, Mangalica) are characterized by high IMF and MUFA, with relatively low PUFA – ideal for traditional dry-cured products. Northern/Eastern European breeds (Pulawska, Zlotnicka, Krskopolje) generally have lower IMF and higher PUFA, resulting in leaner but less oxidative-stable fat. N-6/n-3 ratios are influenced by both genetics and rearing system, but outdoor-raised pigs typically show more favorable ratios.

Uniqueness of meat quality

Regarding carcass characteristics, most breeds demonstrate comparable slaughter yields, yet differ in lean meat content and backfat thickness, reflecting adaptations to specific production goals. Meat quality traits, including pH, colour, water-holding capacity, and texture, further differentiate these breeds. Breeds like Cinta Senese, Gasconne, and Alentejano demonstrate low cooking and drip losses, indicative of strong water-holding capacity, which is a desirable trait in processed meat production. Meanwhile, breeds such as Black Slavonian and Nero Siciliano deliver intensely red meat, appealing to consumer preferences for richer meat colour. Texture and tenderness

vary widely, influenced by genetic factors and meat processing methods. Notably, Gasconne pigs show the lowest shear force, suggesting superior tenderness, while Cinta Senese exhibits the highest, possibly linked to muscle structure and intramuscular fat content. Lastly, the chemical composition, particularly intramuscular fat, confirms the marbling potential of these breeds. While IMF varies across studies due to methodological differences, breeds such as Mangalica and Alentejano offer high marbling, contributing to superior flavour and palatability.

Duality of pig production in modern Europe

Based on this literature review, two main paths in pig production can be brought to light. 1. The path of industrial production breeding and rearing a few commercial breeds and their crosses. This maximises efficiency but reduces resilience, robustness, and uniqueness of meat attributes. 2. The path of sustainable/niche/extensive production, where native breeds are maintained for biodiversity, and premium-quality products. This maximises the genetic diversity and promotes local products.

Future perspectives

The future of native breeds is mainly related to their role as genetic reservoirs for characteristics that may become vital in adapting to future challenges. However, there is one other perspective that may contribute to the increased popularity of these breeds. A modern consumer's attitude toward healthy nutrition shifts, leading them to search for quality products derived from sustainable animal production. EU consumers realise that while industrial systems prioritise quantity, native breeds emphasise unique meat quality from animals produced under high welfare standards. Above that, the role and future of native pig breeds extends beyond cultural preservation to sustainability, biodiversity, and food security. To summarise this review's findings, a recommendations scheme for European native pig breeds has been created (Figure 2). In the case of breeding strategies, future breeding programmes should focus on all three groups of characteristics: productivity, disease resistance and robustness, and meat quality. Native breeds could be crossed with commercial lines to introduce robustness and quality traits without losing growth and feed efficiency performance. Breeding should focus on characteristics aligned with market niches (e.g., high oleic acid for health-conscious consumers, backfat for cured products). Among breed preservation policies, the inclusion of native pig farming in rural development strategies and agrotourism should be encouraged. Consumer awareness should be increased by promoting the cultural heritage, nutritional benefits, and sustainability of native breed pork to stimulate demand and sustain premium markets. From a long-term perspective, native pig breeds should not be viewed as relics of the past but as strategic genetic resources for future food security.

Summary

In summary, native pig breeds in Europe reflect a valuable balance between cultural heritage, genetic diversity, and meat quality. While they often grow more slowly

than commercial hybrids, they provide unique sensory characteristics prized in traditional products. Genetic conservation and sustainable farming practices are essential to maintaining their role in regional economies and food traditions.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Financial support statement

This study was conducted within the project entitled: “Linking extensive husbandry practices to the intrinsic quality of pork and broiler meat” - mEATquality, funded by the European Union’s Horizon 2020 Research and Innovation program under Grant Agreement No 101000344.

References

- Aboagye G., Zappaterra M., Pasini F., Dall’Olio S., Davoli R., Nanni Costa L. (2020). Fatty acid composition of the intramuscular fat in the *longissimus thoracis* muscle of Apulo-Calabrese and crossbreed pigs. *Livest. Sci.*, 232: 103878.
- Addo S., Jung L. (2022). An insight into the runs of homozygosity distribution and breed differentiation in Mangalitsa pigs. *Front. Genet.*, 2967.
- Álvarez-Rodríguez J., Teixeira A. (2019). Slaughter weight rather than sex affects carcass cuts and tissue composition of Bisaro pigs. *Meat Sci.*, 154: 54–60.
- Aquilani C., Sirtori F., Franci O., Acciaioli A., Bozzi R., Pezzati A., Pugliese C. (2019). Effects of protein restriction on performances and meat quality of Cinta Senese pig reared in an organic system. *Animals*, 9: 310.
- Ayuso Carrizosa D., González A., Peña F., Izquierdo M. (2018). Changes in adipose cells of the longissimus dorsi muscle in Iberian pigs raised under extensive conditions. *An. Acad. Bras. Ciênc.*, 90: 247–253.
- Babicz M., Hałabis M., Skąlecki P., Domaradzki P., Litwińczuk A., Kropiwniec-Domańska K., Łukasik M. (2020). Breeding and Performance Potential of Puławska Pigs-A Review. *Ann. Anim. Sci.*, 20: 343–354.
- Belkova J., Rozkot M., Weisbaueroval E., Truneckova J. (2023). Influence of feed substitution with acorns on fattening performance, carcass characteristics, and meat

composition of Prestice Black-Pied pigs finished in a conventional indoor system. *Cogent Food Agric.*, 9:2220182.

- Bocian M., Wojtysiak D., Jankowiak H., Cebulska A., Kapelański W., Migdał W. (2012). Carcass, meat quality and histochemical traits of *m. longissimus lumborum* from Złotnicka spotted pigs and commercial pigs. *Folia Biol. (Czech Republic)*, 60: 181–187.
- Bogucka J., Kapelański W. (2016). Microstructure of longissimus lumborum muscle and meat quality of native Polish pig breeds: Złotnicka Spotted and Puławska. *Ann. Anim. Sci.*, 16: 1199–1210.
- Bozzi R., Gallo M., Geraci C., Fontanesi L., Batorek-Lukač N. (2019). Nero Siciliano Pig. In M. Čandek-Potokar & R. Nieto, *European Local Pig Breeds - Diversity and Performance. A Study of Project TREASURE*, Chapter 18, pp. 1–11.
- Butko D., Senčić D., Antunović Z., Šperanda M., Steiner Z. (2007). Pork carcass composition and the meat quality of the Black Slavonian Pig - The endangered breeds in the indoor and outdoor keeping system. *Poljoprivreda*, 13: 167–171.
- Čandek-Potokar M., Žlender B., Kramar Z., Šegula B., Fazarinc G., Uršic M. (2003). Evaluation of Slovene local pig breed Krškopolje for carcass and meat quality. *Czech J. Anim. Sci.*, 48: 120–128.
- Čandek-Potokar M., Linan R.M.N. (2019). *European Local Pig Breeds—Diversity and Performance*; Intech Open: London, UK.
- Caparra P., Chies L., Scerra M., Foti F., Bognanno M., Cilione C., De Caria P., Claps S., Cifuni, G. F. (2023). Effect of Dietary Ensiled Olive Cake Supplementation on Performance and Meat Quality of Apulo-Calabrese Pigs. *Animals*, 13: 1–13.
- Carrapiso, A. I., Martín-Cabello L., Torrado-Serrano C., Martín L. (2015). Sensory Characteristics and Consumer Preference of Smoked Dry-Cured Iberian Salchichon. *Int. J. Food Prop.*, 18: 1964–1972.
- Charlton, C. E., Reeves Pitts M., Rehm, J. G., Sawyer, J. T., Brandebourg T. D. (2022). Determination of Optimal Harvest Weight for Mangalica Pigs Using a Serial Harvest Approach to Measure Growth Performance and Carcass Characteristics. *Foods*, 11: 3958.
- Charneca R., Martins J., Freitas A., Neves J., Nunes J., Paixim H., Bento P., Batorek-Lukač N. (2019). Alentejano Pig. In M. Čandek-Potokar & R. Nieto, *European Local Pig Breeds - Diversity and Performance. A Study of Project TREASURE*, Chapter 2, pp. 1-24.
- De Tonnac A., Labussière E., Vincent A., Mourot J. (2023). Effects of Pig Dietary n-6/n-3 Polyunsaturated Fatty Acids Ratio and Gender on Carcass Traits, Fatty Acid

Profiles, Nutritional Indices of Lipid Depots and Oxidative Stability of Meat in Medium–Heavy Pigs. *Foods*, 12: 4106.

Debrecéni O., Lípová P., Bucko O., Cebulská A., Kapelánski W. (2018). Effect of pig genotypes from Slovak and Polish breeds on meat quality. *Arch. Anim. Breed.*, 61: 99–107.

Drummond, L. S. Sun D. W. (2006). Feasibility of water immersion cooking of beef joints: Effect on product quality and yield. *J. Food Eng.*, 77: 289–294.

Egerszegi I., Rátky J., Solti L., Brussow K. P. (2003). Mangalica - an indigenous swine breed from Hungary (Review). *Arch. Anim. Breed.*, 46: 245–256.

Eisler, M. C., Lee, M. R. F., Tarlton, J. F., Martin, G. B., Beddington J., Dungait, J. A. J., Greathead H., Liu J., Mathew S., Miller H., Misselbrook, T. H., Murray, P. J., Vinod, V. K., Van Saun R., Winter, M. (2014). Steps to sustainable livestock. *Symp. Soc. Exp. Biol.*, 19: 205–234.

FAO (Food and Agriculture Organization of the United Nations). (2025). Risk status of animal genetic resources. DAD-IS, FAO. <https://www.fao.org/dad-is/risk-status-of-animal-genetic-resources/en/>

FAO (Food and Agriculture Organization of the United Nations). (2025). Browse by country and species. DAD-IS, FAO. <https://www.fao.org/dad-is/browse-by-country-and-species/en/>

Fernández-Barroso M., Silió L., Rodríguez C., Palma-Granados P., López A., Carballo C., Sánchez-Esquiliche F., Gómez-Carballar F., García-Casco, J. M., Muñoz M. (2020). Genetic parameter estimation and gene association analyses for meat quality traits in open-air free-range Iberian pigs. *J. Anim. Breed. Genet.*, 137: 581–598.

Fortina R., Barbera S., Lussiana C., Mimosi A., Tassone S., Rossi A., Zanardi E. (2005). Performances and meat quality of two Italian pig breeds fed diets for commercial hybrids. *Meat Sci.*, 71: 713–718.

Franco D., Carballo J., Bermúdez R., Lorenzo J. M. (2016). Effect of genotype and slaughter age on carcass traits and meat quality of the Celta pig breed in extensive system. *Ann. Anim. Sci.*, 16: 259–273.

Franco D., Vazquez, J. A., Lorenzo J. M. (2014). Growth performance, carcass and meat quality of the Celta pig crossbred with Duroc and Landrace genotypes. *Meat Sci.*, 96: 195–202.

Freitas A. B., Neves J. A., Nunes J. T., Martins J. M. (2007). Effects of a high-fat diet on growth performances and carcass characteristics of Alentejano pigs. In: *Book of Abstracts of the 6th International Symposium on the Mediterranean Pig*, Capo d'Orlando, Messina, Italy.

- Grześkowiak E., Borys A., Borzuta K., Buczyński, J. T., Lisiak, D. (2009). Slaughter value, meat quality and backfat fatty acid profile in Złotnicka White and Złotnicka Spotted fatteners. *Anim. Sci. Pap. Rep.*, 27: 115–125.
- Grześkowiak E., Borzuta K., Lisiak D., Strzelecki J., Janiszewski P. (2010). Właściwości fizykochemiczne i sensoryczne oraz skład kwasów tłuszczowych mięśnia *longissimus dorsi* mieszańców PBZ x WBP oraz PBZ x (D x P). *Żywność. Nauka. Technologia. Jakosc/Food. Science Technology. Quality*, 17: 189–198.
- Hoha, G. V, Pandelea M., Costăchescu E., Nistor, C. E., Măgdicu E., Păsărin B. (2016). The quality of meat to different type Mangalitsa breeds. *Scientific Papers-Animal Science Series: Lucrări Științifice - Seria Zootehnie*, 65: 84–87.
- Imrich I., Mlyneková E., Mlynek J., Halo M., Kank T. (2020). Comparison of the physico-chemical meat quality of the breeds Mangalitsa and Large White with regard to the slaughter weight. *Potr. S. J. F. Sci.*, 14: 135–141.
- Imrich I., Mlyneková E., Mlynek J., Kanka T. (2019). The technological meat quality of the White Mangalitsa breed. *Acta Fytotech. Zootech.*, 22: 124–129.
- Jelić Milković S., Crnčan A., Kristić J., Kralik I., Djurkin Kušec I., Gvozdanović K., Kušec G., Kralik Z., Lončarić R. (2023). Consumer Preferences for Cured Meat Products from the Autochthonous Black Slavonian Pig. *Foods*, 12: 3666.
- Kalinowska A. (2016). Jak przyspieszyć zmiany świadomości, aby spowolnić tempo utraty bioróżnorodności biologicznej. *Polish J. Sust. Develop.*, 20: 67–76.
- Kapelański W., Buczyński, J. T., Bocian M. (2006). Slaughter value and meat quality of the Polish native Złotnicka Spotted pig. *Anim. Sci. Pap. Rep.*, 24: 7–13.
- Kapelanski W., Kortz J., Rak B., Grajewska S., Bocian M. (1999). Meat quality of pietrain and zlotniki spotted pigs and their crossbreds evaluated in 1969 and 1997. *Adv. Agric. Sci.*, 6: 25–31.
- Karolyi D., Salajpal K., Sinjeri, Ž., Kovačić D., Jurić I., Đikić M. (2004). Meat quality, blood stress indicators and trimmed cut yield comparison of Black Slavonian pig with modern pigs in the production of Slavonian kulen. *Acta Agric. Slov.*, 1: 67–72.
- Kastelic A., Čandek-potokar M. (2013). Application of Quality Labels in Support of Conservation of Local Breeds – a Challenge for Slovenian Krškopolje Pig. *Acta Agric. Slov. (Supplement)*, 4: 205–209.
- Kilic S., Oz E., Turhan S., Amarowicz R., Toldra F., Brennan C., Oz F. (2023). The effect of sealing process on the textural and sensory properties and protein profile of cooked meat. *Int. J. Food Sci. Technol.*, 58: 4800–4811.
- Kušec G., Gvozdanović K., Djurkin Kušec I., Margeta V., Galović D., Radišić Ž. (2019). Changes in carcass composition of black Slavonian pigs and their crossbreds with

- Duroc during growth. *J. Cent. Eur. Agric.*, 20: 43–49.
- Lebret B. (2008). Effects of feeding and rearing systems on growth, carcass composition and meat quality in pigs. *Animal*, 2: 1548–1558.
- Lebret B., Lenoir H., Daré S., Fonseca A., Fève K., Riquet J., Mercat, M. (2021). Finishing season and feeding resources influence the quality of products from extensive-system Gascon pigs. Part 1: Carcass traits and quality of fresh loin. *Animal*, 15: 100240.
- Leenhouwers, J. I., Merks J.W.M. (2013). Suitability of traditional and conventional pig breeds in organic and low-input production systems in Europe: Survey results and a review of literature. *Ani. Genet. Resour.*, 53: 169–184.
- Li F., Duan Y., Li Y., Tang Y., Geng M., Oladele, O.A., Kim, S.W., Yin Y. (2015). Effects of dietary n-6 : n-3 PUFA ratio on fatty acid composition, free amino acid profile and gene expression of transporters in finishing pigs. *Br. J. Nutr.*, 113(5), 739–748.
- Ludwiczak A., Kasprowicz-Potocka M., Zaworska-Zakrzewska A., Składanowska-Baryza J., Rodriguez-Estevéz V., Sanz-Fernandez S., Diaz-Gaona C., Ferrari P., Pedersen L.J., Couto M.Y.R., Revilla I., Sell-Kubiak E. (2023). Husbandry practices associated with extensification in European pig production and their effects on pork quality. *Meat Sci.*, 206: 109339.
- Lukač N.B., Tomažin U., Škrlep M., Kastelic A., Poklukar K., Čandek-Potokar M. (2019). Krškopoljski prašič (Krškopolje Pig). In M. Čandek-Potokar & R. Nieto, *European Local Pig Breeds - Diversity and Performance. A Study of Project TREASURE*, Chapter 10, pp. 1-14.
- Martins J.M., Fialho R., Albuquerque A., Neves J., Freitas A., Nunes, J. T., Charneca R. (2020). Growth, blood, carcass and meat quality traits from local pig breeds and their crosses. *Animal*, 14: 636–647.
- Martins J.M., Silva D., Albuquerque A., Neves J., Charneca R., Freitas A. (2021). Physical activity effects on blood parameters, growth, carcass, and meat and fat composition of portuguese alentejano pigs. *Animals*, 11: 1–16.
- Mercat M.-J., Lebret B., Lenoir H., Lukač N.B. (2019). Gascon Pig. In M. Čandek-Potokar & R. Nieto, *European Local Pig Breeds - Diversity and Performance. A Study of Project TREASURE*, Chapter 8, pp. 1-14.
- Micari P., Racinaro L., Sarullo V., Carpino S., Marzullo A. (2009). Zoometric rates, reproductive and productive parameters of the Apulo-calabrian swine, obtained in breeding certified by ANAS Calabria. *Ital. J. Anim. Sci.*, 8: 519–521.
- Migdał W., Radović, Č., Živković V., Gwiazda E., Migdał, Ł., Migdał A., Walczycka

- M., Węsierska E., Zając M., Tkaczewska J., Kulawik P., Krępa-Stefanik K. (2017). Quality of meat from native pigs. Proceedings of the 11th International Symposium Modern Trends in Livestock Production, October 11-13, pp. 189–199.
- Milczarek A. (2021). Carcass composition and quality of meat of Pulawska and Pulawska x PLW crossbred pigs fed rations with naked oats. *Animals*, 11: 3342.
- Milczarek A., Osek M., Banaszkiewicz T. (2019). Chemical composition of meat from the Pulawska breed pigs, depending on their slaughter weight. *J. Elem.*, 24: 639–648.
- Milczarek A., Osek M., Olkowski B., Klocek B., Lipnicka A. (2012). The comparison of fattening results, slaughter value and meat quality of pig crossbreeds Pulawska x Polish Large White and Polish Landrace x Polish Large White. *Acta Sci. Pol. Zootech.*, 11: 31–40.
- Nchama C.N.N., Saccà E., Sepulcri A., Foletto V., Brunner I., Corazzin M., Piasentier E. (2023). Characteristics of carcass, meat and subcutaneous fat of Mangalica pigs reared outdoors. *Inorg. Chem. Industry*, 57: 625–634.
- Nevrkla P., Václavková E., Horký P., Kamanová V., Hadaš Z., Rečková Z., Máchal L. (2018). Growth and Meat Quality of Prestice Black-Pied and (Landrace × Large White) × Duroc Pigs. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, 66(3), 701-705
- Nevrkla P., Václavková E., Rozkot M. (2021). The Indigenous Prestice Black-Pied Pig Breed Differs from a Commercial Hybrid in Growth Intensity, Carcass Value and Meat Quality. *Agriculture*, 11(4), 331.
- Ortiz A., García-Torres S., González E., De Pedro-Sanz, E. J., Gaspar P., Tejerina D. (2020). Quality traits of fresh and dry-cured loin from Iberian x Duroc crossbred pig in the Montanera system according to slaughtering age. *Meat Sci.*, 170: 108242.
- Pardo Z., Fernández-Fígares I., Lachica M., Lara L., Nieto R., Seiquer I. (2021). Impact of heat stress on meat quality and antioxidant markers in Iberian pigs. *Antioxidants*, 10: 1911.
- Parunović N., Petrović M., Matekalo-Sverak V., Trbović D., Mijatović M., Radović Č. (2012). Fatty acid profile and cholesterol content of m. longissimus of free-range and conventionally reared Mangalitsa pigs. *S. Afr. J. Anim. Sci.*, 42: 101–113.
- Proorocu M. (2021). General characteristics of rustic pig breeds (*Sus scrofa*). *Porcine Research*, 11: 12.
- Pugliese C., Bozzi R., Campodoni G., Acciaioli A., Franci O., Gandini G. (2005). Performance of Cinta Senese pigs reared outdoors and indoors. 1. Meat and subcutaneous fat characteristics. *Meat Sci.*, 69: 459–464.

- Pugliese C., Calagna G., Chiofalo V., Moretti, V. M., Margiotta S., Franci O., Gandini G. (2004). Comparison of the performances of Nero Siciliano pigs reared indoors and outdoors: 2. Joints composition, meat and fat traits. *Meat Sci.*, 68: 523–528.
- Pugliese C., Sirtori F., Acciaoli A., Bozzi R., Campodoni G., Franci O. (2013). Quality of fresh and seasoned fat of Cinta Senese pigs as affected by fattening with chestnut. *Meat Sci.*, 93: 92–97.
- Radovic C., Petrovic M., Parunovic N., Radojkovic D., Savic R., Stanišić N., Gogic M. (2017). Carcass and pork quality traits of indigenous pure breeds (Mangalitsa, Moravka) and their crossbreds. *Indian J. Anim. Res.*, 51: 371–376.
- Real Decreto 4/2014. (2014), de 10 de enero, por el que se aprueba la norma de calidad para la carne, el jamón, la paleta y la caña de lomo ibérico. TEXTO CONSOLIDADO Última modificación: 11 de junio de 2016 LEGISLACIÓN CONSOLIDADA. 1–16.
- Robina A., Viguera J., Perez-Palacios T., Mayoral A.I., Vivo J.M., Guillen M.T., Ruiz J. (2013). Carcass and meat quality traits of Iberian pigs as affected by sex and crossbreeding with different Duroc genetic lines. *Span. J. Agric. Res.*, 11: 1057–1067.
- Rodríguez-Estévez V., García A., Peña F., Gómez A.G. (2009). Foraging of Iberian fattening pigs grazing natural pasture in the dehesa. *Livest. Sci.*, 120: 135–143.
- Rodriguez-Estévez V., Sánchez-Rodríguez M., Arce C., Garcia A.R., Perea J.M., Gómez-Castro A.G. (2012). Consumption of acorns by finishing Iberian pigs and their function in the conservation of the dehesa agroecosystem. *AF-BES*, 1–22.
- Rosenvold K., Andersen H.J. (2003). The significance of pre-slaughter stress and diet on colour and colour stability of pork. *Meat Sci.*, 63: 199–209.
- Scerra M., Foti F., Caparra P., Cilione C., Bognanno M., Paolo F., Paolo D. C., Natalello A., Musati M., Chies L. (2024). Effects of feeding bergamot pulp and olive leaves on performance and meat quality in Apulo-Calabrese pigs. *Vet. Anim. Sci.*, 23: 100336.
- Senčić D., Samac D., Steiner Z. (2013). Influence of Nutrition of Black Slavonian Pigs on the Quality of Ham and Cured Ham. *Maced. J. Anim. Sci.*, 3: 57–61.
- Serrano M.P., Valencia D.G., Fuentetaja A., Lázaro R., Mateos G.G. (2009). Influence of feed restriction and sex on growth performance and carcass and meat quality of Iberian pigs reared indoors. *J. Animal Sci.*, 87: 1676–1685.
- Santos Silva J., Pedro Araújo J., Orlando Cerqueira J., Pires P., Alves C., Batorek-Lukač N. (2019). Bísaro Pig. IntechOpen.
- Simopoulos A.P. (2002). The importance of the ratio of omega-6/omega-3 essential fatty

- acids. *Biomed. Pharmacoth.*, 56: 365–379. Simopoulos A. P. (2006). Evolutionary aspects of diet, the omega-6/omega-3 ratio and genetic variation: nutritional implications for chronic diseases. *Biomed. Pharmacoth.*, 60: 502–507.
- Simopoulos A.P. (2016). An increase in the omega-6/omega-3 fatty acid ratio increases the risk for obesity. *Nutrients*, 8(3), 128.
- Sirtori F., Crovetto A., Acciaioli A., Pugliese C., Bozzi R., Campodoni G., Franci O. (2014). Effect of dietary protein level on carcass traits and meat properties of Cinta Senese pigs. *Animal*, 8: 1987–1995. Škorput D., Stupnišek I., Špehar M., Karolyi D., Kaić A., & Luković Z. (2025). Effect of African swine fever on genetic diversity in Black Slavonian pig. *Front. Anim. Sci.*, 6: 1553271.
- Sládek L., Dračková E. (2020). The effect of genotype, sex and intramuscular fat content on the colour of pork. *Acta Univ. Agric. Silvic. Mendel. Brun.*, 68: 101–108.
- Sponenberg D.P., Martin M., Couch C., Beranger J. (2019) Conservation strategies for local breed biodiversity. *Diversity*, 11: 177.
- Stanišić N., Radović, Č., Stajić S., Živković D., Tomašević I. (2015). Physicochemical properties of meat from Mangalitsa pig breed. *XVII(1)*, 156–159.
- Suman S.P., Joseph P. (2013). Myoglobin chemistry and meat color. *Annu. Rev. Food Sci. Technol.*, 4: 79–99. Szulc K., Skrzypczak E., Buczyński, J. T., Stanisławski D., Jankowska-Mąkosza A., and Knecht D. (2012). Evaluation of fattening and slaughter performance and determination of meat quality in Złotnicka Spotted pigs and their crosses with the Duroc breed. *Czech J. Anim. Sci.*, 57: 95–107.
- Szymańska E.J., Dziwulaki M. (2022). Development of African Swine Fever in Poland. *Agruculture* 12, 119.
- Teixeira A., Silva S.R., Hasse M., Almeida J.M.H., Dias L. (2021). Intramuscular fat prediction using color and image analysis of bívaro pork breed. *Foods*, 10: 1–12.
- Tejeda J. F., Hernández-Matamoros A., Paniagua M., González E. (2020). Effect of free-range and low-protein concentrated diets on growth performance, carcass traits, and meat composition of iberian pig. *Animals*, 10: 1–13.
- Tibau J., Torrentó N., Quintanilla Aguado R., González J., Angels Oliver M., Gil M., Jaume J., Batorek-Lukač N. (2019). Negre Mallorquí (Majorcan Black) Pig. In M. Čandek-Potokar & R. Nieto, *European Local Pig Breeds - Diversity and Performance. A Study of Project TREASURE*, 1–13.
- Tomažin U., Batorek-Lukač N., Škrlep M., Prevolnik-Povše M., Čandek-Potokar M. (2018). Meat and fat quality of krškopolje pigs reared in conventional and organic production systems. *Animal*, 13: 1103–1110.
- Tomović V.M., Šević R., Jokanović M., Šojić B., Škaljac S., Tasić T., Ikonić P., Polak

- M.L., Polak T., Demšar L. (2016). Quality traits of longissimus lumborum muscle from White Mangalica, Duroc × White Mangalica and Large White pigs reared under intensive conditions and slaughtered at 150 kg live weight: A comparative study. *Arch. Anim. Breed.*, 59: 401–415.
- Torres Filho R. de A., Cazedey H.P., Fontes P.R., Ramos A. de L. S., Ramos E.M. (2017). Drip loss assessment by different analytical methods and their relationships with pork quality classification. *J. Food Qual.* Traspov A., Deng W., Kostyunin, O., Ji J., Shatokhin K., Lugovoy S., Zinovieva N., Yang B. and Huang L. (2016). Population structure and genome characterization of local pig breeds in Russia, Belorussia, Kazakhstan and Ukraine. *Genet. Sel. Evol.* 48: 16.
- Tsereniuk O.M., Vashchenko P., Khokhlov A.M., Tsybenko V.H., Shostia G.M., Saenko A.M., Peka M.Y., Zhukorskyi O.M. (2023). Comparative characteristics of polymorphisms of melanocortin 4 and ryanodine 1 receptor genes of Myrhorod pigs before and after the African swine fever outbreak. *Regul. Mech. Biosyst.*, 14: 601–608.
- Vujadinović D.P., Grujić R.D., Tomović V.M., Vukić M.S., Jokanović M.R. (2014). Cook Loss as a Function of Meat Heat Treatment and Regime. *Quality of Life (Banja Luka) - APEIRON*, 10(3–4).
- Warriss P.D., Brown S.N. (1995). The relationship between reflectance (EEL value) and colour (L*) in pork loins. *Anim. Sci.*, 61: 145–147.
- Wilkinson J.M. (2011). Re-defining efficiency of feed use by livestock. *Animal*, 5: 1014–1022.
- Wojtysiak D., Połtowicz K. (2014). Carcass quality, physico-chemical parameters, muscle fibre traits and myosin heavy chain composition of m. longissimus lumborum from Puławska and Polish Large White pigs. *Meat Sci.*, 97: 395–403.
- Wood J.D., Enser M., Fisher A.V., Nute G.R., Sheard P.R., Richardson R.I., Hughes S.I., Whittington F.M. (2008). Fat deposition, fatty acid composition and meat quality: A review. *Meat Sci.*, 78: 343–358.

Received: 21 VI 2025

Accepted: 21 X 2025




Figure 1. Methodology for literature search




Figure 2. Recommendations scheme for European native pig breeds

Table 1. Native pig breeds in selected European countries: population data and number of breeding females (FAO, 2024)

Country	Breed	Total number of animals	Number of breeding females
SOUTH-WEST EUROPE			
Italy	Cinta Senese	1 992 (update 2023)	692 (update 2023)
	Apulo-Calabrese	4 230 (update 2023)	704 (update 2023)
	Mora Romagnola	1 233 (update 2023)	259 (update 2023)
	Nero Siciliano	6 281 (update 2023)	698 (update 2023)
Spain	Iberian	3 613 123 (2024)	564 749 (2024)
	Negra Canaria	2 276 (2024)	614 (2024)
	Porc Negre	1 482 (2024)	1 372 (2024)
	Mollorquí	4 173 (2024)	433 (2024)
	Porco Celta		
France	Gasconne	1 760 (2024)	1 591 (2024)
Portugal	Alentejana	27 953 (2024)	3 639 (2024)
	Bisaro	38 772 (2024)	5 028 (2024)

	Malhado	de	3 301 (2024)	429 (2024)
	Alcobaça			
CENTRAL EUROPE				
Slovenia	Krskopolje		2 006 (2024)	296 (2024)
Poland	Pulawska		3 404 (2024)	2 488 (2024)
	Zlotnicka Spotted		1 056 (2024)	936 (2024)
	Zlotnicka White		1 214 (2024)	1 058 (2024)
Czech Republic	Prestice Black Pied		900 (2022)	394 (2022)
Belarus	Belorussian White	Large	2 958 300 (2006)	No data
Croatia	Black Slavonian pig		3 200 (2023)	2 143 (2023)
Hungary	Blond Mangalitsa		3 385 (2023)	3 332 (2023)
	Red Mangalitsa		1 542 (2023)	1 503 (2023)
Serbia	Mangalitsa		4000 (2023)	1954 (2023)
Ukraine	Mirgorod		2 757 (2019)	150 (2019)
	Poltavian Meat		1 392 (2023)	130 (2023)
	Ukrainian White	Steppe	1 513 (2023)	174 (2023)

Table 2. Fattening and slaughter parameters of European native pig breeds

Country	Breed	SW, kg	ADG, g	BF, mm last rib	BF, mm <i>gluteus medius</i>	DY, %	M, %	LMY, %	References
Italy	Cinta Senese	127-147	416	-	20-49.3	82	-	72-76	Pugliese et al. (2005), Sirtori et al. (2014)
	Apulo-Calabrese	144-157	359-413	-	-	83	-	-	Micari et al. (2009), Scerra et al. (2024)
	Mora Romagnola	193	376	57,1	62,1	80,4	-	39	Fortina et al. (2005)
	Nero Siciliano	62-121	162-600	39,1	39-49	79.1-82.5	-	40	Pugliese et al. (2004), Liotta et al. (2005), Bozzi et al. (2019)
Spain	Iberian	152-190	563	7.63-80	65	77.9-81.2	-	-	Tejeda et al. (2020), Serrano et al. (2009), Ayuso et al. (2018)
	Porc Negre Mollorquí	158	410	72	66	80,4	-	-	Tibau et al. (2019)
	Porco Celta	167.3	-	41.6-43.6	48,7	80,5	-	-	Franco et al. (2014)
France	Gasconne	~175.3	439-472	44.7-47	46.1-49	78.7-80	-	-	Mercat et al. (2019), Lebret et al. (2021)
Portugal	Alentejana	100	333-583	38.1-55.6	39,4	73.3-80.9	-	-	Freitas et al. (2007), Charneca et al. (2019), Martins et al. (2021)
	Bisaro	79	396	17- 21,1	-	75.2-76.8	-	-	Álvarez-Rodríguez Teixeira (2019), Charneca et al. (2019)
Slovenia	Krskopolje	nd	-	29	35.4-36.4	79.6-81.1	-	43-46	Čandek-Potokar et al. (2003), Tomažin et al. (2018), Lukač et al.(2019)
Poland	Pulawska	~100	638-810	23	12.73-28.86	75.33-80	49.4-56.11	-	Wojtysiak & Połtowicz (2014), Debrecéni et al. (2018), Babicz et al. (2020), Milczarek (2021)

	Zlotnicka Spotted	107	590	41,3	28.9-34.96	76.51-80.5	41.83-46.28	-	Kapelański et al. (2006), Grzeškowiak et al. (2009), Bocian et al. (2012), Szulc et al. (2012)
	Zlotnicka White	>70	-	27,7	34,7		46,3	-	Grzeškowiak et al. (2009)
Czech Republic	Prestice Black Pied Pig	94.2-115.2	~480	36.07-43.21				48.76-48.94	Nevrkla, Václavková & Rozkot (2021)
Croatia	Black Slavonian pig	~135	-	55		82	-	15-43	Butko et al. (2007), Senčić et al. (2013), Kušec et al. (2019)
	Blond Mangalitsa	140-150	369	-	-	-	-	27	Tomović et al. (2016), Nchama et al. (2023)
Hungary	Mangalica in general term	75-120	268-299	-	50.5-57	73-81.5	-	-	Parunović et al. (2012), Hoha et al. (2016), Radović et al. (2017), Debrecéni et al. (2018), Charlton et al. (2022)
Ukraine	Mirgorod	100	632	30,9	-	72,5	50,4	-	Tsereniuk et al. (2023)

SW- slaughter weight; CW – carcass weight; CCW – cold carcass weight; HCW – hot carcass weight; ADG– Average daily gain; FBW – final body weight; BF – backfat; DY – dressing yield; M – meatiness; LMY – lean meat yield; nd – no data.

* measured at 5 points, ** dressing yield in the loin, *** measured on the back, ****from the age of 90 days to slaughter.

Table 3. Physicochemical characteristics of meat from European native pig breeds

Country	Breed	SW, kg	pH ₄₅ min	pH _{24h}	L*	a*	b*	Cooking loss, %	WBSF, N, kg, kg /cm ²	Drip loss, %	References
Italy	Cinta Senese	127-147	6.2-6.4	5.82-5.78	46-50.1	11-14.9	2.9-5.4	17.9-30.3	94.14-99.87 N, 10.17 kg	0.7-2.14	Pugliese et al. (2005), Sirtori et al. (2014), Aquilani et al. (2019)
	Apulo-Calabrese	144.2	-	-	46.2-49.3	3.9-7.1	5.8-6.1	-	-	-	Caparra et al. (2023), Scerra et al. (2024)
	Nero Siciliano	62-121	6.18-6.29	-	46.7-50.1	14.7-15.3	4.9-5.8	23.5	9.99-10.11 kg	-	Pugliese et al. (2004)
	Mora Romagnola	193	6,57	6,15	42,3	8,7	2,2	-	-	-	Fortina et al. (2005)
Spanish	Iberian	152-190	-	5,62	39.8-41.7	8.7-11.3	6.4-13.6	24,1	4.33 kg/cm ²	-	Serrano et al. (2008), Fernández-Barroso et al. (2020), Tejeda et al. (2020)
	Porco Celta	167.3	-	5-46-5.55	47.3-50.7	10.4-13.1	10.7-11.7	19.7-21.9	3.64-3.8 kg/cm ²	1.8-2.7	Franco et al. (2014;2016)
France	Gasconne	~173.3	6,34	5.57-5.6	44-44.7	9.7-10.04	3.5-3.8	14,5	24.3 N	1,7	Mercat et al. (2019), Lebret et al. (2021)
Portugal	Alentejano	~100	-	5.57-5.7	49.4-50.8	10.3-15.2	3.6-8.7	17.7-26.9	26.6 N	1,3	Martins et al. (2020;2021)
	Bisaro	79	-	5,44	55.1-55.7	3.4-7.8	4-10.9	28,9	47.4 N	3,8	Martins et al. (2020), Teixeira et al. (2021)
Slovenia	Krskopolje	nd	6,33	5.6-5.71	50,4	6,6	0,9	26,9	53.2 N	4,2	Čandek-Potokar et al. (2003), Lukač et al. (2019), Tomažin et al. (2018)
Poland	Pulawska	~100	6.10-6.78	5.58-5.63	52.1-53.17	3.1-15.9	0.8-16.1	20.41-35.9	3.7-5.62 kg/cm ²	1.7-24.89	Wojtysiak & Połtowicz (2014), Bogucka & Kapelański (2016), Debrecéni et al.

											(2018), Milczarek et al. (2019), Milczarek (2021)
	Zlotnicka Spotted	107	6.38-6.57	5.56-5.67	47.9-49.3	4.77-17.2	2.5-5.04	27.46-29.36	43.61 N/cm2	2.36-3.4	Grześkowiak et al. (2009), Bogucka & Kapelański (2016), Bocian et al. (2012), Szulc et al. (2012)
	Zlotnicka White	>70	6,38	5.53-5.72	43.9-49.54	6.3-14.23	3.34-4.9	22.76-27.62	-	3,4	Grześkowiak et al. (2009), Migdał et al. (2017)
Czech Republic	Prestice Black Pied Pig	94.2-115.2	6.29	5.65	50.54	1.28	10.63			2.95	Nevrkla, Václavková & Rozkot (2021)
Croatia	Black Slavonian pig	~153.8		5,82	49,9	20	4,7	-	-	-	Karolyi et al. (2004)
Hungary	Blond Mangalitsa	110-140	6.25-6.37	5.72-5.74	40.3-54	4-11.76	5.3-11.4	18,49	43.1 N	5,9	Tomović et al. (2016), Imrich et al. (2019)
	Mangalica in general	75-120	6,1	5.47-5.64	39.1-50.1	2.7-38.2	10.4-14.6	2.7-26.9	42.6 N, 5.05 kg	4.8-8.8	Stanišić et al. (2015), Radović et al. (2017), Debrecéni et al. (2018), Charlton et al. (2022), Nchama et al. (2023)

SW - slaughter weight; L* (lightness), a* (redness), b* (yellowness); WBSF - Warner–Blatzer shear force (values given in different units depending on the publication).

Table 4. Protein, intramuscular fat content, total MUFA and PUFA fatty acids, and the n-6/n-3 ratio in LTL muscle in European native pig breeds

Country	Breed	SW, kg	Rearing conditions	Protein, %	IMF, %	n-6/n-3 PUFA	Total MUFA (% of total FA)	Total PUFA (% of total FA)	References
Italy	Cinta Senese	84	outdoor	23.09	6.66	12.5	-	-	Aquilani et al. (2019)
		130	indoor	23.03	2.32	24.4	49.9	11.2	Pugliese et al. (2013)
	Apulo-Calabrese	84.8	outdoor	20.66	1.86	8.2	-	-	Caparra et al. (2023)
		144-157	indoor	22	1.91-2.09	15.5*-2	48.8	12.4	Aboagye et al. (2020), Scerra et al. (2024)
	Nero Siciliano	62-121	outdoor	22.24	5.77	10.7	53.3	10.9	Pugliese et al. (2004)
			indoor	23.42	6.25	18.3	42.2	14.4	
	Mora Romagnola	193	indoor	23.3	6.1		47.6	11	Fortina et al. (2005)

Spanish	Iberian	152	free range traditional system	22.44	6.6	-	57.1	6.8	Tejeda et al. (2020)
		145	indoor	21.5-21.6	6.61- 8.6	17.7	52	6.5	Pardo et al. (2021), Serrano et al. (2008)
	Porco Celta	167.3	outdoor	22.65	5.22	17.9	49.6	7.6	Franco et al. (2014)
France	Gasconne	~175.3	outdoor	23.07	2.7	6.1-6.26	53.7	6.6	Mercat et al. (2019)
Portugal	Alentejano	~150	outdoor	23.6	6.9		-	-	Martins et al. (2020)
	Bisaro	~150	outdoor	23	6		-	-	
Slovenia	Krskopolje	nd	indoor	23.2	2.81	15.7	47.5	11.6	Lukač et al. (2019), Tomažin et al. (2018)
Poland	Pulawska	~100	indoor	22.72-23.3	1.9-5.5	8.89- 29.4	56.5	3.4	Debreceni et al. (2018), Milczarek et al. (2021)
	Zlotnicka Spotted	107	indoor	23.27-25.2	2.04- 2.6	10.78- 12.94	49.4	11.1	Grześkowiak et al. (2009), Debreceni et al. (2018)
	Zlotnicka White	>70	indoor	24.5	1.87	10.73	47.4	12.9	Grześkowiak et al. (2009)

Czech Republic	Prestice Pied	Blac	94.2-115.2	indoor	20.12	2.29	6.7	44.07	13.37	Nevrkla, Václavková & Rozkot (2021); Belkova et al. 2023
Croatia	Black Slavonian pig		~135	outdoor indoor	21.25 21.3	5.9 4.95				Butko et al. (2007)
	Blond Mangalitsa		110- 140	indoor	21.83-24.64	1.34-5.86				Tomović et al. (2016), Imrich et al. (2020)
Hungary	Mangalica in general		75-120	outdoor	21.45-21.7	7.57	9.2-12.97	44.8-55.5	6.5-15.6	Parunović et al. (2012), Debrecéni et al. (2018)
			75-120	indoor	19-24.1	1.9	17.92-37.3	57.2	5.9	Parunović et al. (2012), Nchama et al. (2023)

*Content of total PUFA in % of IMF.

SW – slaughter weight; nd- no data; FA – fatty acids, MUFA – monounsaturated fatty acids, PUFA – polyunsaturated fatty acids, IMF – intramuscular fat, LTL – m. longissimus thoracis et lumborum.