

# ASSESSMENT OF LANDSCAPE CONNECTIVITY OF URBAN GREEN INFRASTRUCTURE FOR BIODIVERSITY MANAGEMENT IN IBADAN

RASHEED SEKORE BUSARI<sup>1</sup> AND OLUDARE HAKEEM ADEDEJI<sup>2</sup>

<sup>1</sup>*Department of Technology Education (Building), Emmanuel Alayande University of Education, Oyo,*

<sup>2</sup>*Department of Environmental Management and Toxicology, Federal University of Agriculture, Abeokuta*

*\*Corresponding author email: sekorerasheed@gmail.com, adedejiogh@funaab.edu.ng*

**Received:** 29<sup>th</sup> June 2025, **Accepted:** 1<sup>st</sup> September 2025

## ABSTRACT

Urban Green Infrastructure (UGI) provides multiple benefits, but these are being greatly eroded by urbanization and declining connectivity, which threaten biodiversity in urban centres. This study therefore investigated the connectivity of green spaces in the Ibadan metropolis, Nigeria, using a combination of geospatial technology and landscape metrics. Landsat 5 TM, 7 ETM+, and 8 OLI-TIRS satellite imageries for 2000, 2005, 2010, 2015, and 2020 were used for land use/land cover change detection. Connectivity analysis was conducted with Fragstats metrics using seven indices: Number of Patches (NP), Mean Patch Area (AREA\_MN), Mean Area Perimeter Ratio (PARA\_MN), Mean Patch Contiguity Index (CONTIG\_MN), Largest Patch Index (LPI), Percentage of Landscape (PLAND), and Area-Weighted Mean Patch Fractal Dimension (AWMPFD). The results showed that in 2000, the built-up area covered about 70.66 km<sup>2</sup> of the total land (51.97 %), whereas green areas covered about 51.17 km<sup>2</sup> (37.64 %). By 2020, the built-up area had increased from 51.97 % in 2000 to 69.04 %, to the detriment of other land uses, which decreased drastically over the years. Non-UGI areas accounted for 62.01 % of the total area in 2000 and increased consistently to 77.04 % by 2020. There was also a continual increase in the Number of Patches (NP) in green areas. The study established that UGI connectivity in Ibadan is inadequate, with CONTIG\_MN declining from 0.41 in 2000 to 0.13 in 2020, while the Percentage of Landscape (PLAND) decreased from 10.62 in 2000 to 4.04 in 2020, indicating a high degree of fragmentation of green spaces. The findings therefore recommend halting the destruction of UGI in Ibadan to prevent further biodiversity loss and ecological decline in the city.

**Keywords:** Urban Green Infrastructure, Connectivity, Geospatial Technology, Fragstats Metrics, Fragmentation and Biodiversity

## INTRODUCTION

Urban green infrastructure (UGI) is a vital part of urban areas, consisting of vegetation with a high variability of functions, dimensions, and characteristics, such as parks, forests, community gardens, representative green spaces, street trees, green roofs and walls, and

service or marginal green areas (Fairbrass *et al.*, 2018; Zhang *et al.*, 2022; Menconi *et al.*, 2023; Hutt-Taylor *et al.*, 2024). UGI represents spatial structures that provide benefits from nature to people and aim to enhance nature's ability to deliver multiple valuable ecosystem goods and services, such as clean air and water, carbon sequestration, and mitigation of urban heat (Grover & Singh, 2015; Cheng *et al.*, 2021; Zheng & Chen, 2024). UGI has become an important component of human settlements because of the wide range of services it offers, which provide multiple benefits to human beings and other living organisms (Adegun *et al.*, 2021; Cho *et al.*, 2024; Menconi *et al.*, 2024). Engaging in outdoor recreation and visiting available UGI sites benefits public health, significantly influences attitudes and actions that promote nature conservation, and supports environmental protection (Mygind *et al.*, 2019; Moreira *et al.*, 2021; ONS, 2021; Barragan-Jason *et al.*, 2023). UGI also strengthens connections between communities and nature (Silva *et al.*, 2018; Pauleit *et al.*, 2019; Gong *et al.*, 2024; Reinwald *et al.*, 2024). It is an indispensable prerequisite for good quality of life in cities, comprising both near-natural and culturally shaped open spaces that safeguard ecosystem services for humans and enhance biodiversity (Federal Agency for Nature Conservation, 2017; Menconi *et al.*, 2021; Graffigna *et al.*, 2023). When properly planned and managed, UGI can contribute significantly to urban environmental sustainability (Meijering *et al.*, 2018; Shackleton *et al.*, 2018; Gwedla *et al.*, 2019; Russo & Cirella, 2020; Cocks & Shackleton, 2020; Gelan & Girma, 2021; Puchol-Salort *et al.*, 2021; Stan, 2022). Furthermore, studies have shown that high-quality green areas in urban spaces encourage physical activity, reduce stress, increase resilience, and improve social interactions (Sugiyama *et al.*, 2008; Roe *et al.*, 2013). Research also indicates that urban green spaces positively affect property values. To enhance residents' welfare, green infrastructure policies should therefore prioritize the provision of accessible small- and medium-sized parks or forests near residential areas (Cho *et al.*, 2024). UGI is recognized as a key strategy for combating climate change, halting biodiversity loss, and preventing many ecological crises in densely populated urban centers (Connop *et al.*, 2016; Adegun *et al.*, 2021; Lampinen *et al.*, 2023). It also provides essential ecosystem services for creating liveable cities, on which human well-being depends (Chikaeze, 2019; Menconi *et al.*, 2023; Cho *et al.*, 2024; Kulczyk *et al.*, 2024).

Ecosystem services are the functions provided by GI and natural systems, which are of benefits to society and the economy (Fairbrass *et al.*, 2018; Kulczyk *et al.*, 2024) and these include provisioning, regulating, cultural and supporting services (Hassan *et al.*, 2005; Riungu *et al.*, 2018; Arvidsen *et al.*, 2024). Worldwide, cities depend on GI for the provision of these services and UGI provides numerous ecosystem services and has contributed to increasing physical activity, improved in mental health and improved socialization of community residents and biodiversity functions (Hansen *et al.*, 2014, Moreira *et al.*, 2021; Hannah *et al.* 2023; Cho *et al.*, 2024). Cities depend on green infrastructure for the provision of biodiversity on which human survival depends (Zari, 2018; Pauleit *et al.*, 2019; Reinwald *et al.*, 2024). Biodiversity is a vital component that serves as an essential ecosystem services for liveable, healthy and well-functioning cities which are of immense benefits to the society and the economy (Fairbrass *et al.*, 2018; Scheiber, 2022). However, rapid population growth and urbanization been witnessed in cities of the world have resulted in the conversion of several urban lands into built up structures and in excessive development of the natural ecosystem (Garrah *et al.*, 2017; Areola & Ikporukpo, 2018; Weisser *et al.*, 2021; Gong *et al.*, 2024). This is a major driver of forest fragmentation and the loss of biodiversity (Severijnen, 2018; Kaczorowska & Pont, 2019; Kowarik *et al.*, 2020; Kirk *et al.*, 2021; WBCSD, 2023). The urban landscapes are experiencing drastic change due

to anthropogenic pressures, which include loss of habitats and fragmentation (Graffigna *et al.*, 2023). According to Gong *et al.* (2024) intensity of human activities in the urban area causes landscape alterations to occur rapidly presenting, the challenges of public safety and ecological sustainability that can compromise the urban development framework. Presently, report have shown that 60 % of the world's ecosystem are degraded and used unsustainably (MEA, 2005; UNFAO, 2011; WBCSD, 2023), while about 50 % of the world population live in cities at the commencement of 21<sup>st</sup> century and also 70 % of human population are expected to live in urban areas by 2050 (United Nations Human Settlement Program, 2012; United Nations, 2018; Gong *et al.*, 2024).

UGI is becoming grossly inadequate in major Nigerian cities due to rapid population growth and uncontrolled urban development. The biodiversity components and connectivity of UGI are under serious threat from anthropogenic activities (Kowarik *et al.*, 2020), resulting in fragmentation and habitat loss. In the context of landscape connectivity in urban settings, the concern is not only with increasing green cover, improving the microclimate, and enhancing other ecological functions, but also with bringing people closer to nature to promote better physical and mental health (Jim & Chen, 2003; Pascual-Hortal & Saura, 2006; Pregitzer *et al.*, 2022; Zhao *et al.*, 2023). UGI also contributes to better living environments and enhances economic values such as urban landscape quality and tourism potential (Rudnick *et al.*, 2012). Several studies based on urban morphology have demonstrated the influence of UGI distribution and patterns on various aspects, including thermal adaptation, transportation, energy use, cultural services, and ecological networks (Vance & Hedel, 2007; Wickham *et al.*, 2010; Lin *et al.*, 2024). Conducting a comprehensive inventory of a city's green spaces is an important but challenging prerequisite for improving urban planning and mitigating the ecological impacts of urban expansion (Von Thaden *et al.*, 2021). However, despite the recognized value of UGI in urban settings, its crucial role in constructing urban ecological networks is often overlooked and easily altered by urban expansion (Adegun *et al.*, 2021; Gong *et al.*, 2024). These issues call for a critical assessment of landscape connectivity in urban green infrastructure, particularly for maintaining biodiversity in sub-Saharan cities such as Ibadan, Nigeria, in order to achieve the objectives of the Sustainable Development Goals aimed at creating resilient urban areas. The loss of habitats and UGI poses threats to the physical and psychological well-being of urban residents, as well as to biodiversity itself (Hutt-Taylor *et al.*, 2024).

Landscape connectivity, defined as the ease with which organisms move between landscape elements and the number of connections between patches relative to the maximum potential connections (Brierley *et al.*, 2006; Aune *et al.*, 2011; CIEEM, 2018), faces critical threats from fragmentation and habitat loss (Rudnick *et al.*, 2012). Understanding the composition, quality, and connectivity of green spaces is key to conserving and enhancing the biodiversity assets essential for human well-being in urban environments (Aronson *et al.*, 2017; Jennings *et al.*, 2017). There are two basic types of connectivity: structural and functional (Auffret *et al.*, 2015; Ronan *et al.*, 2020). Structural connectivity refers to the physical (Euclidean) distance between patches, regardless of how organisms move across the landscape (Graffigna *et al.*, 2023). Functional connectivity, by contrast, considers movement behavior by estimating routes between patches that are suitable for dispersal. The degree of structural connectivity provided by green areas (such as public parks, private gardens, squares, and orchards) embedded in the urban matrix can offer insights into ecological functioning and the natural processes taking place there (Guo *et al.*, 2018; Graffigna *et al.*, 2023). Landscape connectivity analysis is in many cases driven by spatial data, and landscape metrics can be measured using spatial models such as Fragstats software (McGarigal, 1995). Spatial metrics are quantitative indices that describe the structure and


patterns of a landscape (Herold *et al.*, 2003). When integrated with GIS, landscape metrics become useful tools for quantifying spatial land-cover characteristics (Kong *et al.*, 2012; Rafiee *et al.*, 2009). They have been widely applied in studies of landscape variability over the years (Sun *et al.*, 2020; Liu *et al.*, 2021), addressing both ecological and visual values (Sahraoui *et al.*, 2021). Fragstats is commonly used to quantify green space patterns using landscape metrics at local scales (Kong & Nakagoshi, 2006; Han *et al.*, 2002). It has also been employed in numerous studies analyzing the distribution of green spaces, particularly in relation to ecological networks that consist mainly of green areas (Li *et al.*, 2015; Kang *et al.*, 2018; Han *et al.*, 2022). Over the years, many green areas in Nigerian cities, including Ibadan, have been lost due to changes in land use and land cover. This loss has negatively affected connectivity and reduced the effectiveness of biodiversity functions (Siehr *et al.*, 2022; Vilanova *et al.*, 2024). Therefore, this study applies Fragstats software as a spatial metric to measure the connectivity of green spaces in the Ibadan urban area, with the aim of improving biodiversity management in the city.

## MATERIALS AND METHODS

### Study Area

The city of Ibadan is located on latitude  $7^{\circ}25'$  to  $7^{\circ}42'N$  and longitude  $3^{\circ}51'$  to  $3^{\circ}9'E$  (Fig. 1), in Oyo State, southwest, Nigeria. It covers 3,080 km<sup>2</sup> with a current metro population of about 4004,000 inhabitants (Ibadan, Nigeria Population, 2024). The area is underlain by Precambrian basement complex rocks mainly of the granite, quartzite and migmatite types and also minor rock types such as pegmatite, aplite and diorite (Ajayi *et al.*, 2012).

**Fig. 1: Map of the Study Area**



It is on a central range of hills and river valley plains with an elevation of 227m above sea level (City of Ibadan, 2013). A very intensive system of rivers and streams such as R. Ogunpa, R. Ona, R. Kudeti, R. Ogbere, R. Alaro and R. Alapata drain the area (Murphy, 1998). It experiences the hot humid climate where annual average temperature ranges between 25 °C and 35°C (Alo & Nwatu, 2018), a mean annual rainfall above 1,505 mm and relative humidity between 60 % and 80 % (Raheem & Adeboyejo, 2016).

### Data and Pre-Processing

Land cover data from 2000 to 2020, collected at five-year intervals, were utilized for this study and were sourced from remote sensing monitoring data. Landsat Thematic (TM) for 2000, 2005 and 2010 and Operational Land Imager (OLI) for 2015 and 2020 were obtained from United State Geological Survey Global Visualization Viewers (USGS-GLOVIS). The dataset has a spatial resolution of 30 m and has been widely applied and validated in different scientific research (Zhang *et al.*, 2023; Gong *et al.*, 2024). Before the LULC classification the imageries has undergone preprocessing such as atmospheric correction and orthorectification. Geometric corrections of collected images were done by Google Earth Images. The Landsat data were pre-referenced with Universal Transverse Mercator (UTM) projection system and WGS-84 datum. The LULC types were categorized into four classes: built-up, green area, bare land, and water bodies (Table 1). The location of LULCs were based on high-resolution images of Google Earth, and then confirmed by ground truthing. The ArcGIS 10.5 software was used for processing the data.

**Table 1: Description of land use classes**

| Serial No | Land Use Class          | Description                                                                                                                                                              |
|-----------|-------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| A         | Built up area/Transport | Built up areas used for residential, commercial, manufacturing, institutions, transportation, communities, and utilities.                                                |
| B         | Green area              | Areas covered with soft landscape elements such as urban forests, woodlands, trees, bush, shrub, grasslands, parks, gardens, greenways, playgrounds, sports fields, etc. |
| C         | Bare surface            | These are bare lands, quarry sites, open spaces, etc.                                                                                                                    |
| D         | Waterbodies             | These are water features such as rivers, streams, lakes, ponds, fountains, etc.                                                                                          |

Source: Adapted from Anderson *et al.* (1976)

### Land use/land cover classification

For the 2000, 2005 and 2010, Landsat TM data bands no. 1–5 and 7 were used for land use mapping, while for the Landsat OLI data 2015 and 2020, bands no. 1–7 were used. The supervised image classification technique with maximum likelihood algorithm was used to show the changes of land use/land cover over time in the study area. The accuracy of the classification was done using an error matrix i.e. the Kappa coefficient (K) (Foody, 1992) for land use map of all four years under consideration. The Kappa index which indicates the presence of misclassification. The Kappa coefficient in this study ranged from 0.72-0.83 representing an accurate classification (Landis & Koch, 1977).

### Landscape Patterns Analysis

Based on the existing published research and the knowledge of the study area seven landscape metrics were to capture the pattern of UGI within the Ibadan Metropolitan Area: Number of patches into (NP), mean patch area (AREA\_MN), mean area perimeter ratio (PARA\_MN), mean patch contiguity index (CONTIG\_MN), largest patch index (LPI),

percentage of landscape (PLAND) and area weighted mean patch area (AWMPFD). The land use land cover analysis obtained from geographic information system analysis served as input into the Fragstats software for the production of connectivity analysis.

**Table 2: Introduction and definition of landscape metrics**

| Abbreviation | Name                          | Definition                                                                                                                                       | Formula                                                                                                                            | Range                         | Unit |
|--------------|-------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------|-------------------------------|------|
| NP           | Number of patches             | Number of patches in the landscape, or total number of patches for a particular class                                                            | $NP = n_i$                                                                                                                         | >0                            | -    |
| AREA_MN      | Mean patch area               | The mean area of patches of a certain class                                                                                                      | $\frac{\sum_{i=1}^m \sum_{j=1}^n x_{ij}}{n_i}$                                                                                     | >0                            | Ha   |
| PARA_MN      | Mean area perimeter ratio     | The simple measure of shape complexity, but without standardization to a simple Euclidean shape                                                  | $PARA = \frac{P_{ij}}{a_{ij}}$                                                                                                     | >0                            | -    |
| CONTIG_MN    | Mean patch contiguity index   | Mean contiguity in patches per class                                                                                                             | $\left[ \frac{\frac{\sum_{r=1}^x c_{ijr}}{a_{ij}}}{\vartheta - 1} \right]$                                                         | $0 \leq \text{CONTIG} \leq 1$ | -    |
| LPI          | Largest patch index           | The area of the most giant patch divided by the total area of the landscape multiplied by 100                                                    | $\frac{A^2}{\sum_{i=1}^m \sum_{j=1}^n a_{ij}^2}$                                                                                   | $0 < \text{LPI} \leq 100$     | %    |
| PLAND        | Percentage of landscape       | Percentage of the area of each class                                                                                                             | $PLAND = P_1 = \frac{\sum_{j=1}^n a_{ij}}{A} (100)$                                                                                | $0 < \text{PLAND} \leq 100$   | %    |
| AWMPFD       | Area weighted mean patch area | The fractal dimension of a patch equals two times the logarithm of patch perimeter (m) divided by the logarithm of patch area (m <sup>2</sup> ). | $AWMPFD = \sum_{i=1}^m \sum_{j=1}^n \left[ \left( \frac{21n(.25p_{ij})}{1na_{ij}} \right) \left( \frac{a_{ij}}{A} \right) \right]$ | $1 \leq \text{AWMPFD} \leq 2$ | None |

## RESULTS AND DISCUSSION

### Land use/land cover change

The results of the classification of land use for year 2000 to 2020 as extracted using geospatial techniques are presented in Fig. 2 (Busari & Adedeji, 2024). During the study period significant changes were witnessed in the proportion and pattern of land use/land cover. For instance in year 2000, Ibadan built-up area covered about 70.66 km<sup>2</sup> of the total land amounting to 51.97 %, whereas green area covered about 51.17 km<sup>2</sup> (37.64 %). At the

same period, bare surface was about 13.64 km<sup>2</sup> (10.04 %), while water bodies covered about 0.49 km<sup>2</sup> (0.36 %). Rapid population growth and urban expansion brings about an increase in land use type such as built-up area for housing, industrialization, recreation, education, and transportation among others. Thus by 2020, the built-up area increased from 51.97 % in the year 2000 to 69.04 % at the detriment of other land uses that decreased drastically over the years (Table 3, Fig. 2). Erstwhile green areas like Samonda, Alalubosa, and Bower Tower downslope were converted into residential and commercial properties. These findings were corroborated by studies such as Barlow *et al.* (2018) and Akingbade *et al.* (2022) that urbanization has led to the degradation and consequent alteration of ecosystem which is responsible for environmental deterioration of the greens.

**Table 3: Land use/Land cover change 2000-2020**

| Year         | 2000                    |          | 2005                    |          | 2010                    |          | 2015                    |          | 2020                    |          |
|--------------|-------------------------|----------|-------------------------|----------|-------------------------|----------|-------------------------|----------|-------------------------|----------|
| Land use     | Area (km <sup>2</sup> ) | Area (%) | Area (km <sup>2</sup> ) | Area (%) | Area (km <sup>2</sup> ) | Area (%) | Area (km <sup>2</sup> ) | Area (%) | Area (km <sup>2</sup> ) | Area (%) |
| Built-up     | 70.66                   | 51.97    | 75.85                   | 55.79    | 77.50                   | 57.00    | 91.27                   | 67.13    | 93.87                   | 69.04    |
| Green Area   | 51.17                   | 37.64    | 47.53                   | 34.97    | 48.00                   | 35.30    | 31.56                   | 21.96    | 31.07                   | 22.84    |
| Bare Surface | 13.64                   | 10.04    | 12.15                   | 8.94     | 12.41                   | 9.13     | 12.84                   | 10.69    | 10.91                   | 8.03     |
| Water bodies | 0.49                    | 0.36     | 0.42                    | 0.31     | 0.40                    | 0.30     | 0.30                    | 0.22     | 0.12                    | 0.09     |
| TOTAL        | 135.96                  | 100.00   | 135.96                  | 100.00   | 135.96                  | 100.00   | 135.96                  | 100.00   | 135.96                  | 100.00   |

### Changes in the UGI Based on Land Use


Urban expansion generates different landscape patterns through the interaction of local ecosystems and land-use decisions (Richards & Belcher, 2020). A curious look at the results of the land use/land cover change analysis show that UGI areas consistently occupied about half of the total study area since 2000 increasing from 70.66 sqkm to 93.87 sqkm in 2020 (Fig. 2). The non-UGI accounted for 62.14 % of the total area and consistently increases over the study period to 77.48 % by 2020. This is an indication that UGI in the study area has been markedly altered by urban expansion and other anthropogenic activities. During study period, UGI decreased between 2000 and 2005, increased between 2005 and 2010 and later decreased between 2005 to 2020 (Table 3). The UGI diminished drastically from 51.17 sqkm to 31.07 sqkm which is significant in proportion from 20 years prior by 2020.

### Landscape Pattern


Land cover maps at the class level were used to calculate landscape metrics in the FRAGSTATS software. These landscape metrics according to Gong *et al.* (2024) indicated a trend towards reduced heterogeneity and decreasing diversity of UGI at the landscape scale. Table 4 displays the results of landscape metrics over the past 20 years as well as a five-year forecast. The highest rate of urban growth is observed between 2010 and 2015 with an increase in the built up area of about 13.77 % within 5 years (Table 4). In the built-up areas, the number of patches (NP) declined from 1143 in 2000 to 349 in 2020. The anticipated NP metrics in 2025 will be 177 showing further decline. The decrease in the number of built-up patches indicates that disjointed patches will progressively come together to create single patches. This is a reflection of the built-up land cover type's expansion and development in the research area. Yang & Liu (2005) have shown that landscape pattern is more fragmented around city centres and along coastlines, where urbanization and human

economic activities are more concentrated. During the research period, NP in bare land decreased from 3097 in 2000 to 1284 in 2020. Moreover, NP of the green area class increased as the year progress, which is an indication of more fragmentation and loss of vegetation. In the green area, the NP measure increase from 1991 in 2000 to 2844 in 2020 and is expected to increase to 2912 by 2025. The continuous rise of number of patches (NP) in the green area could be an indication of fragmented and heterogeneous process of urban growth taking place in the study area (Olayiwola & Fakayode, 2019). At the same time there is the continuing development of scattered and fragmented urban patches in the study area due to a development of small and irregular built up patches around the periphery of the city and in peri-urban regions.


**Fig. 2: Land use analysis between 2000-2020**




**Land use for year 2000**



**Land use for year 2005**




**Land use for year 2010**



**Land use for year 2015**





**Land use for year 2020**

It is important to note that some of the findings of this study are consistent with the findings of Saeedal & Aysan (2020). Adewale & Olayinka (2019) carried out a landscape metrics analysis of land use patterns and changes in suburban local government areas of Ibadan, Nigeria. Though their landscape metrics of interest were slightly different from the ones examined in this study, there were similarities in few common metrics as their study revealed that the NP in 2017 was 809 for the built-up areas, whereas in this study, the NP of built-up area in 2015, the closest study year to 2017, was 898 (Table 4). Disruptors may have separated the patches of green areas into smaller patches as a result of their effect.

The development of residential spaces and the extension of the city, including green area and bare land, are indicated by the increase in the number of patches of green area and the reduction in built-up patches. Furthermore, in the last decades the urban expansion rate has been considerably higher exceeding what is needed to sustain population growth (Liu *et al.* 2020) and this is generating principally negative impacts on biodiversity. This demand for urban construction has placed growing pressure on biodiversity conservation (Guo *et al.*, 2018; Hyseni *et al.*, 2021).

One of the indicators that showed the mean patch size was AREA-MN. The level of interspersions is indicated by the lowering of this metric. The built-up class has seen a rise in this statistic, whereas greenspace and bare land have seen a drop. The level of the metric for the built-up area in 2000 was 7.42 and 12.90 in 2020. The decline of this measure in the green and bare land areas indicated a drop in the mean of these patch types, indicating a drop in the level of interspersions classes. In the year 2000, the average patch size in the green area class was 14.73 which declined to 6.17 in 2020.

The mean area perimeter ratio (PARA-MN) is a basic measure of the shape's complexity because it displays the circumference to area ratio. The closer the shape is to the rotating state, the smaller the ratio. As a result, based on the numbers of the metric in the classes, it was discovered that there is a marked decrease in the green area and built-up classes while bare land increased. The CONTIG-MN measure is one of the metrics for determining the degree of landscape patch integrity as well as the degree of landscape fragmentation. When the patches are entirely distributed and fragmented, the metric is 0. Contiguity is stated to be at its highest level when the landscape is formed by only one type of patch. Between 2000 and 2020, the built-up and bare land classes both increased. Green areas, on the other hand, decreased from 2000 (0.41) to 2020 (0.13), demonstrating increased contiguity and decreased interspersions in these classifications.

**Table 4: Land classes based on landscape metrics**

| Metrics   | LULC       | 2000    | 2005    | 2010    | 2015    | 2020    | 2025    |
|-----------|------------|---------|---------|---------|---------|---------|---------|
| NP        | Built-Up   | 1143    | 1043    | 902     | 898     | 349     | 177     |
|           | Bare land  | 3097    | 2975    | 2702    | 1793    | 1284    | 1007    |
|           | Green Area | 1991    | 2312    | 2505    | 2692    | 2844    | 2912    |
| AREA_MN   | Built-Up   | 7.42    | 8.47    | 9.59    | 11.45   | 12.90   | 14.68   |
|           | Bare land  | 4.23    | 3.43    | 2.39    | 1.38    | 0.98    | 0.45    |
|           | Green Area | 14.73   | 12.73   | 10.45   | 8.45    | 6.17    | 4.70    |
| PARA-MN   | Built-Up   | 1215.29 | 1015.23 | 1002.68 | 987.79  | 876.03  | 748.55  |
|           | Bare land  | 1195.68 | 1278.68 | 1320.21 | 1504.39 | 1799.49 | 1873.46 |
|           | Green Area | 1242.11 | 1142.12 | 1053.55 | 1010.23 | 968.17  | 742.65  |
| CONTIG-MN | Built-Up   | 0.12    | 0.34    | 0.65    | 0.76    | 0.79    | 0.85    |
|           | Bare land  | 0.14    | 0.17    | 0.19    | 0.20    | 0.24    | 0.25    |
|           | Green Area | 0.41    | 0.24    | 0.19    | 0.15    | 0.13    | 0.11    |
| LPI       | Built-Up   | 59.06   | 68.05   | 78.00   | 82.03   | 85.82   | 91.64   |
|           | Bare land  | 17.14   | 13.51   | 8.06    | 3.08    | 2.14    | 1.54    |
|           | Green Area | 13.75   | 11.61   | 9.35    | 4.17    | 2.14    | 1.17    |
| PLAND     | Built-Up   | 62.36   | 67.36   | 83.78   | 97.79   | 96.04   | 98.31   |
|           | Bare land  | 27.02   | 29.02   | 36.23   | 48.94   | 66.92   | 69.26   |
|           | Green Area | 10.62   | 9.62    | 7.22    | 6.27    | 4.04    | 2.46    |
| AWMPFD    | Built-Up   | 1.42    | 1.33    | 1.33    | 1.33    | 1.29    | 1.23    |
|           | Bare land  | 1.32    | 1.31    | 1.13    | 1.12    | 1.28    | 1.17    |
|           | Green Area | 1.24    | 1.12    | 1.31    | 1.31    | 1.15    | 1.21    |

The LPI index shows which patch is the largest in total. According to this indicator, if the largest patch class shrinks, it could be a factor in land fragmentation. In this study, in 2000, the green area class was 13.75 but decline in 2005 to 11.61, the further to 9.35 in 2010 and then drastically to 2.14 in 2020. The amount of the measure in the bare land class declined from 17.14 in 2000 to 2.14 in 2020. In the built-up class, this index increased from 59.06 in 2000 to 85.82 in 2020, indicating a relationship between the patches that created a larger patch. The increasing intensity of buildings in the area increased the LPI of the built-up area leading to a tremendous urbanization pressure over the other land uses (Akintunde *et al.*, 2019). This is aggravated the demand for more houses due to the increasing population, consequently leading to the creation new built-up areas at the urban peripheral areas. Generally in urban landscapes, vegetation patches usually decrease in size and quality along a gradient from periurban rural areas toward the urban center (Williams & Newbold, 2020; Von Thaden *et al.*, 2021). The class percentage of the entire landscape is represented by the PLAND measure. The expansion of both land covers is shown by an increase in this metric in both the built-up and bare land classes. This statistic accounted for 62.36 percent of the built-up class in 2000. In 2020, it was 96.04. The reduction in class fragmentation and the establishment of new built-up regions are represented by the growth in the number and density of built-up patches. During the study period, the percentage of the landscape covered by the green area class declined due to the consumption of the green ecosystem by other land uses and land covers. 10.62, 9.62, 7.22, 6.27, and 4.04 were the scores, respectively. In 2000, 2005, 2010, 2015, and 2020, the outcomes of this metric on the bare land landscape were 27.02, 29.02, 36.23, 48.94, and 66.92, respectively. Over the same period, AWMPFD results across the terrain did not change considerably during the study period. AWMPFD in the

built-up area was 1.42 in 2000 and 1.23 in 2020 while in the green areas; AWMPFD was 1.24 in 2000 and 1.15 in 2020. Higher values were however obtained 2010 and 2015 (1.31). The higher value obtained in during the period shows that the geometry of urban patches is getting more complex over time and the prevalence of fragmented low density development in the fringe areas (Wang *et al.* 2019; Akintunde *et al.*, 2019) having effects on biodiversity. The degree to which the landscape facilitates movement among landscape tracts defines the amount of connectivity (Lepczyk *et al.*, 2017; Ronan *et al.*, 2020).

The landscape connectivity of Ibadan has been badly affected by anthropogenic forces which continues to reduce the green infrastructure available in the city for various ecosystem services and benefits of humans. The ecological linkage in Ibadan remains weak and fragmented, and this connectivity reduces the effectiveness in sustaining biodiversity, restricting species movement and ecosystem resilience. A coherent, ecologically integrated green infrastructure network is imperative for sustaining biodiversity and promoting environmental resilience in Ibadan's rapidly evolving urban landscape.

## CONCLUSION

This study assessed landscape connectivity of urban green infrastructure for biodiversity management in Ibadan, Nigeria. Seven indices of the spatial metrics were used in evaluating the connectivity of these green spaces. Landscape connectivity is low in the city and could be attributed to biodiversity loss caused by anthropogenic activities. It was established that green areas continued to decrease in size from 2000 to 2020 thus leading to reduction in patch sizes and increase in fragmentation. Findings indicate that the number of patches (NP) in the green areas has continued to increase due to urban expansion; there is a noticeable trend towards a reduction in the number of patches in other landscape types. In fact the internal composition of UGI has undergone multifarious spatiotemporal changes during over the study period. This empirical evidence reviewed that existing green spaces in the city need to be effectively managed to halt further degradation of the biodiversity that provides ecosystem services on which human life and organisms well-being depend. Future planning should not only focus on preventing the loss of UGI area but also emphasizes the internal structure and characteristics of UGI. This study have provided valuable insights into how to plan UGI in a rapidly growing city like Ibadan and would help planners and administrators develop a crucial strategy for urban centres to address ecological crises. Therefore, to avoid these ecological crises and reduce unsustainable city expansion, the UGI in urban centres should received significant attention from managers and landscape designers. Efforts should be made towards planting and preserving urban trees, particularly large-stature trees that will address climate change by cooling the environment via transpiration, shading buildings and paved surfaces to reduce energy usage, and storing carbon with high permanence. Furthermore, the negative impact of urbanization on biodiversity can be buffered by blue (e.g., rivers, ponds) and green (e.g., parks, forests) spaces which should be connected by corridors, so that organisms may disperse between sites

## CONFLICT OF INTEREST

The authors declare that they have no competing interests.

## REFERENCES

- Adegun, O.B., Ikudayisi, A.E., Morakinyo, T.E. and Olusoga, O.O. (2021). Urban green infrastructure in Nigeria: A review. *Scientific African*, 14, e01044 <https://doi.org/10.1016/j.sciaf.2021.e01044>.
- Adewale, M.O. and Olayinka, F. (2019). Landscape metrics analysis of land use patterns and changes in suburban local government areas of Ibadan, Nigeria. *South African Journal of Geomatics*, 8: 2.
- Agbelade, J., Aladesanmi, D., Onyekwelu, J.C. and Apogbona, O. (2016). Assessment of urban forest tree species population and diversity in Ibadan, Nigeria. *Environment and Ecology*, 4, 185-192.
- Ajayi, O., Wahab, B., Gbadegesin, M., Taiwo, D., Kolawole, M., Muili, A. And Shiji, F. (2012). Flood Management in an urban setting: A case study of Ibadan metropolis. *Hydrology of Disaster Management*. A Special Edition. 65-81.
- Akintunde, J.A. (2019). Spatial pattern of urban growth using remote sensing and landscape metrics. *Journal of Geomatics*, 13(1), 53-60.
- Alo, A.A. and Nwatu, J.U. (2018). Modelling urban green space dynamics and associated proximate drivers in Ibadan metropolis, Ibadan, Nigeria. *Forest and Forest Products Journal*, 18, 23- 34.
- Anderson, J.R., Hardy, E.E., Roach, J.T. and Nitmar, R.E. (1976). *A land use land cover classification system for use with Remote sensing data*. Geological Survey Professional Paper No. 946. US Government Printing House, Washington DC, 28.
- Areola, A. A. and Ikporukpo, C.O. (2018). Social ecology and urban greenspaces in Ibadan, Nigeria. *Journal of Applied Science and Environmental Management* 22(7): 1111 – 1120.
- Aronson, M.F., Lepczyk, C.A., Evans, K.L., Goddard, M.A., Lerman, S.B., MacIvor, J.S., Nilon, C.H., Vargo, T., (2017). Biodiversity in the city: key challenges for urban green space management. *Frontiers Ecology and Environment*, 15 (4), 189–196. <https://doi.org/10.1002/fee.1480>.
- Auffret, A.G., Plue, J., Cousins, S.A.O., (2015). The spatial and temporal components of functional connectivity in fragmented landscapes. *AMBIO* 44 (1), 51–59. <https://doi.org/10.1007/s13280-014-0588-6>.
- Aune, K., Beier, P., Hilty, J. and Shilling, F. (2011). *Assessment and planning ecological connectivity*. The Wildlife Conservation Society, Bozeman. 82pp.
- Barnosky, A.D., Matzke, N., Tomiya, S., Guinevere, O.U.W., Swartz, B., Quental, T.B., Marshall, C., McGuire, S.L., Lindsey, E.L., Maguire, K.C., Mersey, B. and Ferrer, E.A. (2011). Has the earth's sixth mass extinction already arrived? *Research Review Nature*, 471, 51-57. Doi.1038/nature09678.
- Barragan-Jason, G., Loreau, M., De Mazancourt, C., Singer, M.C., Parmesan, C., (2023). Psychological and physical connections with nature improve both human well-being and nature conservation: a systematic review of meta-analyses. *Biol. Conserv.*, 277, 109842.
- Bennet, G. and Mulongoy, K. (2006). *Review of experience with ecological networks, corridors and buffer zones*. CBO Technical Series No 23. Secretariat of the Convention on Biological Diversity, Montreal. 103pp.
- Bennett, G. (2004). *Integrating biodiversity conservation and sustainable use: Lessons learned from ecological networks*. IUCN, Gland, Switzerland and Cambridge. 66pp.

- Brierley, G. J., Fryirs, K. and Jain, V. (2006). Landscape connectivity: The geographic basis of geomorphic application. *Area*, 38: 165–174.
- Busari, R.S. and Adedeji, O.H. (2024). Green infrastructure as a strategy for building resilience against Stormwater in Ibadan. *Journal of Building Design and Environment*, 3(2) DOI:10.37155/2811-0730-0302-12.
- Cheng, X., Damme, S.V. and Uyttenhove, P. (2021). A review of empirical studies on cultural ecosystem in urban green infrastructure. *Journal of Environmental Management*, 293(112895) DOI:10.1016/j.jenvman.2021.112895.
- Chikaeze, U. (2019). *Green Infrastructure, a conceptual approach for mitigating and adapting climate change in Nigeria cities: Case study of Lagos*. Technical Report for the Research Gate. <https://www.researchgate.net>3307>.
- Cho, C.J., Cheon, K. and Kang, W. (2024). Assessment of the Spatial Variation of the Economic Benefits of Urban Green Spaces in a Highly Urbanized Area. *Land*, 13, 577. <https://doi.org/10.3390/land13050577>
- CIEEM, (2018). *Guidelines for ecological impact assessment in the UK and Ireland: Terrestrial, marine, coastal and marine*. Winchester, UK: Chattered Institute of Ecology and environmental Management <https://cieem.net>2019/02>.
- Cocks, M.L., and Shackleton, C.M. (2020). Situating biocultural relations in city and townscapes: Conclusion and recommendations. In M. L. Cocks and C. M. Shackleton (eds.): *Diversities of nature in cities: enriching belonging, wellbeing and bioculture* (pp. 241–268) (Earthscan/Routledge).
- Connop, S., Vandergert, P., Eisenberg, B., Collier, M.J., Nash, C., Clough, J. and Newport, D. (2016). Renaturing Cities Using a Regionally-Focused Biodiversity-Led Multifunctional Benefits Approach to Urban Green Infrastructure. *Environ. Sci. Policy*, 62, 99–111.
- Fairbrass, A., Jones, K. and McIntosh, A. (2018). *Green Infrastructure for London: A review of the evidence*. The Engineering Exchange, Science of the Environment, Natural Environment Research Council, London's Global University. <https://www.ucl.ac.uk>flies>.
- Federal Agency for Nature Conservation (2017). *Federal green infrastructure concepts*. Nature Conservation Foundations for Plans Adopted by the German Federation [www.bfn.de/bkgi.html](http://www.bfn.de/bkgi.html)
- Foody, G.M. (1992). On the compensation for chance agreement in image classification accuracy assessment. *Photogrammetry, Engineering and Remote Sensing*, 58, 1459–1460.
- Garrah, J., Berton, K. and Chen, S. (2017). Urban Biodiversity through sustainable architecture and urban planning. *McGill Science Undergraduate Research Journal*, 12(1), 25-28.
- Gelan, E. and Girma, Y. (2021). Urban green infrastructure accessibility for the achievement of SDG 11 in rapidly urbanizing cities of Ethiopia. *GeoJournal*, 87(2):2883-2902.
- Gong, C., Pang, H., Olhnuud, A., Hao, F. and Lyu, F. (2024). Green Infrastructure Fluctuations in Urban Agglomeration of Shanxi Province, China: Implications for Controlling Ecological Crises. *Land*, 600. <https://doi.org/10.3390/land13050600>.
- Graffigna, S., Gonzalez-Vaquero, R.A., Torretta, J.P. and Marrero, H.J. (2023). Importance of urban green areas' connectivity for the conservation of pollinators. *Urban Ecosystems*, <https://doi.org/10.1007/s11252-023-01457-2>.
- Graham, B. and Wit, P. (2001). *The development and application of ecological networks: A review of proposals, plans and program*. Advice and Research for Development and Environment. The World Conservation Union, IUCN. AIDE, Netherlands. 1-15pp.

- Grover, A. and Singh, R. (2015). Analysis of Urban Heat Island (UHI) in Relation to Normalized Difference Vegetation Index (NDVI): A Comparative Study of Delhi and Mumbai. *Environments*, 2, 125–138.
- Guo, S., Saito, K., Yin, W and Su, C. (2018). Landscape Connectivity as a Tool in Green Space Evaluation and Optimization of the Haidan District, Beijing. *Sustainability*, 10, 1979. doi:10.3390/su10061979.
- Gwedla, N., and Shackleton, C.M. (2019). Perceptions and preferences for urban trees across multiple socio-economic contexts in the Eastern Cape, South Africa. *Landscape and Urban Planning*, 189: 225–234.
- Hall, D.M., Camilo, G.R. and Tonietto, R.K. et al (2017). The city as a refuge for insect pollinators. *Conserv. Biol.*, 31:24–29. <https://doi.org/10.1111/cobi.12840>.
- Han, Y., Yang, D., Zhang, Y., Cao, L. (2022). Optimizing Urban Green Space Layouts for Stormwater Runoff Treatment in Residential Areas: A Case Study in Tianjin, China. *Water*, 14: 2719.
- Hanna, E., Bruno, D. and Comin, F.A. (2023). The ecosystem services supplied by urban green infrastructure depends on their naturalness, functionality and imperviousness. *Urban Ecosystem*, 27, 187- 202 DOI: 10.1007/s11252-023-01442-9.
- Hansen, R., Rall, E. L. and Pauleit, S. (2014). A transatlantic lens on green infrastructure planning and ecosystem services: Assessing implementation in Berlin and Seattle. In D. Czechowski, T. Hauck and G. Hausladen, G. 2017. *Revising green infrastructure: Concept between nature and design* (pp. 247-265). Taylor P. Francis, London. DOI: 10.1201/b17639-17
- Herold, M., Goldslein, N.C. and Clarlie, K.C. (2003). The Spatiotemporal form of urban growth: Measurement, analysis and modelling. *Remote sensing of Environment* 86: 286 – 302.
- Hutt-Taylor, K., Bassett, C.G., Kinnunen, R.P., Frei, B and Ziter, C.D. (2024). Existing evidence on the effect of urban forest management in carbon solutions and avian conservation: a systematic literature map. *Environmental Evidence*, 13:23 <https://doi.org/10.1186/s13750-024-00344-3>.
- Hyseni, C., Heino, J., Bini, L.M., Bjelke, U. and Johansson, F. (2021). The importance of blue and green landscape connectivity for biodiversity in urban ponds. *Basic and Applied Ecology* 57,129-145 <https://doi.org/10.1016/j.baae.2021.10.004>.
- Jennings, V., Floyd, M.F., Shanahan, D., Coutts, C., Sinykin, A., (2017). Emerging issues in urban ecology: implications for research, social justice, human health, and well-being. *Population Environment*, 39 (1), 69–86. <https://doi.org/10.1007/s11111-017-0276-0>.
- Jim, C.Y. and Chen, S.S. (2003). Comprehensive greenspace planning based on landscape ecology principles in compact Nanjing city, China. *Landscape and Urban Planning*, 65, 95–116.
- Kaczorowska, A. and Pont, M.B. (2019). Modelling urban environment to promote ecosystem services and biodiversity: case study of Stockholm. *International Journal of E-planning Research*, 8(3), 1-12.
- Kang, J.E., Choi, H.S., Hwang, H.S. and Lee, S. (2018). Analysis of ecological network according to invalidation of decision on urban parks: Focused on Busan. *Journal of Environmental Impact Assessments*, 27: 618–634.
- Kirk, H., Garrard, G.E., Croeser, T., Backstron, A., Berthon, K., Furlong, C. Hurley, J.,

- Thomas, F., Webb, A. and Bekessy, S.A. (2021). Building biodiversity into the urban fabric: A case study in applying biodiversity sensitive urban design. *Urban Forestry and urban Greening*, 62, 1-14.
- Kong, F. and Nakagoshi, N. (2006). Spatial-temporal gradient analysis of urban green spaces in Jinan, China. *Landscape and Urban Planning*, 78: 147–164.
- Kong, F., Yin, H., Nakagoshi, N. and James, P. (2012). Simulating urban growth processes incorporating a potential model with spatial metrics. *Ecological Indicators*, 20: 82–91.
- Kowarik, I., Kendal, D. and Fischer, L.K. (2020). Biodiversity conservation and sustainable urban development. *Sustainability*, 12, 4964. doi:10.3390/su/12124964.
- Kulczyk, S. Grzyb, T., Wozniak, E andDerek, M. (2024). Nature in urban green spaces: Main attractor or nice background? Drivers and dynamics of cultural ecosystem services provision. *Urban Forestry and Urban Greening*, 9(128328) <https://doi.org/10.1016/j.ufug.2024.128328>.
- La Point, S., Balkenhol, N., Hale, J., Sadler J. and Van der Ree, R. (2015). Ecology of organisms in urban environment. *Functional Ecology*, 29, 868-878.
- Lampinen, J., García-Antúnez, O., Lechner, A.M., Stahl Olafsson, A., Gulsrud, N.M. and Raymond, C.M. (2023). Mapping Public Support for Urban Green Infrastructure Policies across the Biodiversity-Climate-Society -Nexus. *Landscape and Urban Planning*, 239, 104856.
- Landis, J.R. and Koch, G.G. (1977). An application of hierarchical kappa-type statistics in the assessment of majority agreement among multiple observers. *Biometrics*, 363–74.
- Lepczyk, C.A., Aronson, M.F.J., Evans, K.L., Goddard, M.A., Lerman, S.B., MacIvor, J.S., (2017). Biodiversity in the city: fundamental questions for understanding the ecology of urban green spaces for biodiversity conservation. *BioScience*, 67 (9), 799–807. <https://doi.org/10.1093/biosci/bix079>.
- Li, H., Chen, W. and He, W. (2015). Planning of green space ecological network in urban areas: An example of Nanchang, China. *International Journal Environmental Resources and Public Health*, 12, 12889–12904.
- Lin, J., Wei, K. and Guan, Z. (2024). Exploring the Connection between Morphological Characteristic of Built-up Areas and Surface Heat Islands Based on MSPA. *Urban Clim.*, 53, 101764.
- Lindermayer, D.B., Cunningham, R.B. and Fisher, J. (2005). Native Vegetation cover thresholds and species responses. *Biological Conservation*, 124(3): 311-316.
- Liu, X., Huang, Y., Xu, X., Li, X. et al. (2020). High-spatiotemporal-resolution mapping of global urban change from 1985 to 2015. *Nat Sustain.*, 3:564–570. <https://doi.org/10.1038/s41893-020-0521-x>.
- Liu, M., Li, X., Song, D. and Zhai, H. (2021). Evaluation and Monitoring of Urban Public greenspace Planning Using Landscape Metrics in Kunming. *Sustainability* 13: 3704.
- McDonald, R.I., Mansur, A.V., Ascensao, F., Crossman, K., Elmqvist, T., Gonzalez, A., Guneralp, B., Haase, D., Hamann, M. and Huler, O. (2020). Research Gaps in knowledge of the impact of urban growth on biodiversity. *Natural Sustainability*, 3, 16-24.
- McGarigal, K. (1995). *FRAGSTATS: Spatial Pattern Analysis Program for Quantifying Landscape Structure*; U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: Washington, DC, USA.
- Meijering, J.V., Tobi, H. and Kern, K. (2018). Defining and measuring urban sustainability

in Europe: A Delphi study on identifying its most relevant components. *Ecological Indicators*, 90: 38–46.

Menconi, M., Abbate, R., Simone, L. and Grohmann, D. (2023). Urban Green System Planning Insights for a Spatialized Balance between PM10 Dust Retention Capacity of Trees and Urban Vehicular PM10 Emissions. *Sustainability*, 15, 5888.

Menconi, M.E., Bonciarelli, L. and Grohmann, D. (2024). Assessment of the Potential Contribution of the Urban Green System to the Carbon Balance of Cities. *Environments*, 11, 98. <https://doi.org/10.3390/environments11050098>.

Menconi, M.E., Sipone, A. and Grohmann, D. (2021). Complex Systems Thinking Approach to Urban Greenery to Provide Community-Tailored Solutions and Enhance the Provision of Cultural Ecosystem Services. *Sustainability*, 13, 11787.

Millenium Ecosystem Assessment (MEA), (2005). *Ecosystems and well-being: Synthesis*. Island Press, Washington DC. <https://www.millenniumassessment.org>.

Milwaukee Metropolitan Sewage District, (2018). *Using Green Infrastructure to enhance urban biodiversity in the MMSD planning Area*. Milwaukee, WI: Milwaukee Metropolitan Sewage District. <https://www.freshcoastguardians.com>

Moreira, T.C.L., Polize, J.L., Brito, M., da Silva Filho, D.F., Chiavegato Filho, A.D.P., Viana, M.C., Andrade, L.H. and Mauad, T. (2021). Assessing the impact of urban environment and green infrastructure on mental health: results from the São Paulo Megacity Mental Health Survey. *Journal of Exposure Science & Environmental Epidemiology*, <https://doi.org/10.1038/s41370-021-00349-x>.

Mygind, L., Kjeldsted, E., Hartmeyer, R.D., Mygind, E., Bolling, M., Bentsen, P., (2019). Immersive nature-experiences as health promotion interventions for healthy, vulnerable, and sick populations? A systematic review and appraisal of controlled studies. *Front Psychol.* 10, 943.

National Biodiversity Report (2015). *Nigeria: Fifth National Biodiversity Report*, Abuja, Nigeria <https://www.cbd.int>

Nigeria. (2015). *Nigeria: 5<sup>th</sup> National Biodiversity Report* <https://www.cbd.int/world>.

Nilon, C.H., Aranso, M.F.J., Cilliers, S.S., Dobbc, C., Frazee, L.J., Goddard, M.A., O'Neil, K.M., Roberts, D., Stander, E.K., Werner, P., Winter, M. and Yocom, K.P. (2017). Planning the future of urban diversity: A global review of city-scale initiatives. *Bioscience*, 67(4), 332-342.

Olayiwola, A.M. and Fakayode, O. (2019). Landscape metrics analysis of land use patterns and changes in suburban local government areas of Ibadan, Nigeria. *South African Journal of Geomatics*, 8(2), 209-220. DOI: <http://dx.doi.org/10.4314/sajg.v8i2.7>.

ONS, (2021). *Tourism and Outdoor Leisure Accounts*. 2021. Office for National Statistics, natural capital, UK.

Pascual-Hortal, L. and Saura, S. (2006). Comparison and development of new graphy-based landscape connectivity indices: Towards the prioritization of habitat patches and corridors for conservation. *Landscape Ecology*, 21, 959–967.

Pauleit, S., Andersson, E., Anton, B., Buijs, A., Haase, D., Hansen, R., Kowarik, I., Stahl Olafsson, A. and Van der Jagt, S. (2019). Urban Green Infrastructure—Connecting People and Nature for Sustainable Cities. *Urban Forest and Urban Greening*, 40, 1–3. [doi:10.1016/j.ufug.2019.04.007](https://doi.org/10.1016/j.ufug.2019.04.007).

Popoola, A.A., Medayese, S.O., Olaniyan, O. M., Onyemenam, P.I. and Adeleye, B.M.



- (2016). Users' perception of urban parks and green networks in Ibadan, *Singapore Journal of Business, Economics and Management Studies*, 51(3401), 1–15.
- Pregitzer, C.C., Hanna, C., Charlop-Powers, S. and Bradford, M.A. (2022). Estimating carbon storage in urban forests of New York City. *Urban Ecosystems*, 25(2), 617–631.
- Puchol-Salort, P., O'Keeffe, J., van Reeuwijk, M. and Mijic, A. (2021). An urban planning sustainability framework: Systems approach to blue green urban design. *Sustainable Cities and Society*, 66: 102677.
- Rafiee, R., Mahiny, A.S. and Khorasani, N. (2009). Assessment of changes in urban green spaces of Mashad city using satellite data. *International Journal of Applied Earth Observation and Geoinformation*, 11: 431–438.
- Raheem, W.M. and Adeboyejo, A.T. (2016). Urban greening and city sustainability in Ibadan metropolis, Nigeria. *Ethiopian Journal of Environment Studies and Management*, 9(3), 287–302.
- Reinwald, F., Weichselbaumer, R., Schindelegger, A. and Damyanovic, D. (2024). From Strategy to Implementation: Mainstreaming Urban Green Infrastructure in Austrias Spatial Planning Instruments for Climate Change Adaptation. *Urban Forestry and Urban Greening*, 94, 128232.
- Richards, D.R., Belcher, R.N., (2020). Global changes in urban vegetation cover. *Remote Sens.* 12 (1), 23. <https://doi.org/10.3390/rs12010023>.
- Riungu, G.K., Peterson, B.A., Beeco, J.A., Brown, G., (2018). Understanding visitors' spatial behavior: a review of spatial applications in parks. *Tour. Geogr.* 20, 833–857.
- Roe, J., Thompson, C., Aspinall, P., Brewer, M., Duff, E., Miller, D, et al. (2013). Green space and stress: evidence from cortisol measures in deprived urban communities. *Int. J. Environ. Res. Public Health*. 10:4086–103.
- Ronan, M., Majid, I., Branko, H., Jahan, K., Wagner, H.H., (2020). Conceptual framework and uncertainty analysis for large-scale, species-agnostic modelling of landscape connectivity across Alberta, Canada. *Sci. Rep. (Nature Publisher Group)* 10 (1). <https://doi.org/10.1038/s41598-020-63545-z>.
- Rudnick, D., Beier, P., Cushman, S., Dieffenbach, F., Epps, C.W., Gerber, L., Hartter, J., Jenness, J., Kintsch, J., Merenlender, A.M., Perkle, R.M., Preziosi, D.V., Ryan, S.J., and Trombulak, S.C. (2012). The Role of Landscape Connectivity in Planning and Implementing Conservation and Restoration Priorities. *Issues in Ecology*. Report No. 16. Ecological Society of America. Washington, DC.
- Russo, A. and Cirella, G. T. (2020). Edible green infrastructure for urban regeneration and food security: Case Studies from the Campania Region. *Agriculture*, 10(8): 358.
- Saeedeh N. and Aysan I. N. (2020). Assessment of urban greenspace fragmentation using landscape metrics (case study: district 2, Tehran city). *Modeling Earth Systems and Environment* <https://doi.org/10.1007/s40808-020-00809-7>.
- Sahraoui, Y., Clauzel, C. and Foltete, J. C. (2021). A metrics-based approach for modelling covariation of visual and ecological landscape qualities. *Ecological Indicators*, 123(9): 107331.
- Scheiber, S. (2022). Re-Designing Urban Open Spaces to Act as Green Infrastructure—The Case of Malta. *Transp. Res. Procedia*, 60, 148–155.
- Science Environment Policy, (2015). *Ecosystem service and the Environment*. Indepth report II produced for the European Commission, DG Environment by the Science Communication unit.

- Severijnen, P. (2018). *Biodiversity by design: Maximizing the biodiversity of Rivierenwulk, Utrecht by landscape architecture design*. M.Sc Thesis submitted to the Wageningen University, Landscape Architecture Group.
- Shackleton, C.M., Blair, A., De Lacy, P., Kaoma, H., Mugwagwa, N., Dalu, M.T., et al. (2018). How important is green infrastructure in small and medium-sized towns? Lessons from South Africa. *Landscape and Urban Planning*, 80: 273–281.
- Siehr, S.A., Sun, M. and Aranda Nucamendi, J.L. (2022). Blue-Green Infrastructure for Climate Resilience and Urban Multifunctionality in Chinese Cities. *WIREs Energy Environ.* 11, e447.
- Silva, C.S., Viegas, I., Panagopoulos, T. and Bell, S. (2018). Environment justice in accessibility to green infrastructure in two European countries. *Land*, 7(134): 1 -23.
- Simon, F. (2015). *Prevalence and Usage of Open Recreational Spaces in Ibadan, Southwest Nigeria*, Department of Architecture, Covenant University, Canaan Land, Ota, Ogun State, Nigeria, Doctoral Thesis.
- Stan, M.I. (2022). Are public administrations the only ones responsible for organizing the administration of green spaces within the localities? An assessment of the perception of the citizens of Constanța municipality in the context of sustainable development. *Technium Social Sciences Journal*, 31: 58-74.
- Sugiyama, T., Leslie, E., Giles-Corti, B. and Owen, N. (2008). Associations of neighbourhood greenness with physical and mental health: do walking, social coherence and local social interaction explain the relationships? *Journal of Epidemiology and Community Health*, 62:e9–e9.
- Sun, X., Tan, X., Chen, K., Song, S., Zhu, X. and Hou, D. (2020). Quantifying landscape-metrics impacts on urban green-spaces and waterbodies cooling effect: The study of Nanjing, China. *Urban for Urban Greening*, 55: 126838.
- UNFAO, (2011). *The state of world's land and water resources for food and agriculture. Rome and earthscan*, London. <https://www.fao.org>.
- United Nations (2018). *Department of Economic and Social Affairs/Population Division World Urbanization Prospects. The 2018 Revision*. United Nations. New York. ESA/P/WP/205.
- United Nations Human Settlements Program, (2012). *State of the world's cities 2012/2013. Prosperity of cities*. United Nations Human Settlements Program, Nairobi, Kenya <https://unhabitat.org>files>.
- Vance, C. and Hedel, R. (2007). The Impact of Urban Form on Automobile Travel: Disentangling Causation from Correlation. *Transportation*, 34, 575–588.
- Vilanova, C., Ferran, J.S. and Concepción, E.D. (2024). Integrating Landscape Ecology in Urban Green Infrastructure Planning: A Multi-Scale Approach for Sustainable Development. *Urban Forestry and Urban Greening*, 94, 128248.
- Vittekk, M., Brink, A., Donnay, F., Simonetti, B. and Deslee, B. (2014). Landcover change monitoring using landsatmss/Tm satellite image data cover, West Africa between 1975 and 1990. *Remote Sensing*, 6, 658 –676. <http://www.mdpi.com/journal/remotesensing>.
- Von Thaden, J., Badillo-Montaño, R., Lira-Noriega, A., García-Ramírez, A., Benítez, G., Equihua, M., Looker, N. and Pérez-Maqueo, O. (2021). Contributions of green spaces and isolated trees to landscape connectivity in an urban landscape. *Urban Forestry and Urban Greening*, 64, 127277. <https://doi.org/10.1016/j.ufug.2021.127277>.

- Wang, X., Blanchet, F. G. and Koper, N. 2014. Measuring habitat fragmentation: An evaluation of landscape patterns metric. *Methods in Ecology and Evolution*, 5(7), 634–646 <https://doi.org/10.1111/2041-210X.12198>
- WBCSD, (2023). *Landscape Connectivity: A call to action*. World Business Council for Sustainable Development, Syngenta, Geneva. [www.wbcsd.org](http://www.wbcsd.org).
- Weisser, W.W., Hensel, M., Barath, S., Culshaw, V., Grobman, Y.J., Hauck, T.E., Jaschinski, J., Ludwig, F., Mimet, A., Perini, K., Enrica, R. Schlofer, M., Schwartz, A., Hensel, D.S. and Volger, V. (2022). Creating ecologically sound buildings by integrating ecology, architecture and computation design. *People and Nature*, 5, 4–20.
- Wickham, J.D., Riitters, K.H., Wade, T.G. and Vogt, P.A. (2010). National Assessment of Green Infrastructure and Change for the Conterminous United States Using Morphological Image Processing. *Landscape and Urban Planning*, 94, 186–195.
- Williams, J.J. and Newbold, T. (2020). Local climatic changes affect biodiversity responses to land use: a review. *Divers. Distrib.* 26 (1), 76–92. <https://doi.org/10.1111/ddi.12999>.
- World Population Review (2024). *Nigeria Population 2024*. <https://worldpopulationreview.com/cities/nigeria>
- Yang, X.J., Liu, Z. (2005). Quantifying landscape pattern and its change in an estuarine watershed using satellite imagery and landscape metrics, *International Journal of Remote Sensing*, 26(23): 5297–5323, doi:10.1080/01431160500219273.
- Zari, M.P. (2018). The importance of urban diversity – An ecosystem services approach. *Biodiversity International Journal*, 2(4), 357–360.
- Zhang, Y. Smith, J.P., Tong, D. and Turner II, B.L. (2022). Optimizing the co-benefits of food desert and urban heat mitigation through community garden planning. *Landscape and Urban Planning*, 226: 104488.
- Zhang, Y., Wang, Y., Ding, N. and Yang, X. (2023). Assessing the Contributions of Urban Green Space Indices and Spatial Structure in Mitigating Urban Thermal Environment. *Remote Sens.*, 15, 2414.
- Zhao, J., Zhao, X., Wu, D., Meili, N. and Fatichi, S. (2023). Satellite-based evidence highlights a considerable increase of urban tree cooling benefits from 2000 to 2015. *Global Change Biology*, 00, 1–13.
- Zheng, Y. and Chen, L. (2024). Modeling the Effect of Green Roofs for Building Energy Savings and Air Pollution Reduction in Shanghai. *Sustainability*, 16, 286.
- Zipperer, W. C., Northrop, R. and Anddrea, M. (2020). *Urban development and environmental degradation*. Oxford Research Encyclopedia of environmental Science <https://doi.org/10.1093/acrefore/9780199389414.013.97>.