

The use of pre-rooted *in-vitro* grapevine shoot segments for elimination of Arabis mosaic virus with chemotherapy

Krisztina Oláh, Mihály Turcsán, Gizella Jahnke, Tamás Deák, Róbert Oláh* and Diána Ágnes Nyitrainé Sárdy

Hungarian University of Agriculture and Life Sciences, Institute for Viticulture and Oenology
HU-1118 Budapest, Villányi Str. 29-43

*Corresponding author: olah.robert@uni-mate.hu

Abstract

Chemotherapy is a widely used technique for virus elimination in crop plants. However, phytotoxicity of the applied antiviral drugs may encumber maintenance of sensible grape plantlets on different sanitation media. We tested the virus eradication efficacy of ribavirin (RBV) on arabis mosaic virus (ArMV) infected *Vitis vinifera* 'Csókaszőlő' *in-vitro* shoot cultures. Due to the phytotoxicity of RBV, pre-rooted 1 cm long shoot tips were placed on 25 mg/l RBV containing medium. The application of pre-rooted shoots instead of unrooted shoot segments promoted plant development and increased the reliability of ArMV removal. An effective method for rooting of 'Csókaszőlő' shoot tips was developed as well.

Keywords: *Vitis vinifera*, ribavirin, virus elimination, antiviral drugs, phytotoxicity

Zusammenfassung

Die Verwendung von vorbereiteten *in-vitro* Traubentriebsegmenten zur Eliminierung des Arabis Mosaic Virus mit Chemotherapie. Chemotherapie ist eine weit verbreitete Technik der Viruseliminierung bei Kulturpflanzen. Aufgrund der Phytotoxizität der eingesetzten antiviralen Chemikalien kann es jedoch schwierig werden, die höchst empfindlichen jungen Pflanzen auf den verschiedenen Sanierungsmedien zu erhalten. In unseren Experimenten haben wir die Wirksamkeit von Ribavirin (RBV) auf mit dem Arabis Mosaikvirus (ArMV) infizierten *Vitis vinifera* 'Csókaszőlő' *in-vitro* Sprosskulturen getestet. Aufgrund der Phytotoxizität von RBV wurden vorbereitete 1 cm lange Triebspitzen auf 25 mg/l RBV-haltiges Medium gesetzt. Die Verwendung von vorbereiteten Triebsegmenten förderte die Pflanzenentwicklung und erhöhte die Zuverlässigkeit der ArMV-Eliminierung. Wir haben auch eine wirksame Methode zur Bewurzelung von 'Csókaszőlő' Triebspitzen entwickelt.

Schlagerwörter: *Vitis vinifera*, Ribavirin, Viruseliminierung, antivirale Chemikalien, Phytotoxizität

Introduction

Arabis mosaic virus (ArMV) is a member of the *Secoviridae* family (genus: *Nepovirus*), and infects a wide range of plant species, for example cherry, grapevine, lily, narcissus, strawberry, and rhubarb (Gao et al., 2016; Komorowska et al., 2021). In the case of grapevine, ArMV is one of the viral agents causing infectious degeneration, and is mainly transmitted by nematodes and grafting (Digiario et al., 2017).

Chemotherapy is a widely used virus elimination technique (Chauhan et al., 2019), and ribavirin (RBV), despite of its phytotoxicity, is one of the most effectively used antiviral drugs in case of plants (Panattoni et al., 2013; Magyar-Tábori et al., 2021). RBV was successfully applied earlier for ArMV eradication on rose (Modarresi et al., 2016), against e.g. grapevine fleck virus (GFKV), grapevine leafroll-associated virus-1 (GLRaV-1), grapevine rupestris stem pitting-associated virus (GRSPaV) on grapevine (Eichmeier et al. 2019, Hu et al., 2018), or apple scar skin viroid in apple (Hu et al., 2022). In addition, its use in combination with thermotherapy has proven to be effective in many cases (Wang et al., 2018).

‘Csókaszőlő’ is a traditional and lately rediscovered Hungarian grapevine cultivar (Galbacs et al., 2009), from which a mother plant was selected and tested by RT-PCR for different viruses; such as ArMV, grapevine fanleaf virus, grapevine leafroll-associated virus-1-3, grapevine rupestris stem pitting-associated virus, and grapevine virus A and B. The selected mother plant proved to be free of most tested viruses (data not shown) except for ArMV. For the sanitation, *in-vitro* shoot cultures were established and maintained on media free of plant growth regulators (PGRs), but the rooting of the excised shoot-segments was very slow and weak on RBV containing medium limiting the shoot development and the effectivity of the sanitation. Therefore, pre-rooted shoot segments were used

on RBV media, and the effectiveness of this method on ArMV eradication was analysed.

Materials and Methods

Production of *in-vitro* shoot cultures and rooting experiments

An ArMV infected *Vitis vinifera* ‘Csókaszőlő’ mother plant was used, in which ArMV was detected with RT-PCR. Ca. 1 cm long shoot tips were prepared from the mother plant and following surface sterilization (70% ethanol 30 s, ca. 0.6% NaOCl 10 min, steril water 3x rinsing), the explants were placed on plant growth regulator (PGR)-free Murashige and Skoog (MS) based medium which contained MS macroelements in half, MS microelements and vitamins in full concentration supplemented with 10 g/l sucrose and 3 g/l gelrite (pH 5.8) ($1/2$ MS). The rooted shoots were tested with RT-PCR and one selected ArMV positive *in-vitro* mother plant was propagated on the same medium. ArMV infection of the micropropagated mother plants used for rooting experiments and RBV treatment was regularly checked by RT-PCR (10 *in-vitro* plants/month). The 1 cm long shoot tips of these *in-vitro* mother plants were either used directly for RBV experiments as a control, or these shoot tips were rooted prior to RBV treatments (Fig. 1). To obtain rooted shootsegments for the RBV treatments we tested the effectivity of indole-3-acetic acid (IAA); indole-3-butyric acid (IBA) and 1-naphthaleneacetic acid (NAA) in different concentrations (Tab. 1). The $1/2$ MS culture medium described above supplemented with 0.5, 0.8 and 1 mg/l IAA, or 0.5 mg/l IBA, or 0.5 mg/l NAA was compared to the same PGR-free medium containing 10 or 20 g/l sucrose. For the evaluation, the number of appearing 1 mm long roots and rooting time (days) were measured. Shoots were growing between 1 and 2 cm on the media with different PGR supplementation, but these data were not evaluated in this experiment. The best PGR combination for rooting was used to produce pre-rooted shoots for RBV treatments.




Fig. 1: Main steps of RBV treatment on 'normal' and pre-rooted 1 cm long shoot tips

Ribavirin treatment of pre-rooted shoots

1 cm long shoot tips with roots were placed on PGR-free solid $\frac{1}{2}$ MS medium containing 25 mg/l RBV (10 g/l sucrose). At 7 different points in time (between 14-68 days after treatment initiation; Tab. 2) 2 mm long shoot tips were excised from the shoots and cultivated on solid $\frac{1}{2}$ MS medium supplemented with 0.02 mg/l BA, 0.01 mg/l NAA and 30 g/l sucrose for a month. After one month micro shoots were placed on PGR-free $\frac{1}{2}$ MS medium (10 g/l sucrose) for two weeks, followed by 0.01 mg/l IAA and 1 mg/l kinetin containing MS medium (10 g/l sucrose) for a month. Developed shoots were rooted in the same way as the shoot segments prior to RBV treatment.

As control, 1 cm long unrooted shoot tips were treated in the same way, but 2 mm long shoot tips were excised only at two points in time: 30 and 60 days after treatment initiation (17 and 33 shoot tips, respectively).

Virus detection using RT-PCR

For isolation of total nucleic acid, a modified CTAB method was used (Xu et al., 2004). Briefly: ca. 50 mg plant material from *in-vitro* leaves was lysed in

1 ml extraction buffer (the washing step of the original protocol was omitted). Chloroform:isoamyl alcohol extraction was applied two times, and 0.8 volumes of isopropanol was used for nucleic acid precipitation (30 min at room temperature). The pellet was finally dissolved in 30 μ l sterile water. After isolation, nucleic acid concentration was measured by NanoPhotometer®N60 (Implen GmbH, Germany). For cDNA synthesis Revert Aid First Strand cDNA kit was used (Thermo Scientific, #K1622) with random hexamer primers, according to the protocol provided by the manufacturer. For reverse transcription 0.2-0.4 μ g template was applied in a final reaction volume of 10 μ l. For PCR reactions 1 μ l 2-fold diluted cDNA was added to 9 μ l reaction mix. Success of DNA synthesis was confirmed with tub-fw2/tub-rev2 primers (Szegedi et al., 2018). ArMV specific PCR amplification applying ArFw and ArRev primers (Gambino and Gribaudo, 2006) and DreamTaq™ (Thermo Scientific, #EP0701) polymerase enzyme was carried out according to the protocol provided by the manufacturer with the following parameters: 94 °C 30 s denaturation, 58 °C 30 s primer annealing, 72 °C 60 s elongation). PCR products were separated in 1,5 (w/v) % agarose gel, and stained with ethidium bromide. The number of samples is shown in Tab. 2.

Statistical analysis

For the evaluation of the obtained data of *in-vitro* shoot cultures we used ART ANOVA (Wobbrock et al., 2011; Elkin et al., 2021). For the estimation of the correlation between the length of ribavirin treatment and the ratio of ArMV-free plants Pearson's product-moment correlation was used in R (R Core Team R; Data Visualization with R and Ggplot2).

Results

Rooted *in-vitro* plants obtained from the shoot tips of the ArMV positive mother plant were tested with RT-PCR using ArMV specific primers (Fig. 2), and PCR amplification using ArMV specific primers was successful in the case of 9 *in-vitro* plants (9/10). A single ArMV positive *in-vitro* mother plant was selected for further micropropagation.

All of the tested PGRs were appropriate for the rooting of the isolated shoot tips based on the number of roots and rooting time (Tab. 1.), however, the application of NAA caused callus formation in addition to rooting, roots induced by IBA were abnormally thick, while in case of IAA new roots showed normal development. Based on the effects on root development, 0.8 mg/l IAA with 10 g/l sucrose was chosen for the propagation of *in-vitro* plantlets. The variance analysis shows a significant difference ($p < 0.001$) for all PGR treatments regarding the number of rooting days except between NAA and IBA (Tab. 1). The effect of applied PGRs on root number per shoots was significant ($p < 0.001$) for all PGR relative to the null control, and the difference between IAA and IBA was significant as well (Fig. 3). Our results match the rooting results of Shatnawi and coworkers (Shatnawi et al., 2011) with the same PGRs on 'Salty Kodary' cultivar.




Fig. 2: Amplification of the 402 bp long fragments of the ArMV genome with PCR test from the first obtained *in-vitro* plants. One *in-vitro* plant (line 8) was selected for further experiments. M: H3 RTU DNA ladder (NIPPON Genetics Europe GmbH, Cat. no.: MWD 100). 0: Nucleic acid-free control. +: Positive control. Lanes 1 to 10 are PCR products from cDNAs of *Vitis vinifera* 'Csókaszó'




Fig. 3: The effect of PGRs on root number per shoots. Letters (a-c) show the groups of the post-hoc test

Tab. 1: The effect of the applied PGRs and sucrose concentration on rooting of the 1 cm long ‘Csókaszóľő’ shoot segments. Number of samples: IAA (0.5 mg/l)=28, IAA (0.8 mg/l)=62, IAA (1 mg/l)=15, IBA (0.5 mg/l)=30, IBA (0.8 mg/l)=33, NAA (0.5 mg/l)=5, NULL=30

PGR	PGR concentration mg/l	Sucrose g/l	Number of shoots on the rooting medium	Average number of days until the appearance of 1 mm long roots	Ratio of rooted shoots % (rooted shoots/all shoots)	Average number of roots/shoot
-	-	20	15	34.66	78.9 (19/15)	1.7
-	-	10	15	33.73	83,3 (15/18)	1.9
IAA	0,5	20	4	20	100 (4/4)	2.5
IAA	0,5	10	24	18	100 (24/24)	2.8
IAA	0,8	10	62	14.63	100 (62/62)	2.98
IAA	1	20	6	21.5	100 (6/6)	2
IAA	1	10	9	13.66	100 (9/9)	3.44
IBA	0,5	20	7	26.57	100 (7/7)	7.1
IBA	0,5	10	23	25	95,8 (24/23)	5.1
IBA	0,8	10	33	22.8	91,6 (36/33)	6.6
NAA	0,5	20	5	25	100 (5/5)	3.6

2 mm long shoot tips were isolated at different time points from the pre-rooted and RBV-treated shoots. After regeneration, the presence of ArMV content in the *in-vitro* plants was tested by RT-PCR (Fig. 4, Tab. 2). After 30 days of treatment over 50% of the plants were free from ArMV. The ratio of ArMV-free plants and the length of RBV treatment shows strong linear correlation based on Pearson’s product-moment correlation method (95 percent confidence interval, cor = 0.8261759).

If 1 cm long non-rooted shoot tips were treated for 30 or 60 days (Fig. 1) on the same RBV containing medium, then 50% or 69% of the regenerated plants were ArMV-free respectively. Our results show that RBV treatment of pre-

rooted shoots proved to be successful in ArMV elimination, and pre-rooting of the shoots did not change substantially the effectiveness of sanitation in case of ‘Csókaszőlő’ cultivar.




Fig. 4: Amplification of the 402 bp long fragments of ArMV with RT-PCR test from *in-vitro* plants regenerated from the 2 mm long shoot tips of RBV-treated plants. M: H3 RTU DNA ladder (NIPPON Genetics Europe GmbH, Cat. no.: MWD 100). 0: Nucleic acid-free control. +: Positive control. Lanes 1 to 9 are PCR products from cDNAs of *Vitis vinifera* ‘Csókaszőlő’.

Tab. 2: RT-PCR test results of regenerated plants from 2 mm long shoot tips of pre-rooted and RBV-treated shoots.

Length of RBV treatment (day)	Number of isolated 2 mm long shoot tips from the long shoot tips from the RBV-treated shoots	Number of PCR-tested plants	Ratio of ArMV negative plants (%)
14	14	8	12.5
21	11	5	40
26	25	10	30
30	32	13	61.5
42	24	18	55
60	35	26	54
68	31	22	82

Conclusions

Pre-rooting of *in-vitro* shoot segments for RBV treatment is a possible solution to limit the adverse effects of chemotherapy. Using of 0.8 mg/l IAA reduced the rooting time of the shoot segments and increased the average number of roots. At the same time the application of pre-rooted shoot segments did not reduce the efficiency of RBV treatment, but this method

decreased the ratio of dead or non-developing shoots from 73% to 14% (after 30 days of RBV treatment). This method could help in the development of the different chemotherapy treatments by increasing chance of survival of treated shoot segments. ArMV elimination effectiveness with pre-rooted shoot segments in our experiments is comparable to the results obtained on rose *in-vitro* shoot segments (Modarresi et al. 2016). In the case of grapevine, somatic embryogenesis proved to be 100% successful for ArMV removal (Olah et al., 2022), but to the best of our knowledge this is the first report about ArMV elimination with chemotherapy in grapevine. On the other hand, RBV was effective against some grapevine viruses, such as GFKV, grapevine fanleaf virus (GFLV), and grapevine Pinot gris virus (GPGV) (Komínek et al., 2016; Weiland et al., 2004), but lower efficiency was experienced in the case of GLRaV-1, GRSPaV, and grapevine virus A (GVA) (Gutá and Buciumeanu, 2011; Hu et al., 2018; Komínek et al., 2016; Skiada et al., 2013). The efficiency of RBV treatment depends on the virus, on the cultivar and on experimental setting (Komínek et al., 2016), and our method (pre-rooting, 25 mg/l RBV, min. 30 day) proved to be reliable and effective for ArMV elimination in ‘Csókaszőlő’.

Acknowledgments

M.T. is PhD student at Hungarian University of Agriculture and Life Sciences at the Doctoral

School of Horticultural Sciences. This research was funded by National Research, Development and Innovation Office (NKFIH) grant number: K131679.

References

- Chauhan, P., Singla, K., Rajbhar, M., Singh, A., Das, N., Kumar, K.** 2019: A systematic review of conventional and advanced approaches for the control of plant viruses. *Journal of Applied Biology and Biotechnology* 7(4): 89-98.
- Data Visualization with R and Ggplot2 | the R Graph Gallery** Available online: <https://r-graph-gallery.com/ggplot2-package.html> (28.02.2023)
- Digiario, M., Elbeaino, T., Martelli, G.P.** 2017: Grapevine fanleaf virus and other old world nepoviruses. In: Meng, B., Martelli, G., Golino, D., Fuchs, M. (eds) *Grapevine Viruses: Molecular Biology, Diagnostics and Management*. - Springer, Cham, 2017, S. 47-82.
- Eichmeier, A., Kominkova, M., Pecenka, J., Kominek, P.** 2019: High-throughput small RNA sequencing for evaluation of grapevine sanitation efficacy. *Journal of Virological Methods* 267: 66-70.
- Elkin, L.A., Kay, M., Higgins, J.J., Wobbrock, J.O.** 2021: An aligned rank transform procedure for multifactor contrast tests. *UIST 2021 - Proceedings of the 34th Annual ACM Symposium on User Interface Software and Technology*, S. 754–768. doi:10.1145/3472749.3474784.
- Galbacs, Z.S., Molnar, S., Halasz, G., Kozma, P., Hoffmann, S., Kovács, L., Veres, A., Galli, Z.S., Szőke, A., Heszky, L., Kiss, E.** 2009: Identification of grapevine cultivars using microsatellite-based DNA barcodes. *VITIS-Journal of Grapevine Research* 48(1): 17-24.
- Gambino, G., Gribaudo, I.** 2006: Simultaneous detection of nine grapevine viruses by multiplex reverse transcription-polymerase chain reaction with coamplification of a plant RNA as internal control. *Phytopathology* 96(11): 1223-1229.
- Gao, F., Lin, W., Shen, J., Liao, F.** 2016: Genetic diversity and molecular evolution of arabis mosaic virus based on the CP gene sequence. *Archives of Virology* 161(4): 1047-1051.
- Guță, I.C., Buciumeanu, E.C.** 2011: Grapevine chemotherapy for elimination of multiple virus infection. *Romanian Biotechnological Letters* 16(5): 6535-6539.
- Hu, G., Dong, Y., Zhang, Z., Fan, X., Ren, F., Li, Z., Zhang, S.** 2018: Elimination of Grapevine rupestris stem pitting-associated virus from *Vitis vinifera* 'Kyoho' by an antiviral agent combined with shoot tip culture. *Scientia Horticulturae* 229: 99-106.
- Hu, G., Dong, Y., Zhang, Z., Fan, X., Ren, F.** 2022: Inefficiency of ribavirin to eliminate apple scar skin viroid from apple plants. *Plant Cell, Tissue and Organ Culture (PCTOC)* 151(1): 189-197.
- Komínek, P., Komínková, M., Jandová, B.** 2016: Effect of repeated Ribavirin treatment on grapevine viruses. *Acta Virologica* 60(4): 400-403.
- Komorowska, B., Hasiów-Jaroszewska, B., Budzyńska, D.** 2021: Genetic variability and molecular evolution of arabis mosaic virus based on the coat protein gene sequence. *Plant Pathology* 70(9): 2197-2206.
- Magyar-Tábori, K., Mandler-Drienyovszki, N., Hanász, A., Zsombik, L., Dobránszki, J.** 2021: phytotoxicity and other adverse effects on the *in-vitro* shoot cultures caused by virus elimination treatments: Reasons and Solutions. *Plants* 10(4): 670.
- Modarresi Chahardehi, A., Rakhshandehroo, F., Mozafari, J., Mousavi, L.** 2016: Efficiency of a chemo-thermotherapy technique for eliminating

Arabis mosaic virus (ArMV) and Prunus necrotic ringspot virus (PNRSV) from *in vitro* rose plantlets. Journal of Crop Protection 5(4): 497-506.

Olah, R., Turcsan, M., Olah, K., Farkas, E., Deak, T., Jahnke, G., Sardy, D.A.N. 2022: Somatic Embryogenesis: A Tool for Fast and Reliable Virus and Viroid Elimination for Grapevine and other Plant Species. Horticulturae 8(6): 508.

Panattoni, A., Luvisi, A., Triolo, E. 2013: Review. Elimination of Viruses in Plants: Twenty Years of Progress. Span. J. Agric. Res. 11: 173–188. <https://doi.org/10.5424/sjar/2013111-3201>.

R Core Team R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. Available online: <http://www.r-project.org/>.

Shatnawi, M., Anfoka, G., Shibli, R., Al-Mazra'awi, M., Shahrour, W., Arebiat, A. 2011: Clonal propagation and cryogenic storage of virus-free grapevine (*Vitis vinifera* L.) via meristem culture. Turkish Journal of Agriculture and Forestry 35(2): 173-184.

Skiada, F.G., Maliogka, V.I., Katis, N.I., Eleftheriou, E.P. 2013: Elimination of Grapevine rupestris stem pitting-associated virus (GRSPaV) from two *Vitis vinifera* cultivars by *in vitro* chemotherapy. European journal of plant pathology 135: 407-414.

Szegedi, E., Deák, T., Turcsán, M., Szénási, M., Bordé, Á., Oláh, R. 2018: Evaluation of intron containing potential reference gene-specific primers to validate grapevine nucleic acid samples prepared for conventional PCR and RT-PCR. Vitis 57(2): 69-73.

Xu, Q., Wen, X., Deng, X. 2004: A simple protocol for isolating genomic DNA from chestnut rose (*Rosa roxburghii* Tratt) for RFLP and PCR analyses. Plant Molecular Biology Reporter 22: 301-302.

Wang, M.R., Cui, Z.H., Li, J.W., Hao, X.Y., Zhao, L., Wang, Q.C. 2018: *In vitro* thermotherapy-based methods for plant virus eradication. Plant methods 14(1): 1-18.

Weiland, C.M., Cantos, M., Troncoso, A., Perez-Camacho, F. 2004: Regeneration of virus-free plants by *in vitro* chemotherapy of GFLV (Grapevine fanleaf virus) infected explants of *Vitis vinifera* L. Cv'Zalema'. Acta Hort. 652: 463-466.

Wobbrock, J.O., Findlater, L., Gergle, D., Higgins, J.J. 2011: The Aligned Rank Transform for Nonparametric Factorial Analyses Using Only ANOVA Procedures. Conference on Human Factors in Computing Systems – Proceedings. S. 143–146. doi:10.1145/1978942.1978963.

Received April, 26th, 2023