DOI: 10.2478/agr-2024-0010

THE ANNALS OF "VALAHIA" UNIVERSITY OF TARGOVISTE

OPTIMIZING FEED FORMULATION STRATEGIES FOR ATTAINING OPTIMAL NUTRITIONAL BALANCE IN HIGH-PERFORMING DAIRY GOATS IN INTENSIVE FARMING PRODUCTION SYSTEMS

Cosmin Șonea[1], Raluca Aniela Gheorghe-Irimia[1*], Makki Khalaf Hussein Al Dulaimi[2], Lavinia Udrea[3], Dana Tăpăloagă[1], Paul-Rodian Tăpăloagă[1]

[1]University of Agronomic Sciences and Veterinary Medicine of Bucharest, 59 Marasti Blvd, District 1, Bucharest, Romania

[2]Al-Furat Al-Awsat Technical University, Baghdad, Iraq

[3]Valahia University of Targoviste, 13 Sinaia Alley, Targoviste, Romania

[*]Corresponding author email: raluca.irimia@fmv.usamv.ro

Abstract

Genetic factors play a pivotal role in shaping feed formulation strategies for dairy goats, influencing milk production, composition, and overall productivity through the modulation of genetic parameters and polymorphisms associated with milk protein fractions and yield. Moreover, environmental challenges, including heat stress and feed availability fluctuations, pose significant hurdles in achieving optimal nutrition and performance, necessitating precision livestock farming and the utilization of local by-products to mitigate their impact on herd health and productivity. In addition, nutritional interventions, such as supplement utilization and alternative feed incorporation, offer promising avenues for addressing specific deficiencies and enhancing overall performance, from fatty acid supplementation to herbal feed additives, while balancing rations to meet evolving nutritional needs and ensuring long-term sustainability remains paramount for optimizing herd nutrition, health, and performance in intensive dairy goat production systems. The aim of this study was to explore the multifaceted influences of genetic factors, environmental challenges, and nutritional interventions on feed formulation strategies and overall herd nutrition, health, and performance in intensive dairy goat production systems.

Keywords: dairy goats, feed formulation, genetic factors, environmental challenges, nutritional interventions

1. INTRODUCTION

Maximizing feed efficiency in dairy goat husbandry plays a pivotal role in augmenting milk productivity and safeguarding the overall well-being of the livestock. The economic significance of dairy goat farming resonates across diverse locales, including the Mediterranean basin, China, and sub-Saharan Africa, where a burgeoning appetite for dairy goat commodities persists [1,2,3]. Pertinent challenges, such as operational efficacy, waste management intricacies, and animal welfare prerogatives, are discernible within the dairy goat sector, notably prevalent in the Americas [3]. Additionally, the burgeoning demand for goat-derived dairy products catalyzes the proliferation of commercial dairy goat enterprises in the United States [3].

The availability and caliber of feed reservoirs exert profound ramifications on dairy goat agribusiness. Within regions constrained by land scarcity, the judicious exploitation of diversified feedstock, encompassing crop residues, agro-industrial byproducts, and alimentary discards, becomes indispensable in surmounting productivity thresholds [4]. Moreover, the inadequacy of grazing pastures, particularly during arid spells, prompts the incorporation of imported conventional feed constituents into dairy goat rations [5]. The dietary regimen for dairy ovine and caprines in Greece, for instance, predicates on a synergistic amalgam of seasonal herbage foraging and supplementary nourishment, accentuating the exigency of an equilibrated and sustainable alimentation stratagem [6].

Beyond dietary composition, the deployment of feed additives and supplements constitutes a focal point of scholarly inquiry. Investigative endeavors have scrutinized the efficacy of feed supplements vis-à-vis dairy goat cohorts, with a view to ameliorating feed protein utilization efficiency and overarching lactogenic proficiency [7,8]. Furthermore, the ramifications of chronic thermal stress on lactational aptitude in dairy goats have undergone scientific scrutiny, underscoring the pivotal role of environmental variables in feed optimization and lacteal output [9].

The optimization of feed dynamics in dairy goat production transcends mere yield augmentation, encapsulating a broader paradigm of animal welfare enhancement. Pioneering feed management methodologies, such as the assimilation of feed additives, hold promise in fortifying environmental integrity and elevating the nutritional quotient of dairy goat derivates [10]. Moreover, delving into the nexus between feed regimes and the transcriptomic milieu of lactating dairy goat leukocytes furnishes invaluable insights into the physiological responses of livestock to divergent nutritional stimuli [11].

The aim of this review paper is to comprehensively assess the current state of knowledge regarding feed formulation strategies in dairy goat production systems, with a focus

on optimizing nutritional balance to enhance milk production and ensure the overall health and welfare of the animals. Additionally, the paper aims to explore the economic, environmental, and physiological factors influencing feed optimization practices, while also examining emerging trends and innovative approaches in the field.

2. NUTRITIONAL REQUIREMENTS OF HIGH-PERFORMING DAIRY GOATS

The energy requisites of high-performing dairy goats are subject to multifaceted influences, encompassing factors such as heat stress, dietary supplementation modalities, and metabolic intricacies. Scientific investigations have delineated how heat stress perturbations can engender alterations in the thermophysiological parameters and lactational performances of dairy goats, thereby exerting discernible ramifications on their energy equilibrium and overarching metabolic milieu [11,12]. Additionally, inquiries into the effects of propylene glycol supplementation amidst heat-stressed dairy goats have elucidated its impact on milk productivity and energetic metabolism across divergent environmental milieus, underscoring its potential significance [13]. Furthermore, the evaluation of multinutrient blocks, integrating constituents like avocado pulp and peels, has shed light on alternative nutritional paradigms aimed at bolstering energy requisites, thereby enriching our understanding of dietary strategies in dairy goat management [14].

Moreover, scientific endeavors have scrutinized the efficacy of dietary supplements, such as zinc-methionine, in ameliorating milk production outcomes and udder health in dairy goats, accentuating the pivotal role of specific nutrients in fulfilling the energy demands of high-performing dairy goats [15]. Additionally, investigations into the ameliorative effects of berberine supplementation on energy balance, glucose homeostasis, and hepatic functionality in transition dairy goats have underscored the potential for targeted nutritional interventions in fortifying energy metabolism dynamics within dairy goat cohorts [16].

The protein exigencies of high-performing dairy goats are similarly subject to diverse influences, including heat stress dynamics, dietary supplementation paradigms, and metabolic responses. Scientific inquiry has elucidated how heat stress perturbations can instigate modifications in the thermophysiological parameters and lactational performances of dairy goats, thereby impinging upon their energy equilibrium and overarching metabolic milieu [12]. Furthermore, investigations into the efficacy of propylene glycol supplementation amidst heat-stressed dairy goats have shed light on its consequential impact on milk productivity and energetic metabolism across varying environmental contexts, thus delineating its potential relevance [13].

Moreover, the exploration of dietary interventions, such as multinutrient blocks, has yielded insights into alternative nutritional strategies aimed at supporting protein requirements, thereby enriching our understanding of dietary modulation in dairy goat husbandry.

Furthermore, scientific inquiries have delved into the interplay between water requisites and various factors such as heat stress dynamics, dietary constitution, and lactational performance among high-performing dairy goats. Investigations have evidenced that heat stress exerts a pronounced influence on water intake patterns among dairy goats, with lactating does experiencing a notable surge in water consumption under such thermal exigencies [12]. Additionally, inquiries into dietary cation and anion differences (DCAD) have elucidated their impact on water balance dynamics, wherein goats from the DCAD group exhibited augmented apparent water balance over a 24-hour period [17]. Moreover, studies on dietary roughage composition have revealed discernible effects on water intake behaviors, with the inclusion of spineless cactus notably diminishing water consumption via traditional drinkers [18]. Additionally, investigations into water preference modalities, such as water nipples versus water bowls, have yielded insights into the nuanced drinking patterns and water intake propensities among dairy goat populations [19]. Furthermore, scientific inquiries into the physiological responses and lactational performances of late-lactation dairy goats under heat stress conditions have unveiled a substantial augmentation in total water evaporation rates among heat-stressed goats vis-à-vis those reared under thermoneutral conditions [20].

3. FACTORS INFLUENCING FEED FORMULATION IN INTENSIVE DAIRY GOAT PRODUCTION

Genetic determinants wield substantial influence over feed formulation and management strategies within intensive dairy goat production systems. The genomic architecture inherent to dairy goat breeds intricately influences myriad aspects of feed formulation, encompassing milk production, compositional attributes, and overall productivity metrics. Empirical studies have delineated the genetic underpinnings of total lactation yields encompassing milk, fat, protein, and somatic cell score in dairy goats, thereby elucidating the genetic imprints upon both qualitative and quantitative facets of milk production [21].Moreover, investigations into the genetic variability within dairy goat cohorts have underscored its consequential impact on a spectrum of milk production, reproductive, and immunity traits, thereby spotlighting the genetic bedrock underpinning productivity and health parameters in dairy goat populations [22,23].

Furthermore, the genetic architecture delineated by polymorphic variations in key genes, such as CSN1S1 and CSN2, has been correlated with discernible variances in

milk protein fractions and milk yield, thus furnishing insights into the intricate genetic regulation governing milk composition and output dynamics within dairy goat cohorts [24,25]. Delving deeper into the genetic milieu, explorations into the genetic diversity and ancestral origins of dairy goat breeds have provided invaluable insights into the foundational genetic drivers shaping productivity and adaptability across diverse demographic clusters [26].

Beyond milk production metrics, genetic determinants extend their purview to encompass conformational traits, the investigation of which has yielded illuminating insights into their interplay with milk yield trajectories across lactation cycles, thereby elucidating the genetic modulation of structural attributes and milk productivity in dairy goat populations [27]. Moreover, the recognition of genetic enhancement avenues has emerged as a pivotal imperative in augmenting productivity and performance metrics, thereby underscoring the indispensable role of genetic selection and breeding paradigms within intensive dairy goat production ecosystems [28, 4].

Concurrently, environmental dynamics emerge as pivotal determinants in shaping feed formulation and management paradigms within intensive dairy goat production systems. The valorization of agricultural by-products, such as residual fruits from tomato and cucumber processing, has been scrutinized for its efficacy in augmenting nutrient utilization, ruminal fermentation dynamics, microbial abundances, and milk yield and composition profiles in dairy goat herds [29]. This highlights the potential of leveraging by-products to curtail feeding costs and ameliorate environmental repercussions associated with waste accumulation.

Moreover, the intensification trajectory of dairy goat farming has garnered attention due to its discernible impact on the environmental footprint of milk production. Comparative evaluations across farm clusters, characterized by varying intensification levels, have unveiled akin environmental performances on a product-centric basis, thereby illuminating the intricate interplay between intensification paradigms, management modalities, and environmental ramifications within dairy goat production domains. Furthermore, the advent of precision livestock farming holds promise in augmenting environmental sustainability benchmarks and curbing greenhouse gas emissions within intensive dairy goat production landscapes [30].

Additionally, the adoption of efficient reproductive and herd health management practices has been discerned as pivotal in sculpting milk yield trajectories, reproductive efficiency indices, and overall productivity metrics within dairy goat enterprises. Insights gleaned from structural equation modeling have further elucidated the intricate interplay between education, training interventions, and management frameworks, ultimately reverberating upon the production performance dynamics of dairy goat enterprises.

Furthermore, the judicious utilization of agricultural by-products and the integration of pasture management methodologies have emerged as efficacious management modalities shaping feed formulation dynamics and nutrient utilization metrics within dairy goat production landscapes. The substitution of conventional ingredients with by-products has been underscored as a means to slash feeding costs and bolster profit margins, thereby accentuating the transformative potential of sustainable feed resources within dairy goat husbandry.

Moreover, the strategic deployment of feeding and nutrition management strategies assumes paramount significance in augmenting growth performance and productivity benchmarks within dairy goat cohorts. Investigations into the supplementation regimens incorporating mineral blocks and a cocktail of fatty acids, organic acids, and minerals have shed light on their consequential impact on milk yield trajectories and hematological profiles among dairy goat populations, thus underscoring the pivotal role of efficacious feeding management strategies in dairy goat farming.

4. FEED INGREDIENTS AND THEIR ROLE IN DAIRY GOAT NUTRITION

Forages serve as linchpins in the nutritional paradigm and overall welfare of dairy goats. The integration of forages into dairy goat diets has undergone extensive scrutiny due to its profound implications on milk production, nutrient assimilation, and animal well-being. These forages impart indispensable nutrients, such as fiber, protein, and minerals, pivotal for sustaining the health and productivity of dairy goats [31]. Their inclusion in dietary regimens exerts discernible influences on milk production, composition, and sensory attributes, underscoring their pivotal role in optimizing both the quality and quantity of milk output [32].

Moreover, the nutritional constitution inherent to forages significantly shapes the quality and constituents of goat milk. Investigations have delved into the fatty acid profiles, protein contents, and mineral compositions of goat milk vis-à-vis distinct forage species and dietary regimes, illuminating the profound repercussions of forages on the nutritional milieu of milk [33,34]. Furthermore, the incorporation of forages has been linked to nuanced shifts in ruminal microbiota, growth kinetics, carcass attributes, and meat quality among fattening goats, underscoring the manifold impacts of forages on animal vitality and productivity [35].

Additionally, the accessibility and caliber of forages are pivotal considerations in feed formulation for dairy goat husbandry. Given the substantial heterogeneity in the nutritional profiles of forages, their nutritive potency can markedly sway the overall health and performance metrics

of dairy goats [31]. Adroit forage management practices encompassing selection, harvest, and storage regimes are indispensable for securing a reservoir of premium-grade forages conducive to dairy goat nutrition.

Supplements occupy a pivotal niche in the dietary mosaic of dairy goats, concomitantly bolstering milk quality, animal wellness, and overall productivity. The incorporation of supplements into dairy goat diets has undergone exhaustive scrutiny for its ramifications on milk production dynamics, nutrient utilization efficiencies, and animal welfare indices. An array of supplements ranging from fatty acids and microalgae to vitamins has been interrogated for their potential in enhancing the nutritional profile of milk, amplifying milk yields, and fortifying animal health.

Scholarly inquiry has delineated that the integration of fatty acid supplements into dairy goat diets orchestrates discernible modulations in milk lipid biosynthesis and lipolysis pathways, thereby effectuating shifts in milk fat contents and fatty acid spectra. Furthermore, the supplementation of microalgae in caprine diets has evinced augmentations in the concentrations of docosahexaenoic acid and total n-3 fatty acids in milk and cheese matrices, thereby accentuating the nutritional exquisiteness of milk.

Moreover, the supplementation regimen incorporating dietary L-carnitine has been associated with elevated blood carnitine levels in dairy goats, attesting to the potential of nutritional adjuncts in influencing metabolic dynamics and animal vitality. Concurrently, the incorporation of high-palmitic acid fat supplements has garnered attention for its purported efficacy in optimizing lactational performances among high-producing dairy goats, thereby underscoring the pivotal role of dietary fats in optimizing milk output.

Furthermore, the judicious utilization of supplements in dairy goat nutrition has been correlated with ameliorative shifts in milk yield, milk quality, and serum biochemical profiles. Scientific research has evidenced that the supplementation of ecotrofin™ engenders favorable outcomes on milk yield, milk quality, and serum biochemistry in lactating goats, thereby contributory to the amelioration of animal welfare and productivity.

By-products serve as formidable constituents in dairy goat nutrition, proffering a cornucopia of nutritional dividends and espousing tenets of sustainable agricultural practices. By-products stemming from sundry agricultural and industrial processes have been harnessed in dairy goat diets, furnishing an alternate reservoir of nutrients and redounding to the mitigation of waste burdens. The assimilation of by-products into dairy goat nutrition has been scrutinized for its ramifications on milk production dynamics, nutrient assimilation efficiencies, ruminal fermentation kinetics, and overarching animal health.

By-products such as pistachio remnants, dried distillers grains with solubles (DDGS), and dried citrus pulp (DCP) can be efficaciously assimilated into dairy goat diets, eliciting nuanced shifts in ruminal fermentation modalities, blood metabolic profiles, and milk fatty acid compositions [36]. Moreover, the harnessing of agricultural by-products has been probed for its influence on milk attributes, nutrient assimilation kinetics, ruminal fermentation cascades, and methane abatement in dairy goat cohorts, thus accentuating the potential of by-products in enhancing milk nutritional profiles and curtailing feeding overheads.

Furthermore, the incorporation of by-products into dairy goat nutrition has been conjoined with ameliorative transformations in milk technological indices, sensory attributes, and health dividends. Other research suggests that the inclusion of artichoke by-products in dairy goat rations precipitates salutary impacts on milk technological and sensory dimensions, fostering a reduction in feeding expenditures without compromising milk quality [37]. Additionally, the exploration of fermented goat dairy products has unearthed their putative health benefits, spanning anti-inflammatory, cardiovascular, and metabolic arenas, thereby catalyzing the emergence of novel dairy formulations enriched with enhanced nutritional and functional attributes [38].

Concentrates assume an indispensable mantle in undergirding the nutritional edifice and overall performance matrix of dairy goats. The integration of concentrates into dairy goat diets has been the subject of intensive inquiry, courtesy of its far-reaching repercussions on milk production, nutrient utilization kinetics, and animal health indices [39]. High-yielding dairy goats necessitate diets replete with copious starchy concentrates to underpin their energy requisites, particularly in attaining apogean lactational benchmarks [39].

Their inclusion in dairy goat diets engenders palpable effects on milk output, compositional nuances, and gustatory attributes, thus underscoring the pivotal role of concentrates in optimizing milk quality and yield [40]. The nutritional constitution intrinsic to concentrates markedly shapes the quality and constituents of goat milk. Research elucidation has probed into the fatty acid profiles, protein titres, and mineral compositions of goat milk in response to disparate concentrate formulations and dietary configurations, thereby accentuating the imprint of concentrates on milk nutritional profiles [41,42].

Moreover, the incorporation of concentrates has been correlated with discernible perturbations in ruminal microbiota, growth kinetics, carcass attributes, and meat quality among fattening goats, thus spotlighting the multifaceted impacts of concentrates on animal vigor and productivity [43]. Furthermore, the accessibility and caliber of concentrates assume paramount importance in feed formulation for dairy goat enterprises.

Given the marked heterogeneity in the nutritional spectra of concentrates, their nutritive valor can substantially sway the overall health and performance metrics of dairy goats [31]. Hence, judicious concentrate management encompassing selection, formulation, and feeding modalities emerges as quintessential for furnishing a reservoir of premium-grade concentrates conducive to dairy goat nutrition.

5. STRATEGIES FOR OPTIMIZING FEED FORMULATION

The strategic balancing of rations to meet the exacting nutritional requisites of dairy goats stands as a linchpin for optimizing milk production, animal vitality, and overall productivity. Achieving equilibrium in ration formulation mandates meticulous scrutiny of the nutrient profiles inherent to feed constituents, ensuring precise alignment with the dietary imperatives of dairy goats. This meticulous process constitutes a sine qua non for attaining peak milk yields, superior milk quality, and enhanced animal welfare.

Research underscores that the formulation of total mixed rations (TMR) necessitates judicious calibration to furnish a harmonious amalgam of metabolizable protein, minerals, and vitamins, thereby regulating energy intake and curtailing the total dry matter ingested by dairy goats [44]. Moreover, the meticulous balancing of rations sourced from a diverse array of locally available feeds reverberates in discernible alterations in greenhouse gas emissions stemming from dairy goat operations, thus accentuating the environmental ramifications of ration formulation [45].

Further research underscores the pivotal role of ration balancing in satisfying the protein and energy exigencies of lactating goats, alongside modulating nutritional metabolism during critical transitional phases [46,47]. The strategic harmonization of rations is inextricably linked to improved shifts in milk production, milk quality, and animal welfare, thereby underpinning augmented performance and productivity among dairy goats [47,48].

Additionally, the deployment of precision diet formulation and the nuanced balancing of rations to optimize essential amino acid ratios hold promise for bolstering net incomes and curbing environmental footprints in dairy farming milieus [49]. Moreover, the meticulous balancing of rations is allied with enhancements in nitrogen efficiency, nutrient uptake, and farm profitability within dairy goat production ecosystems [50].

In a complementary vein, the formulation of feeds tailored to augment digestibility and assimilation in dairy goats constitutes a cornerstone for optimizing milk production, nutrient uptake, and overall animal health. The discerning curation and balancing of feed constituents are pivotal for meeting the exacting nutritional demands of dairy goats and fostering efficient feed utilization.

Assorted investigations have probed into the impacts of diverse feed components and supplements on nutrient digestibility, rumen fermentation, and milk production in dairy goat populations.

For instance, the integration of mulberry leaf biomass into dietary regimens has evinced pronounced enhancements in fiber digestion and utilization, thereby eliciting superior lactational outcomes in both goats and dairy cattle [51]. Moreover, supplementation with fibrolytic enzymes has demonstrated efficacy in augmenting feed digestion and milk production in dairy cows [52]. Additionally, the inclusion of non-traditional feed ingredients such as tomato and cucumber waste fruits has evinced potential in efficaciously supplanting cereal-based concentrates in ruminant diets, thereby underscoring the efficiency of feed utilization [53].

Furthermore, the partial substitution of concentrate with feed blocks has been associated with ameliorations in nutrient utilization, microbial nitrogen flux, and milk yield and composition in goats [54]. Likewise, reductions in dietary crude protein levels have been shown to enhance nutrient digestion, serum metabolites, and nitrogen utilization in growing goat kids [55]. Notably, the compaction of complete feed pellets has been found to augment acceptability, digestibility, and nutrient utilization, thereby eliciting favorable shifts in animal performance [56].

Moreover, supplementation with glycerin has evinced efficacy in enhancing feed digestibility and bolstering microbial protein production in the rumen of cattle [57]. The incorporation of non-traditional feed resources such as Azolla pinnata and mango seeds has been found to heighten nutrient digestibility, milk yield, and feed conversion efficiency in dairy goats [58,59]. Additionally, the treatment of feed resources employing biochemical agents such as effective microbes and urea molasses has been demonstrated to enhance feed digestibility and nutritive value [60].

Addressing specific nutritional lacunae in dairy goats assumes paramount significance in safeguarding their health and optimizing productivity. Myriad investigations have delved into the repercussions of specific nutritional deficits and the potential ameliorative interventions to redress them. Notably, iron deficiency aenemia (IDA) emerges as a pervasive nutritional shortfall globally, with research delineating the potential of fermented goat milk consumption to ameliorate brain molecular functions during IDA convalescence, thereby proffering salutary benefits in addressing this deficiency [61]. Furthermore, calcium-enriched goat's milk has been documented to expedite iron status recovery more efficaciously than calcium-enriched cow's milk in rats afflicted with nutritional ferropenic aenemia, thus spotlighting the potency of specific dietary interventions in rectifying iron deficits [62].

Moreover, selenium deficiency looms as a recurrent concern in dairy goat husbandry, with supplementation utilizing barium selenate emerging as a recommended prophylactic measure for forestalling clinical mastitis and ameliorating milk quality in selenium-depleted locales [63]. Copper deficiency constitutes another salient nutritional apprehension, with studies documenting clinical manifestations of deficiency in both dairy goats and offspring, thereby underscoring the imperative of redressing specific mineral deficits in dairy goat nutrition [64].

Recently, gamut of alternative feed sources and supplements to enrich the nutritional caliber of dairy goat diets and bolster milk production were researched. For instance, the adjunct of vegetable oils to dairy goat diets has been demonstrated to elevate the lipid profile of milk and its derivatives, furnishing a viable alternative to supplemental feeding during the dry period [65]. Additionally, date fruit waste has been probed as a prospective feed alternative for lactating dairy goats, thus showcasing its viability for inclusion in dietary formulations [66].

Supplementation with glycerin has emerged as a promising energy feedstuff alternative for dairy animals, evincing potential benefits in augmenting milk composition and thermal stability in dairy goats [67]. Mineral blocks and other supplements have evinced utility in enhancing milk quality and amplifying the productivity of lactating goats [68]. Furthermore, the deployment of herbal feed additives has been shown to modulate intestinal lactic acid bacteria counts in dairy goats, potentially impacting their gastrointestinal health [69].

Additionally, dietary supplements such as zinc-methionine and L-carnitine have been scrutinized for their potential to elevate milk production and bolster resistance to udder stress in dairy goats [70]. Furthermore, sodium butyrate supplementation has been probed for its capacity to mitigate apoptosis induced by high-concentrate diets in mammary cells of dairy goats [71]. The incorporation of marine algae supplementation has been shown to amplify the docosahexaenoic acid (DHA) content in milk and favorably influence the n6 to n3 ratio in dairy goat populations [72].

6. CHALLENGES AND LIMITATIONS IN OPTIMIZING FEED FORMULATION

Fine-tuning feed formulation for dairy goats poses myriad challenges and constraints that exert pronounced effects on their nutritional status, well-being, and productivity. These challenges encompass heat stress, feed sorting phenomena, and the requisite for alternative feedstuffs and supplementary nutrients. Heat stress instances have been shown to curtail feed intake and compound issues stemming from diminished energy intake, thus exerting adverse ramifications on milk yield and animal welfare [73]. Moreover, the incidence of feed sorting behaviors targeting shorter particles can engender diminished energy consumption, thereby further impinging upon the nutritional equilibrium of dairy goats [73].

Furthermore, the utilization of alternative feedstuffs and supplements can encounter impediments rooted in factors such as their availability, cost implications, and the exigency for addressing specific nutrient deficits. Environmental vicissitudes and managerial exigencies, typified by instances of heat stress, possess the capacity to disrupt feed intake and nutrient assimilation among dairy goats, thereby exerting tangible repercussions on their holistic health and productivity [3]. Additionally, the scarcity of high-yielding animals acclimated to local environments, dearth of premium-grade feeds, and logistical hurdles in transporting products to markets represent formidable challenges confronting dairy goat farmers across diverse locales, thereby encumbering their capacity to optimize feed formulation and elevate animal nutrition standards [74].

Furthermore, the imperative of remedying specific nutritional inadequacies, such as deficiencies in iron and selenium, poses formidable hurdles in devising diets attuned to the precise requisites of dairy goats [75]. The utilization of alternative feedstuffs and supplements is also circumscribed by considerations such as palatability, digestibility, and the potential for eliciting deleterious effects on milk composition and animal well-being. Additionally, the imperative of precision in formulating diets to align with the nuanced nutritional mandates of dairy goats engenders challenges in safeguarding optimal nutrient utilization and animal performance [76,77,78,79]. Moreover, the integration of supplements to redress specific nutritional lacunae may encounter constraints stemming from factors such as their availability, cost considerations, and potential interactions with other dietary constituents.

7. CONCLUSIONS

In conclusion, this review underscores the intricate interplay of genetic, environmental, and nutritional factors in optimizing feed formulation and management practices for dairy goats in intensive production systems.

Firstly, genetic factors significantly influence feed formulation strategies, as the genetic makeup of dairy goat breeds impacts milk production, composition, and overall productivity. Understanding genetic parameters and polymorphisms, such as those associated with milk protein fractions and yield, offers crucial insights into tailoring diets to specific breed requirements.

Secondly, environmental challenges, notably heat stress and fluctuating feed availability, pose substantial hurdles in achieving optimal nutrition and performance. Strategies to mitigate these challenges, including precision livestock farming and utilization of local by-products, are essential for maintaining herd health and productivity.

Thirdly, the review underscores the pivotal role of nutritional interventions, such as supplement utilization and alternative feed incorporation, in addressing specific deficiencies and enhancing overall performance. From fatty acid supplementation to herbal feed additives, a diverse array of strategies exists to optimize nutrient utilization and support dairy goat health.

Moreover, the review emphasizes the importance of precision and adaptability in feed formulation, considering factors like palatability, digestibility, and cost-effectiveness. Balancing rations to meet the evolving nutritional needs of dairy goats, particularly during critical transition stages, is paramount for sustaining productivity and profitability.

8. REFERENCES

[1] Bouattour, M., Casals, R., Albanell, E., Such, X., & Caja, G. (2008). Feeding soybean oil to dairy goats increases conjugated linoleic acid in milk. Journal of Dairy Science, 91(6), 2399-2407. https://doi.org/10.3168/jds.2007-0753

[2] Zhu, H., Ma, J., Du, R., Zheng, L., Wu, J., Song, W., … & Hua, J. (2014). Characterization of immortalized dairy goat male germline stem cells (mgscs). Journal of Cellular Biochemistry, 115(9), 1549-1560. https://doi.org/10.1002/jcb.24812

[3] Lu, C. and Miller, B. (2019). Current status, challenges and prospects for dairy goat production in the americas. Asian-Australasian Journal of Animal Sciences, 32(8), 1244-1255. https://doi.org/10.5713/ajas.19.0256

[4] Bett, R., Kosgey, I., Kahi, A., & Peters, K. (2008). Analysis of production objectives and breeding practices of dairy goats in kenya. Tropical Animal Health and Production, 41(3), 307-320. https://doi.org/10.1007/s11250-008-9191-9

[5] Romero-Huelva, M., Ramírez-Fenosa, M., Planelles-González, R., García-Casado, P., & Molina-Alcaide, E. (2017). Can by-products replace conventional ingredients in concentrate of dairy goat diet?. Journal of Dairy Science, 100(6), 4500-4512. https://doi.org/10.3168/jds.2016-11766

[6] Tsiplakou, E., Kotrotsios, V., Hadjigeorgiou, I., & Zervas, G. (2010). Differences in sheep and goats milk fatty acid profile between conventional and organic farming systems. Journal of Dairy Research, 77(3), 343-349. https://doi.org/10.1017/s0022029910000270

[7] Bampidis, V., Azimonti, G., Bastos, M., Dusemund, B., Durjava, M., Kouba, M., … & Revez, J. (2022). Assessment of the efficacy of the feed additive consisting of saccharomyces cerevisae cncm i-1077 (levucell® sc) for dairy cows, cattle for fattening, minor ruminant species and camelids (lallemand sas). Efsa Journal, 20(7). https://doi.org/10.2903/j.efsa.2022.7431

[8] Leiber, F., Arnold, N., Heckendorn, F., & Werne, S. (2020). Assessing effects of tannin-rich sainfoin supplements for grazing dairy goats on feed protein efficiency. Journal of Dairy Research, 87(4), 397-399. https://doi.org/10.1017/s0022029920000965

[9] Contreras-Jodar, A., Salama, A., Hamzaoui, S., Vailati-Riboni, M., Caja, G., & Loor, J. (2018). Effects of chronic heat stress on lactational performance and the transcriptomic profile of blood cells in lactating dairy goats. Journal of Dairy Research, 85(4), 423-430. https://doi.org/10.1017/s0022029918000705

[10] Mosolov, A., Gorlov, I., Nikolaev, D., Slozhenkina, M., Kudryashova, O., & Vasilyeva, M. (2022). Ensuring environmental safety of goat milk production based on the integration of innovations in feeding. Iop Conference Series Earth and Environmental Science, 981(2), 022098. https://doi.org/10.1088/1755-1315/981/2/022098

[11] Contreras-Jodar, A., Nayan, N., Hamzaoui, S., Caja, G., & Salama, A. (2019). Heat stress modifies the lactational performances and the urinary metabolomic profile related to gastrointestinal microbiota of dairy goats. Plos One, 14(2), e0202457. https://doi.org/10.1371/journal.pone.0202457

[12] Contreras-Jodar, A., Salama, A., Hamzaoui, S., Vailati-Riboni, M., Caja, G., & Loor, J. (2018). Effects of chronic heat stress on lactational performance and the transcriptomic profile of blood cells in lactating dairy goats. Journal of Dairy Research, 85(4), 423-430. https://doi.org/10.1017/s0022029918000705

[13] Hamzaoui, S., Caja, G., Such, X., Albanell, E., & Salama, A. (2020). Milk production and energetic metabolism of heat-stressed dairy goats supplemented with propylene glycol. Animals, 10(12), 2449. https://doi.org/10.3390/ani10122449

[14] Evan, T., Carro, M., Yepes, J., Haro, A., Arbesú, L., Romero-Huelva, M., … & Molina-Alcaide, E. (2020). Effects of feeding multinutrient blocks including avocado pulp and peels to dairy goats on feed intake and milk yield and composition. Animals, 10(2), 194. https://doi.org/10.3390/ani10020194

[15] Salama, A., Caja, G., Albanell, E., Such, X., Casals, R., & Plaixats, J. (2003). Effects of dietary supplements of zinc-methionine on milk production, udder health and zinc metabolism in dairy goats. Journal of Dairy Research, 70(1), 9-17. https://doi.org/10.1017/s0022029902005708

[16] Ghavipanje, N., Nasri, M., Farhangfar, S., Ghiasi, S., & Vargas-Bello-Pérez, E. (2021). Regulation of

nutritional metabolism in transition dairy goats: energy balance, liver activity, and insulin resistance in response to berberine supplementation. Animals, 11(8), 2236. https://doi.org/10.3390/ani11082236

[17] Nguyen, T., Chaiyabutr, N., Chanpongsang, S., & Thammacharoen, S. (2017). Dietary cation and anion difference: effects on milk production and body fluid distribution in lactating dairy goats under tropical conditions. Animal Science Journal, 89(1), 105-113. https://doi.org/10.1111/asj.12897

[18] Pereira, G., Neto, J., Gracindo, Â., Silva, Y., Difante, G., Gurgel, A., … & Lima, G. (2021). Replacement of grain maize with spineless cactus in the diet of dairy goats. Journal of Dairy Research, 88(2), 134-138. https://doi.org/10.1017/s0022029921000352

[19] Bøe, K., Ehrlenbruch, R., & Andersen, I. (2011). The preference for water nipples vs. water bowls in dairy goats. Acta Veterinaria Scandinavica, 53(1). https://doi.org/10.1186/1751-0147-53-50

[20] Hamzaoui, S., Salama, A., Albanell, E., Such, X., & Caja, G. (2013). Physiological responses and lactational performances of late-lactation dairy goats under heat stress conditions. Journal of Dairy Science, 96(10), 6355-6365. https://doi.org/10.3168/jds.2013-6665

[21] Scholtens, M., Lopez-Villalobos, N., Garrick, D., Blair, H., Lehnert, K., & Snell, R. (2019). Genetic parameters for total lactation yields of milk, fat, protein, and somatic cell score in new zealand dairy goats. Animal Science Journal, 91(1). https://doi.org/10.1111/asj.13310

[22] Xiong, J., Bao, J., Hu, W., Shang, M., & Zhang, L. (2023). Whole-genome resequencing reveals genetic diversity and selection characteristics of dairy goat. Frontiers in Genetics, 13. https://doi.org/10.3389/fgene.2022.1044017

[23] Lashmar, S., Visser, C., & Marle-Köster, E. (2016). Snp-based genetic diversity of south african commercial dairy and fiber goat breeds. Small Ruminant Research, 136, 65-71. https://doi.org/10.1016/j.smallrumres.2016.01.006

[24] Anggraeni, A., Syifa, L., Sari, O., Ishak, A., & Sumantri, C. (2021). Polymorphism of csn1s1 (g.12164g>a) and csn2 (g.8913c>a) genes in pure and cross dairy goats. Bio Web of Conferences, 33, 02001. https://doi.org/10.1051/bioconf/20213302001

[25] Mastrangelo, S., Sardina, M., Tolone, M., & Portolano, B. (2013). Genetic polymorphism at the csn1s1 gene in girgentana dairy goat breed. Animal Production Science, 53(5), 403. https://doi.org/10.1071/an12242

[26] Wang, G., Pi, X., Ji, Z., Qin, Z., Hou, L., Chao, T., … & Wang, J. (2015). Investigation of the diversity and origins of chinese dairy goats via the mitochondrial dna d-loop. Journal of Animal Science, 93(3), 949. https://doi.org/10.2527/jas.2014-8420

[27] McLaren, A., Mucha, S., Mrode, R., Coffey, M., & Conington, J. (2016). Genetic parameters of linear conformation type traits and their relationship with milk yield throughout lactation in mixed-breed dairy goats. Journal of Dairy Science, 99(7), 5516-5525. https://doi.org/10.3168/jds.2015-10269

[28] Kahi, A. and Wasike, C. (2019). Dairy goat production in sub-saharan africa: current status, constraints and prospects for research and development. Asian-Australasian Journal of Animal Sciences, 32(8), 1266-1274. https://doi.org/10.5713/ajas.19.0377

[29] Romero-Huelva, M., Ramírez-Fenosa, M., Planelles-González, R., García-Casado, P., & Molina-Alcaide, E. (2017). Can by-products replace conventional ingredients in concentrate of dairy goat diet?. Journal of Dairy Science, 100(6), 4500-4512. https://doi.org/10.3168/jds.2016-11766

[30] Sejian, V., Silpa, M., Nair, M., Devaraj, C., Krishnan, G., Bagath, M., … & Bhatta, R. (2021). Heat stress and goat welfare: adaptation and production considerations. Animals, 11(4), 1021. https://doi.org/10.3390/ani11041021

[31] Paskaš, S., Miocinovic, J., Vejnović, B., & Becskei, Z. (2019). The nutritional quality of feedstuffs used in dairy goat nutrition in vojvodina. Biotechnology in Animal Husbandry, 35(2), 163-178. https://doi.org/10.2298/bah1902163p

[32] Chilliard, Y. and Ferlay, A. (2004). Dietary lipids and forages interactions on cow and goat milk fatty acid composition and sensory properties. Reproduction Nutrition Development, 44(5), 467-492. https://doi.org/10.1051/rnd:2004052

[33] Min, B., Hart, S., Sahlu, T., & Satter, L. (2005). The effect of diets on milk production and composition, and on lactation curves in pastured dairy goats. Journal of Dairy Science, 88(7), 2604-2615. https://doi.org/10.3168/jds.s0022-0302(05)72937-4

[34] Bonanno, A., Grigoli, A., Montalbano, M., Bellina, V., Mazza, F., & Todaro, M. (2013). Effects of diet on casein and fatty acid profiles of milk from goats differing in genotype for αs1-casein synthesis. European Food Research and Technology, 237(6), 951-963. https://doi.org/10.1007/s00217-013-2069-8

[35] Wu, Z., Yang, X., Zhang, J., Wang, W., Liu, D., Hou, B., … & Xia, Y. (2023). Effects of forage type on the rumen microbiota, growth performance, carcass traits,

and meat quality in fattening goats. Frontiers in Veterinary Science, 10. https://doi.org/10.3389/fvets.2023.1147685

[36] Ghaffari, M., Tahmasbi, A., Khorvash, M., Naserian, A., & Vakili, A. (2013). Effects of pistachio by-products in replacement of alfalfa hay on ruminal fermentation, blood metabolites, and milk fatty acid composition in saanen dairy goats fed a diet containing fish oil. Journal of Applied Animal Research, 42(2), 186-193. https://doi.org/10.1080/09712119.2013.824889

[37] Muelas, R., Monllor, P., Romero, G., Sayas-Barberá, E., Navarro, C., Díaz, J., … & Sendra, E. (2017). Milk technological properties as affected by including artichoke by-products silages in the diet of dairy goats. Foods, 6(12), 112. https://doi.org/10.3390/foods6120112

[38] Chen, L., Bagnicka, E., Chen, H., & Shu, G. (2023). Health potential of fermented goat dairy products: composition comparison with fermented cow milk, probiotics selection, health benefits and mechanisms. Food & Function, 14(8), 3423-3436. https://doi.org/10.1039/d3fo00413a

[39] Levesque, J., Dion, S., Rico, D., Brassard, M., Gervais, R., & Chouinard, P. (2022). Milk yield and composition in dairy goats fed extruded flaxseed or a high-palmitic acid fat supplement. Journal of Dairy Research, 89(4), 355-366. https://doi.org/10.1017/s0022029922000784

[40] Chilliard, Y., Ferlay, A., Rouel, J., & Lamberet, G. (2003). A review of nutritional and physiological factors affecting goat milk lipid synthesis and lipolysis. Journal of Dairy Science, 86(5), 1751-1770. https://doi.org/10.3168/jds.s0022-0302(03)73761-8

[41] Gebereyowhans, S. (2023). Inclusion of microalgae in the caprine diet improves nutritional profile of milk and its camembert cheese. International Journal of Dairy Technology, 76(4), 801-812. https://doi.org/10.1111/1471-0307.12994

[42] Miroshina, T. and Chalova, N. (2023). Dairy goat breeding in russia and the world (review). E3s Web of Conferences, 380, 01004. https://doi.org/10.1051/e3sconf/202338001004

[43] Musco, N., Morittu, V., Mastellone, V., Spina, A., Vassalotti, G., D'Aniello, B., … & Lombardi, P. (2021). Effects of ecotrofin™ on milk yield, milk quality and serum biochemistry in lactating goats. Journal of Animal Physiology and Animal Nutrition, 105(S1), 26-33. https://doi.org/10.1111/jpn.13592

[44] Drackley, J. and Cardoso, F. (2014). Prepartum and postpartum nutritional management to optimize fertility in high-yielding dairy cows in confined tmr systems. Animal, 8, 5-14. https://doi.org/10.1017/s1751731114000731

[45] Broderick, G. (2018). Review: optimizing ruminant conversion of feed protein to human food protein. Animal, 12(8), 1722-1734. https://doi.org/10.1017/s1751731117002592

[46] Aguilera, J., Prieto, C., & Fonolla, J. (1990). Protein and energy metabolism of lactating granadina goats. British Journal of Nutrition, 63(2), 165-175. https://doi.org/10.1079/bjn19900104

[47] Ghavipanje, N., Nasri, M., Farhangfar, S., Ghiasi, S., & Vargas-Bello-Pérez, E. (2021). Regulation of nutritional metabolism in transition dairy goats: energy balance, liver activity, and insulin resistance in response to berberine supplementation. Animals, 11(8), 2236. https://doi.org/10.3390/ani11082236

[48] Berthel, R., Simmler, M., Dohme-Meier, F., & Keil, N. (2022). Dairy sheep and goats prefer the single components over the mixed ration. Frontiers in Veterinary Science, 9. https://doi.org/10.3389/fvets.2022.1017669

[49] Binggeli, S., Lapierre, H., Charbonneau, E., Ouellet, D., & Pellerin, D. (2021). Economic and environmental effects of revised metabolizable protein and amino acid recommendations on canadian dairy farms. Journal of Dairy Science, 104(9), 9981-9998. https://doi.org/10.3168/jds.2020-19893

[50] Fadul-Pacheco, L., Pellerin, D., Chouinard, P., Wattiaux, M., Duplessis, M., & Charbonneau, E. (2017). Nitrogen efficiency of eastern canadian dairy herds: effect on production performance and farm profitability. Journal of Dairy Science, 100(8), 6592-6601. https://doi.org/10.3168/jds.2016-11788

[51] Hassan, F., Arshad, M., Li, M., Rehman, M., Loor, J., & Huang, J. (2020). Potential of mulberry leaf biomass and its flavonoids to improve production and health in ruminants: mechanistic insights and prospects. Animals, 10(11), 2076. https://doi.org/10.3390/ani10112076

[52] Yang, W., Beauchemin, K., & Rode, L. (1999). Effects of an enzyme feed additive on extent of digestion and milk production of lactating dairy cows. Journal of Dairy Science, 82(2), 391-403. https://doi.org/10.3168/jds.s0022-0302(99)75245-8

[53] Romero-Huelva, M., Ramos-Morales, E., & Molina-Alcaide, E. (2012). Nutrient utilization, ruminal fermentation, microbial abundances, and milk yield and composition in dairy goats fed diets including tomato and cucumber waste fruits. Journal of Dairy Science, 95(10), 6015-6026. https://doi.org/10.3168/jds.2012-5573

[54] Molina-Alcaide, E., Morales-García, Y., Martín-García, A., Salem, H., Nefzaoui, A., & Sanz-Sampelayo, M. (2010). Effects of partial replacement of concentrate with feed blocks on nutrient utilization, microbial n flow, and milk yield and composition in goats. Journal of Dairy Science, 93(5), 2076-2087. https://doi.org/10.3168/jds.2009-2628

[55] Zhu, W., Xu, W., Wei, C., Zhang, Z., Jiang, C., & Chen, X. (2020). Effects of decreasing dietary crude protein level on growth performance, nutrient digestion, serum metabolites, and nitrogen utilization in growing goat kids (capra. hircus). Animals, 10(1), 151. https://doi.org/10.3390/ani10010151

[56] Patil, P., Gendley, M., Dubey, M., Dhok, A., Gade, N., & Khune, V. (2023). Effect of feeding gram straw-based complete feed pellets on the performance, nutrient utilization and rumen fermentation of goats. Asian Journal of Dairy and Food Research, (Of). https://doi.org/10.18805/ajdfr.dr-2031

[57] Chanjula, P., Pakdeechanuan, P., & Wattanasit, S. (2014). Effects of dietary crude glycerin supplementation on nutrient digestibility, ruminal fermentation, blood metabolites, and nitrogen balance of goats. Asian-Australasian Journal of Animal Sciences, 27(3), 365-374. https://doi.org/10.5713/ajas.2013.13494

[58] Hassanein, H., Maggiolino, A., El-Fadel, M., Palo, P., El-Sanafawy, H., Hussein, A., … & Salem, A. (2023). Inclusion of azolla pinnata as an unconventional feed of zaraibi dairy goats, and effects on milk production and offspring performance. Frontiers in Veterinary Science, 10. https://doi.org/10.3389/fvets.2023.1101424

[59] El-Sanafawy, H., Maggiolino, A., El-Esawy, G., Riad, W., Zeineldin, M., Abdelmegeid, M., … & Salem, A. (2023). Effect of mango seeds as an untraditional source of energy on the productive performance of dairy damascus goats. Frontiers in Veterinary Science, 10. https://doi.org/10.3389/fvets.2023.1058915

[60] Mekuriaw, S., Tsunekawa, A., Ichinohe, T., Tegegne, F., Haregeweyn, N., Kobayashi, N., … & Fievez, V. (2020). Effect of feeding improved grass hays and eragrostis tef straw silage on milk yield, nitrogen utilization, and methane emission of lactating fogera dairy cows in ethiopia. Animals, 10(6), 1021. https://doi.org/10.3390/ani10061021

[61] Moreno-Fernandez, J., López-Aliaga, I., García-Burgos, M., Alférez, M., & Díaz-Castro, J. (2019). Fermented goat milk consumption enhances brain molecular functions during iron deficiency anemia recovery. Nutrients, 11(10), 2394. https://doi.org/10.3390/nu11102394

[62] Nestares, T., Barrionuevo, M., Díaz-Castro, J., López-Aliaga, I., Alferez, M., & Campos, M. (2008). Calcium-enriched goats' milk aids recovery of iron status better than calcium-enriched cows' milk, in rats with nutritional ferropenic anaemia. Journal of Dairy Research, 75(2), 153-159. https://doi.org/10.1017/s0022029908003178

[63] Sánchez, J., Montes, P., Jiménez, A., & Andrés, S. (2007). Prevention of clinical mastitis with barium selenate in dairy goats from a selenium-deficient area. Journal of Dairy Science, 90(5), 2350-2354. https://doi.org/10.3168/jds.2006-616

[64] Almeida, V., Lima, T., Filho, G., Bom, H., Fonseca, S., Evêncio-Neto, J., … & Mendonça, F. (2022). Copper deficiency in dairy goats and kids. Pesquisa Veterinária Brasileira, 42. https://doi.org/10.1590/1678-5150-pvb-7162

[65] Medeiros, E., Queiroga, R., Oliveira, M., Medeiros, A., Sabedot, M., Bomfim, M., … & Madruga, M. (2014). Fatty acid profile of cheese from dairy goats fed a diet enriched with castor, sesame and faveleira vegetable oils. Molecules, 19(1), 992-1003. https://doi.org/10.3390/molecules19010992

[66] Yuniarti, E., Evvyernie, D., & Astuti, D. (2015). Production and energy partition of lactating dairy goats fed rations containing date fruit waste. Media Peternakan, 39(1), 27-33. https://doi.org/10.5398/medpet.2016.39.1.27

[67] Thoh, D., Pakdeechanuan, P., & Chanjula, P. (2017). Effect of supplementary glycerin on milk composition and heat stability in dairy goats. Asian-Australasian Journal of Animal Sciences, 30(12), 1711-1717. https://doi.org/10.5713/ajas.17.0066

[68] Noor, M., Rusli, N., Mat, K., Hasnita, C., & Mira, P. (2020). Milk composition and milk quality of saanen crossbreed goats supplemented by mineral blocks. Tropical Animal Science Journal, 43(2), 169-175. https://doi.org/10.5398/tasj.2020.43.2.169

[69] Foksowicz-Flaczyk, J., Wójtowski, J., Dankow, R., Mikołajczak, P., Pikul, J., Gryszczyńska, A., … & Stanisławski, D. (2022). The effect of herbal feed additives in the diet of dairy goats on intestinal lactic acid bacteria (lab) count. Animals, 12(3), 255. https://doi.org/10.3390/ani12030255

[70] Salama, A., Caja, G., Albanell, E., Such, X., Casals, R., & Plaixats, J. (2003). Effects of dietary supplements of zinc-methionine on milk production, udder health and zinc metabolism in dairy goats. Journal of Dairy Research, 70(1), 9-17. https://doi.org/10.1017/s0022029902005708

[71] Chang, G., Yan, J., Ma, N., Liu, X., Dai, H., Bilal, M., … & Shen, X. (2018). Dietary sodium butyrate supplementation reduces high-concentrate diet feeding-induced apoptosis in mammary cells in dairy goats. Journal of Agricultural and Food Chemistry, 66(9), 2101-2107. https://doi.org/10.1021/acs.jafc.7b05882

[72] Nudda, A., Cannas, A., Correddu, F., Atzori, A., Lunesu, M., Battacone, G., … & Pulina, G. (2020). Sheep and goats respond differently to feeding strategies directed to improve the fatty acid profile of milk fat. Animals, 10(8), 1290. https://doi.org/10.3390/ani10081290

[73] Mehaba, N., Salama, A., Such, X., Albanell, E., & Caja, G. (2019). Lactational responses of heat-stressed dairy goats to dietary l-carnitine supplementation. Animals, 9(8), 567. https://doi.org/10.3390/ani9080567

[74] Liang, J. and Paengkoum, P. (2019). Current status, challenges and the way forward for dairy goat production in asia – conference summary of dairy goats in asia. Asian-Australasian Journal of Animal Sciences, 32(8), 1233-1243. https://doi.org/10.5713/ajas.19.0272

[75] Battini, M., Barbieri, S., Vieira, A., Stilwell, G., & Mattiello, S. (2016). Results of testing the prototype of the awin welfare assessment protocol for dairy goats in 30 intensive farms in northern italy. Italian Journal of Animal Science, 15(2), 283-293. https://doi.org/10.1080/1828051x.2016.1150795

[76] Goetsch, A. (2019). Recent advances in the feeding and nutrition of dairy goats. Asian-Australasian Journal of Animal Sciences, 32(8), 1296-1305. https://doi.org/10.5713/ajas.19.0255

[77] Gheorghe-Irimia, R. A., Tăpăloagă, D., Tăpăloagă, P. R., Ilie, L. I., Șonea, C., Serban, A. I. (2022). Mycotoxins and Essential Oils—From a Meat Industry Hazard to a Possible Solution: A Brief Review. Foods, 11(22), 3666.

[78] Tudor, L., Pițuru, M. T., Gheorghe-Irimia, R. A., Șonea, C., & Tăpăloagă, D. (2023). Optimizing milk production, quality and safety through essential oil applications. Farmacia, 71(5)

[79] Irimia, R. A., Georgescu, M., Tudoreanu, L., & Militaru, M. (2020). Testing The Effect Of Nigella Sativa Essential Oil Solution On Chicken Breast pH And Total Volatile Base Nitrogen During Refrigeration. Scientific Works. Series C, Veterinary Medicine, 66(2).