

ELECTROSTATIC INTERACTION IN STOCHASTIC ELECTRODYNAMICS

ION SIMACIU^{1*}, ZOLTAN BORSOS², VIOREL DRAFTA³,
GHEORGHE DUMITRESCU⁴

¹Retired lecturer, Petroleum-Gas University of Ploiești, Ploiești 100680, Romania

²Petroleum-Gas University of Ploiești, Ploiești 100680, Romania

³Independent researcher

⁴High School Toma N. Socolescu, Ploiești, Romania

*Corresponding author e-mail: isimaciu@yahoo.com

Citation:

SIMACIU ION, BORSOS ZOLTAN,
DRAFTA VIOREL, DUMITRESCU
GHEORGHE

ELECTROSTATIC INTERACTION IN
STOCHASTIC ELECTRODYNAMICS

BULLETIN OF THE POLYTECHNIC
INSTITUTE OF IAȘI, 2022, 68(72), 4,
29-40;

DOI: 10.2478/bipmf-2022-0017.

Received: October 25th, 2022

Accepted: November 22, 2022

Publication: December 13, 2022

© 2022 Simaciu I. et al.

This is an Open access article licensed
under the Creative Commons
Attribution-NonCommercial –
NoDerivatives 4.0 International
License (CC BY-NV-ND 4.0)

Abstract. In this paper, the expression of the electrostatic interaction force between two charged particles is derived in the framework of Stochastic Electrodynamics. The fundamental assumption is that the electrically charged particle can be modeled as a two-dimensional oscillator that scatters the classical zero point field background radiation. The correct expression of the electrostatic force is obtained if the natural pulsation of the oscillator is equal to the Zitterbewegung angular velocity.

Keywords: Stochastic Electrodynamics (SED), classical zero-point field (CZPF), particle model; PACS: 03.50.-z; 02.50.Ey; 14.60.Cd; 01.40.-d .

1. Introduction

The Stochastic Electrodynamics (SED) studies the interaction of charged particles with the subquantum environment. In SED, the subquantum environment is supposed to be the background of electromagnetic waves with random phases. This background is the analogous of the background of

quantum fluctuations with zero temperature (or Zero point field (ZPF) in quantum electrodynamics (QED)).

Stochastic Electrodynamics was substantiated by: Brafford, P., Surdin, M., Tzara, C., Taroni, A. [1-3] as a result of the Wheeler - Feynman theory of radiation. The theory was deepened and developed by: Marshall, T. [5, 6], Boyer T. H. [7-9], L. de la Peña-Auerbach and A. M. Cetto [10, 11].

According to the analysis made by T. H. Boyer [12, 13] on SED, it is based on the principles of Classical Electrodynamics (CED) but with different boundary conditions. Although it does not use the concepts of quantum and quantification but reaches the same results obtained by QED: the van der Waals forces and Casimir effect [14-18], the Planck blackbody spectrum [19-21], the third law of thermodynamics [22], rotator and oscillator specific heats [23], the diamagnetism of free particles [24, 25], the thermal effect of acceleration [26-28], the stationary state of microscopic systems [29-31], the Lamb shift (radiative corrections) [32], etc.

This paper generalizes the particle model, in SED picture, in order to study the electrostatic interaction of two different particles. According to the model developed in [33] and improved in this paper, the electrostatic interaction between two charged particles is figured as an effect of interaction between the CZPF background and charged particles.

Since the concept of scattering cross-section must be used in this paper, then the intensity of the scattered radiation background is averaged in time and does not depend on the oscillation phases of the two oscillators [34]. For this reason, the model does not allow the explanation that there are two types of electric charge.

Both attractive and repulsive behavior of the forces may occur between oscillators if we consider that the two oscillators can oscillate in phase or in phase opposition. The phenomenon is analogous to the forces of the interaction between two oscillating bubbles, i.e. the secondary Bjerknes forces [35-37].

At the microscopic level, the electron model is characterized by a scattering cross-section dependent on angular frequency ω [33]. If the electron is at rest, the averaged electromagnetic scattering cross-section (Appendix 3) is proportional to the square of the electrostatic radius, i.e., the Thomson cross-section [38].

In the elementary charged particle model in Stochastic Electrodynamics [33], the particle at rest is an oscillator that scatters the CZPF background and the scattering cross-section depends on the angular frequency ω of the scattered electromagnetic wave and the natural angular frequency of the oscillator ω_i . In this model, electric charge is a measure of the ability of the microscopic system to

scatter the stochastic background of radiation. This interpretation of electric charge is supported by the connection between discrete electric charge, relativity and radiation thermodynamics, highlighted, in the classical (non quantum) context, in the paper [39].

2. Electrostatic force in SED

We will further model a charged particle (in particular, the electron) as a two-dimensional oscillator that scatters CZPF (Classical Zero Point Field) background radiation. Our derivation is related to the model of the non-relativistic charged oscillator [38- Sch.17.8, pp. 801-806].

The oscillator has the mass m_i and the electric charge $q_i = n_i|q_e|$, ($n_i = \frac{1}{3}, \frac{2}{3}, 1$). Its scattering cross-section with the plane electromagnetic wave [33, 38] is

$$\sigma_i(\omega) = \frac{6\pi c^2 \omega^4 \Gamma_i^2}{\omega_i^4 \left[(\omega_i^2 - \omega^2)^2 + \frac{(\Gamma_i^2 \omega^6)}{\omega_i^4} \right]}, \quad i = 1, 2. \quad (1)$$

In previous relationships, Γ_i is the radiative decay constant

$$\Gamma_i = \frac{2R_i}{3c} \omega_i^2, \quad (2)$$

and the total decay constant $\Gamma_{ti}(\omega) = \Gamma_i' + \left(\frac{\omega^2}{\omega_i^2} \right) \Gamma_i \cong \Gamma_i$ because the absorptive width (absorptive decay constant) $\Gamma_i' \ll \Gamma_i$. In Eq. (2), R_i is the electrostatic radius of the particle,

$$R_i = \frac{q_i^2}{4\pi\epsilon_0 m_i c^2} = \frac{n_i^2 e^2}{m_i c^2}. \quad (3)$$

Replacing Γ_i from Eq. (2) in Eq. (1), yields

$$\sigma_i(\omega) = \frac{8\pi R_i^2 \omega^4}{3 \left[(\omega_i^2 - \omega^2)^2 + \left(\frac{2R_i}{3c} \right)^2 \omega^6 \right]}, \quad i = 1, 2. \quad (4)$$

According to Eq. (26) of paper [33], the elementary force of interaction between two oscillators that scatter the CZPF background has the expression

$$\delta F_{12}(r, \omega) = \frac{\rho(\omega) \sigma_1(\omega) \sigma_2(\omega) d\omega}{3\pi r^2}. \quad (5)$$

The CZPF background is characterized by the spectral density of the energy [8]

$$\rho(\omega, T = 0) = \rho(\omega) = \frac{\hbar \omega^3}{2\pi^2 c^3}. \quad (6)$$

Replacing Eq. (4) and Eq. (6) in Eq. (5) and integrating, is obtained

$$F_{12}(r) = \frac{2^5 \hbar R_1^2 R_2^2}{3^3 \pi r^2 c^3} \int_0^\Omega \frac{\omega^{11} d\omega}{\left[(\omega_1^2 - \omega^2)^2 + \left(\frac{2R_1}{3c} \right)^2 \omega^6 \right] \left[(\omega_2^2 - \omega^2)^2 + \left(\frac{2R_2}{3c} \right)^2 \omega^6 \right]} = \frac{2^5 \hbar R_1^2 R_2^2}{3^3 \pi r^2 c^3} I_\Omega, \quad \Omega > \omega_2 \neq \omega_1. \quad (7)$$

The integral I_Ω in Eq. (7) is estimated according to the saddle point method [40 - Sch. 41.2, 33] that allows an analytical evaluation. We consider that the particles have the angular frequency approximately equal. This is the situation for the constituent particles of nucleons, the quarks [42]. Under these conditions, $\omega_2 \cong \omega_1 = \omega_0$, the integral I_Ω , according to Appendix 1, is

$$I_\Omega = \int_0^\Omega \frac{\omega^{11} d\omega}{\left[(\omega_0^2 - \omega^2)^2 + \left(\frac{2R_1}{3c} \right)^2 \omega^6 \right] \left[(\omega_0^2 - \omega^2)^2 + \left(\frac{2R_2}{3c} \right)^2 \omega^6 \right]} \cong \frac{3^3 \pi c^3 \omega_0}{2^4 R_1 R_2 (R_1 + R_2)}, \quad \Omega > \omega_0. \quad (8)$$

Replacing Eq. (8) in Eq. (7), yields

$$F_{12}(r) = \frac{2^5 \hbar R_1^2 R_2^2}{3^3 \pi r^2 c^3} \frac{3^3 \pi c^3 \omega_0}{2^4 R_1 R_2 (R_1 + R_2)} = \frac{2 \hbar \omega_0 R_1 R_2}{r^2 (R_1 + R_2)}. \quad (9)$$

Replacing Eq. (3) in Eq. (9), it leads to

$$F_{12}(r) = \frac{2^5 \hbar R_1^2 R_2^2}{3^3 \pi r^2 c^3} \frac{3^3 \pi c^3 \omega_0}{2^4 R_1 R_2 (R_1 + R_2)} = \left(\frac{n_1 n_2 e^2}{r^2} \right) \frac{2 \hbar \omega_0 n_1 n_2}{c^2 (n_1^2 m_2 + n_2^2 m_1)}. \quad (10)$$

The expression for the force (10) is like the expression for the Coulomb force, $F_{12}(r) = n_1 n_2 e^2 / r^2$ if

$$\frac{2 \hbar \omega_0 n_1 n_2}{c^2 (n_1^2 m_2 + n_2^2 m_1)} = 1 \text{ or } \hbar \omega_0 = \frac{c^2 (n_1^2 m_2 + n_2^2 m_1)}{2 n_1 n_2} = \frac{c^2}{2} \left(\frac{n_2}{n_1} m_1 + \frac{n_1}{n_2} m_2 \right). \quad (11)$$

The quarks are characterized by two types of mass [42, 43]: the current/naked/bare quark mass ($m_{un} = 1.8 - 2.8 \text{ MeV}/c^2 = 3.52 - 5.48 m_e$, $n_1 = n_u = \frac{2}{3}$; $m_{dn} = 4.3 - 5.2 \text{ MeV}/c^2 = 8.41 - 10.17 m_e$, $n_2 = n_d = \frac{1}{3}$) and the constituent quark mass ($m_{dc} = 340 \text{ MeV}/c^2 = 665.36 m_e$, $m_{uc} = 336 \text{ MeV}/c^2 = 657.53 m_e$). In the case of up and down quarks, the expression (11) is just the resonance condition.

When particles have different natural angular frequencies (the case of interactions between the protons, electrons, muons ($n_p = n_e = n_\mu = 1$), etc.), the integral must be averaged for the two natural angular frequencies $I_\Omega = \frac{(I_{\Omega 1} + I_{\Omega 2})}{2}$. According to the results of Appendix 2

$$I_{\Omega av} = \frac{1}{2} (I_{\Omega 1} + I_{\Omega 2}) = \frac{1}{2} \left(\frac{3 \pi c \omega_1^7}{2^2 \omega_2^4} + \frac{3^3 \pi c^3 \omega_2}{2^4 R_1^2 R_2} \right) \cong \frac{3^3 \pi c^3 \omega_2}{2^5 R_1^2 R_2}, \quad \Omega > \omega_2 > \omega_1, \quad (12)$$

Replacing Eq. (12) in Eq. (7), it follows

$$F_{12}(r) = \frac{2^5 \hbar R_1^2 R_2^2}{3^3 \pi r^2 c^3} \frac{3^3 \pi c^3 \omega_2}{2^5 R_1^2 R_2} = \frac{\hbar \omega_2 R_2}{r^2} = \left(\frac{n_1 n_2 e^2}{r^2} \right) \left(\frac{n_2 \hbar \omega_2}{n_1 m_2 c^2} \right). \quad (13)$$

The expression for the force (13) is like the expression for the Coulomb force, $F_{12}(r) = n_1 n_2 e^2 / r^2$, if

$$\frac{n_2 \hbar \omega_2}{n_1 m_2 c^2} = 1 \text{ or } \hbar \omega_2 = \frac{n_1 m_2 c^2}{n_2}. \quad (14)$$

Since for protons, electrons and muons, $n_p = n_e = n_\mu = 1$ or $n_1 = n_2 = 1$, then the condition for resonance is

$$\hbar\omega_i = m_i c^2, \quad i = 1, 2. \quad (15)$$

The asymmetry of the expression of the force (13) is the result of the approximation of the integral I_Ω by the average $I_{\Omega av} = \frac{(I_{\Omega 1} + I_{\Omega 2})}{2} \cong \frac{I_{\Omega 2}}{2}$ which was required by the necessity to obtain an analytical expression of the integrals.

3. The average scattering cross-section

At the microscopic level (relative to the CZPF background), the scattering cross-section of an electrically charged particle depends on the angular frequency ω . In order to obtain the scattering cross-section independent of the angular frequency, we will average the expression of the cross-section given by Eq. (4). According to Appendix 3, the averaged scattering cross-section has the expression

$$\langle \sigma_s \rangle_\omega = \frac{8\pi^2 c R_i \omega_i^3}{\Omega^4}. \quad (16)$$

Assuming the energy density of the CZPF radiation background, $\int_0^\infty \rho(\omega) d\omega$, is infinite, then it is necessary to enter a limit for the angular frequency, i.e., $\Omega > \omega_i$. If this is the Thomson cross-section, $\frac{8\pi R_i^2}{3}$, then the value of the maximum angular frequency Ω is

$$\Omega(\omega_i) = \omega_i \left(\frac{3\pi c}{\omega_i R_i} \right)^{\frac{1}{4}} = \omega_i \left(\frac{3\pi m_i c^3}{\omega_i n_i^2 e^2} \right)^{\frac{1}{4}} = \omega_i \left(\frac{3\pi \hbar c}{n_i^2 e^2} \right)^{\frac{1}{4}} > \omega_i \quad (17)$$

and it depends on the natural angular frequency ω_i .

This limit of the frequency must be also used to calculate numerically the integral of the interaction force between two oscillators. The difficulty that arises when calculating the integral is that this frequency is different for the two types of particles. And this occurs because the natural frequency depends on the parameters of particles. One can solve this issue by improving the particle model. We assume to be necessary to consider the phenomenon of coupling of the oscillators in interaction. Our assumption is based on the analogy with the phenomenon that occurs at the interaction of several bubbles in a cluster [44]. Also, because the oscillators have an accelerated motion, they perceive the CZPF background as a Planckian radiation background with temperatures proportional to the average acceleration [26]. The relative average acceleration of the two oscillators depends on the parameters of the two interacting particles.

4. Conclusions

In the particle model proposed in this paper, the interaction of the oscillator with the homogeneous background of CZPF radiation generates, by absorption-emission (scattering), an isotropic and inhomogeneous background. Then this primary scattering generates a secondary radiation addressed to their reciprocal scattering and which can depict the electrostatic interaction.

The source of the energy of the interaction is the energy absorbed and scattered from the background of CZPF [45]. The electric charge is a measure of the capacity of fundamental particles to scatter the CZPF radiation background. This result does not surprise us, because the SED is based on the Maxwell's equations. And these equations also involve the Coulomb force of interaction of two charges.

In our approach of the electrically charged particle is for interest to highlight the interpretation of electric charge as the scattering capacity of the CZPF background. Analogous to the quantum model (in QED, particles change energy carried by photons), particles modelled in SED change energy carried by electromagnetic waves. This phenomenon has an analogous approach in the physics of the interaction between two oscillating bubbles in liquid [35-37, 46, 47]. The electrical acoustic charge is of two types because oscillators can oscillate in phase or phase opposition. The problem of the sign for electro-acoustic charge is unresolved [48] because the acoustic particle (the oscillating bubble) has no spin. A possible solution to the problem of the internal kinetic moment of a charged acoustic particle is a model in which the oscillating bubble is captured by a vortex in the fluid [49-54]. Highlighting this property is only possible by modelling the interactions between the two oscillators in the SED formalism, without using the notion of interaction cross-section. The spin problem for the two-dimensional oscillator was solved in the paper [55]. The particle model proposed in the SED is also important for highlighting an attractive interaction when the oscillators absorb some of the scattered energy from the CZPF background. This phenomenon we will study in a forthcoming paper.

The proposed model also has shortcomings. The Zitterbewegung condition obtained in classical relativistic model of the electron [56] is $\hbar\omega_0 = 2mc^2$ and not the one used in our paper $\hbar\omega_0 = mc^2$. This discrepancy may exist because the motion of the oscillator is treated non-relativistically. Also, the model does not explain the nature of the elastic field that ensures the internal oscillation. A possible solution to this problem would be to model the charged particles as an oscillating vacuum bubble [57].

Appendix 1. The Integral I_Ω

$$\begin{aligned}
 I_\Omega &= \int_0^\Omega \frac{\omega^{11} d\omega}{\left[(\omega_0^2 - \omega^2)^2 + \left(\frac{2R_1}{3c} \right)^2 \omega^6 \right] \left[(\omega_0^2 - \omega^2)^2 + \left(\frac{2R_2}{3c} \right)^2 \omega^6 \right]} = \\
 &= \int_0^\Omega \frac{\omega^{11} d\omega}{\left[(\omega_0 - \omega)^2 (\omega_0 + \omega)^2 + \left(\frac{2R_1}{3c} \right)^2 \omega^6 \right] \left[(\omega_0 - \omega)^2 (\omega_0 + \omega)^2 + \left(\frac{2R_2}{3c} \right)^2 \omega^6 \right]} \cong \\
 &= \int_{\omega_0}^{\omega_0 - \Omega} \frac{\omega_0^{11} (-dx)}{\left[4\omega_0^2 x^2 + \left(\frac{2R_1}{3c} \right)^2 \omega_0^6 \right] \left[4\omega_0^2 x^2 + \left(\frac{2R_2}{3c} \right)^2 \omega_0^6 \right]} = \frac{(-\omega_0^7)}{2^4} \int_{\omega_0}^{\omega_0 - \Omega} \frac{dx}{\left[x^2 + \left(\frac{R_1 \omega_0^2}{3c} \right)^2 \right] \left[x^2 + \left(\frac{R_2 \omega_0^2}{3c} \right)^2 \right]} = \\
 &= \frac{(-\omega_0^7)}{2^4} \int_{\omega_0}^{\omega_0 - \Omega} \frac{dx}{x^4 + x^2 \left(\frac{\omega_0^2}{3c} \right)^2 (R_1^2 + R_2^2) + \left[R_1 R_2 \left(\frac{\omega_0^2}{3c} \right)^2 \right]^2} = \frac{(-\omega_0^7)}{2^4} I_x, \omega_0 - \omega = x, \Omega > \omega_0. \quad (A1.1)
 \end{aligned}$$

The I_x integral is solved using formulas 2.161 and 2.103 of the book [41]

$$\begin{aligned}
 I_x &= \int_{\omega_0}^{\omega_0 - \Omega} \frac{dx}{x^4 + x^2 \left(\frac{\omega_0^2}{3c} \right)^2 (R_1^2 + R_2^2) + \left[R_1 R_2 \left(\frac{\omega_0^2}{3c} \right)^2 \right]^2} = \\
 &= \frac{2}{\left(\frac{\omega_0^2}{3c} \right)^2 (R_1^2 - R_2^2)} \left[\int_{\omega_0}^{\omega_0 - \Omega} \frac{dx}{2x^2 + 2R_2^2 \left(\frac{\omega_0^2}{3c} \right)^2} - \int_{\omega_0}^{\omega_0 - \Omega} \frac{dx}{2x^2 + 2R_1^2 \left(\frac{\omega_0^2}{3c} \right)^2} \right] = \\
 &= \frac{1}{\left(\frac{\omega_0^2}{3c} \right)^2 (R_1^2 - R_2^2)} \left[\int_{\omega_0}^{\omega_0 - \Omega} \frac{dx}{x^2 + \left(\frac{R_2 \omega_0^2}{3c} \right)^2} - \int_{\omega_0}^{\omega_0 - \Omega} \frac{dx}{x^2 + \left(\frac{R_1 \omega_0^2}{3c} \right)^2} \right] = \\
 &= \frac{1}{\left(\frac{\omega_0^2}{3c} \right)^2 \omega_0^2 (R_1^2 - R_2^2)} \left[\frac{3c}{R_2 \omega_0^2} \arctan \frac{3cx}{R_2 \omega_0^2} \Big|_{\omega_0}^{\omega_0 - \Omega} - \frac{3c}{R_1 \omega_0^2} \arctan \frac{3cx}{R_1 \omega_0^2} \Big|_{\omega_0}^{\omega_0 - \Omega} \right] = \\
 &= \frac{(-3^3 \pi c^3)}{\omega_0^6 (R_1^2 - R_2^2)} \left(\frac{R_1 - R_2}{R_1 R_2} \right) = \frac{(-3^3 \pi c^3)}{\omega_0^6 R_1 R_2 (R_1 + R_2)} \frac{3c}{R_i \omega_0} \gg 1, \frac{3c\Omega}{R_i \omega_0^2} \gg 1, \Omega > \omega_0. \quad (A1.2)
 \end{aligned}$$

$$h = \left(\frac{\omega_0^2}{3c} \right)^4 (R_1^2 + R_2^2)^2 - 4 \left[R_1 R_2 \left(\frac{\omega_0^2}{3c} \right)^2 \right]^2 = \left(\frac{\omega_0^2}{3c} \right)^4 (R_1^2 - R_2^2)^2 > 0,$$

$$\left(\frac{\omega_0^2}{3c} \right)^2 (R_1^2 + R_2^2) - \sqrt{h} = 2R_2^2 \left(\frac{\omega_0^2}{3c} \right)^2$$

$$\left(\frac{\omega_0^2}{3c} \right)^2 (R_1^2 + R_2^2) + \sqrt{h} = 2R_1^2 \left(\frac{\omega_0^2}{3c} \right)^2, \text{ if } R_1 \neq R_2. \quad (A1.3)$$

From the conditions $\frac{3c}{R_i \omega_0} \gg 1$ and $\frac{3c\Omega}{R_i \omega_0^2} \gg 1$, yields

$$\begin{aligned}\frac{3c}{R_i \omega_0} &= \frac{3}{n_i^2} \frac{\hbar c}{e^2} \frac{m_i c^2}{\hbar \omega_0} \gg 1 \text{ or } \frac{m_i c^2}{\hbar \omega_0} \gg \frac{n_i^2 e^2}{3 \hbar c}, \\ \frac{3c\Omega}{R_i \omega_0^2} &= \frac{3}{n_i^2} \frac{\hbar c}{e^2} \frac{m_i c^2}{\hbar \omega_0} \frac{\Omega}{\omega_0} \gg 1 \text{ or } \frac{m_i c^2}{\hbar \omega_0} \frac{\Omega}{\omega_0} \gg \frac{n_i^2 e^2}{3 \hbar c}.\end{aligned}\quad (\text{A1.4})$$

Appendix 2. The Integrals $I_{\Omega 1}$ and $I_{\Omega 2}$

$$\begin{aligned}I_{\Omega 1} &= \int_0^{\Omega} \frac{\omega^{11} d\omega}{[(\omega_1^2 - \omega^2)^2 + \left(\frac{2R_1}{(3c)}\right)^2 \omega^6]} = \int_{\omega_1}^{\omega_1 - \Omega} \frac{\omega_1^{11} (-dx)}{4\omega_1^6 \left(x^2 + \left(\frac{R_1 \omega_1^2}{(3c)}\right)^2\right) \left[\left(\frac{\omega_2^2}{\omega_1^2} - 1\right)^2 + \left(\frac{2R_2 \omega_1}{(3c)}\right)^2\right]} = \\ &= \frac{\omega_1^5}{4 \left[\left(\frac{\omega_2^2}{\omega_1^2} - 1\right)^2 + \left(\frac{2R_2 \omega_1}{(3c)}\right)^2\right]} \int_{\omega_1}^{\omega_1 - \Omega} \frac{dx}{\left[x^2 + \left(\frac{R_1 \omega_1^2}{(3c)}\right)^2\right]} \cong \frac{\omega_1^5}{4 \left[\left(\frac{\omega_2^2}{\omega_1^2} - 1\right)^2 + \left(\frac{2R_2 \omega_1}{(3c)}\right)^2\right]} \left[\frac{3c}{R_1 \omega_1^2} \arctan \frac{3cx}{R_1 \omega_1^2} \right]_{\omega_1}^{\omega_1 - \Omega} \cong \\ &\cong \frac{(-\omega_1^5)}{4 \left[\left(\frac{\omega_2^2}{\omega_1^2} - 1\right)^2 + \left(\frac{2R_2 \omega_1}{(3c)}\right)^2\right]} \left(\frac{-3c\pi}{R_1 \omega_1^2} \right) = \frac{3\pi c \omega_1^3}{2^2 \left[\left(\frac{\omega_2^2}{\omega_1^2} - 1\right)^2 + \left(\frac{2R_2 \omega_1}{(3c)}\right)^2\right]} \cong \frac{3\pi c \omega_1^7}{2^2 \omega_2^4}, \\ \omega_1 - \omega &= x, \quad \Omega > \omega_2 > \omega_1, \quad \frac{3c}{R_1 \omega_1} \gg 1, \quad \frac{3c\Omega}{R_1 \omega_1^2} \gg 1, \quad \omega_2^4 \gg \left(\frac{2R_2 \omega_1}{(3c)}\right)^2.\end{aligned}\quad (\text{A2.1})$$

$$\begin{aligned}I_{\Omega 2} &= \int_{\omega_2}^{\omega_2 - \Omega} \frac{\omega_1^{11} (-dy)}{4\omega_2^6 \left[y^2 + \left(\frac{R_2 \omega_2^2}{(3c)}\right)^2\right] \left[\left(\frac{\omega_1^2}{\omega_2^2} - 1\right)^2 + \left(\frac{2R_1 \omega_2}{(3c)}\right)^2\right]} = \frac{\omega_1^5}{2^2 \left[\left(\frac{\omega_1^2}{\omega_2^2} - 1\right)^2 + \left(\frac{2R_1 \omega_2}{(3c)}\right)^2\right]} \int_{\omega_2}^{\omega_2 - \Omega} \frac{dy}{\left[y^2 + \left(\frac{R_2 \omega_2^2}{(3c)}\right)^2\right]} \cong \\ &\cong \frac{\omega_2^5}{2^2 \left[\left(\frac{\omega_1^2}{\omega_2^2} - 1\right)^2 + \left(\frac{2R_1 \omega_2}{(3c)}\right)^2\right]} \left[\frac{3c}{R_2 \omega_2^2} \arctan \frac{3cy}{R_2 \omega_2^2} \right]_{\omega_2}^{\omega_2 - \Omega} \cong \frac{(-\omega_2^5)}{2^2 \left[\left(\frac{\omega_1^2}{\omega_2^2} - 1\right)^2 + \left(\frac{2R_1 \omega_2}{(3c)}\right)^2\right]} \left(\frac{-3c\pi}{R_2 \omega_2^2} \right) = \\ &\cong \frac{3\pi c \omega_2^3}{2^2 \left[\left(\frac{\omega_1^2}{\omega_2^2} - 1\right)^2 + \left(\frac{2R_1 \omega_2}{(3c)}\right)^2\right]} \cong \frac{3^3 \pi c^3 \omega_2}{2^4 R_1^2 R_2}, \\ \omega_2 - \omega &= y, \quad \Omega > \omega_2 > \omega_1, \quad \frac{3c}{R_2 \omega_2} \gg 1, \quad \frac{3c\Omega}{R_2 \omega_2^2} \gg 1, \quad \left(\frac{2R_1 \omega_2}{(3c)}\right)^2 \gg 1.\end{aligned}\quad (\text{A2.2})$$

The integrals, in the variables x and y , are solved according to the formulas 2.103 of the book [41, p. 67]

Appendix 3. The averaged scattering cross-section

$$\langle \sigma_s \rangle_{\omega} = \frac{\int_0^{\Omega} \sigma_{si}(\omega) \rho(\omega) d\omega}{\int_0^{\Omega} \rho(\omega) d\omega} = \frac{\int_0^{\Omega} \left[\frac{8\pi R_i^2 \omega^4}{3 \left[(\omega_i^2 - \omega^2)^2 + \left(\frac{2R_i}{(3c)}\right)^2 \omega^6 \right]} \frac{\hbar \omega^3}{2\pi^2 c^3} \right] d\omega}{\int_0^{\Omega} \frac{\hbar \omega^3}{2\pi^2 c^3} d\omega}, \quad \Omega > \omega_i. \quad (\text{A3.1})$$

To calculate the integral from the denominator, we use the property that the section has a maximum for the angular frequencies close to its natural angular frequency. In integral from the denominator we make the change of variable $\omega_i - \omega = x$ and it becomes

$$\begin{aligned} \langle \sigma_s \rangle_\omega &= \frac{\int_0^\Omega \frac{8\pi R_l^2 \omega^7 d\omega}{3 \left[(\omega_i^2 - \omega^2)^2 + \left(\frac{2R_l}{3c} \right)^2 \omega^6 \right]}}{\int_0^\Omega \omega^3 d\omega} = \frac{\int_0^\Omega \frac{8\pi R_l^2 \omega^7 d\omega}{3 \left[(\omega_i^2 - \omega^2)^2 + \left(\frac{2R_l}{3c} \right)^2 \omega^6 \right]}}{\frac{\Omega^4}{4}} = \frac{32\pi c R_l^2}{3\Omega^4} \int_0^\Omega \frac{\omega^7 d\omega}{\left[(\omega_i - \omega)^2 (\omega_i + \omega)^2 + \left(\frac{2R_l}{3c} \right)^2 \omega^6 \right]} \cong \\ &= \frac{32\pi c R_l^2 \omega_i^5}{3\Omega^4} \int_{\omega_i}^{\omega_i - \Omega} \frac{(-dx)}{4\omega_i^2 \left[x^2 + \left(\frac{\omega_i^2 R_l}{3c} \right)^2 \right]} \cong \frac{(-8\pi c R_l^2 \omega_i^5)}{3\Omega^4} \left(\frac{-3\pi c}{\omega_i^2 R_l} \right) = \frac{8\pi^2 c R_l \omega_i^3}{\Omega^4}. \end{aligned} \quad (A3.2)$$

The integral, in the variable x , are solved according to the formulas 2.103 of the book [41, p. 67].

REFERENCES

- [1] Braffort P. and Tzara C., *Energie de l'oscillateur harmonique dans le vide*, C. R. Acad. Sci. Paris, **239**, p. 1779, (1954).
- [2] Braffort P., Surdin M. and Taroni A., *L'énergie Moyenne d'un Oscillateur Harmonique Non Relativiste en Electrodynamique Aleatoire*, C. R. Acad. Sci., **261**, p. 4339, (1965).
- [3] Surdin M., *Stochastic Electrodynamics an Overview*. In: van der Merwe A. (eds) *Old and New Questions in Physics, Cosmology, Philosophy, and Theoretical Biology*. Springer, Boston, MA. (1983), https://doi.org/10.1007/978-1-4684-8830-2_33
- [4] Wheller I.A. and Feynman R., *Interaction with the Absorber as the Mecanism of Radiation*, Rev., Mod. Phys., **17**, 157, (1959).
- [5] Marshall T.W., *Random electrodynamics*, Proc. R. Soc. London, Ser. A, **276**, 475, (1963).
- [6] Marshall T.W., *Statistical electrodynamics*, Proc. Cambridge Phil. Soc. **61**, 537, (1965).
- [7] Boyer T.H., *Quantum electromagnetic zero-point energy and retarded dispersion forces*, Phys. Rev., **174**, 1631 (1968).
- [8] Boyer T.H., *A Brief Survey of Stochastic Electrodynamics*, In: Barut A. O. (Eds.) *Foundations of Radiation Theory and Quantum Electrodynamics*. Springer, Boston, MA., (1980), <https://doi.org/10.1007/978-1-4757-0671-05>
- [9] Boyer T.H., *Stochastic Electrodynamics: The Closest Classical Approximation to Quantum Theory*, Atoms. **7**, 1, 29-39, (2019).
- [10] de la Peña-Auerbach L. and Cetto A.M., *Quantum mechanics derived from stochastic electrodynamics*, Found. Phys., **8**, 3-4, 191, (1978).
- [11] de la Peña-Auerbach L. and Cetto A.M., *The quantum dice: an introduction to stochastic electrodynamics*, Kluwer Academic Publishers, Dordrecht, The Netherlands, p. 528, (1996), ISBN 0-7923-3818-9.
- [12] Boyer T.H., *Random Electrodynamics: The theory of classical electrodynamics with classical electromagnetic zero-point radiation*, Phys. Rev. D, **11**, 790, (1975).

- [13] Boyer T.H., *General Connection Between Random Electrodynamics and Quantum Electrodynamics for Free Electromagnetic Fields and for Dipol Oscillator Systems*, Phys. Rev. D, **11**, 809, (1975).
- [14] Marshall T.W., *A classical treatment of blackbody radiation*, Nuovo Cimento, **38**, 1, pp. 206-215, (1965).
- [15] Boyer T.H., *Asymptotic Retarded van der Waals Forces Derived from Classical Electrodynamics with Classical Electromagnetic Zero-Point Radiation*, Phys. Rev. A, **5**, p. 1799, (1972).
- [16] Boyer T.H., *Van der Waals forces and zero-point energy for dielectric and permeable materials*, Phys. Rev. A, **9**, p. 2078, (1974).
- [17] Boyer T.H., *Retarded van der Waals forces at all distances derived from classical electrodynamics with classical electromagnetic zero-point radiation*, Phys. Rev. A, **7**, p. 1832, (1973).
- [18] Boyer T.H., *Quantum Electromagnetic Zero - Point Energy of Conducting Spherical Shell and Casimir Model for a Charged Particle*, Phys. Rev., **174**, p. 1764, (1968).
- [19] Surdin M., Braffort P. and Taroni A., *Black-body radiation law deduced from stochastic electrodynamics*, Nature 210, 405, (1966).
- [20] Boyer T.H., *Derivation of the blackbody radiation spectrum from the equivalence principle in classical physics with classical electromagnetic zeropoint radiation*, Phys. Rev. D29, 1096, (1984).
- [21] Boyer T.H., *Thermodynamics of the harmonic oscillator: derivation of the Planck blackbody spectrum from pure thermodynamics*, European Journal of Physics, Volume 40, Issue 2, p. 025101, (2019).
- [22] Boyer T.H., *Third Law of Thermodynamics and Electromagnetic Zero-Point Radiation*, Phys. Rev. D1, 1526, (1970).
- [23] Boyer T.H., *Specific Heat of a Classical, Plane, Rigid, Dipole Rotator in Electromagnetic Zero-Point Radiation*, Phys. Rev. D1, 2257, (1970).
- [24] Boyer T.H., *Diamagnetism of a free particle in classical electron theory with classical electromagnetic zero-point radiation*, Phys. Rev. A 21, 66, (1980).
- [25] de la Peña-Auerbach L. and Jauregui A., *Stochastic electrodynamics for the free particle*, J. Math. Phys.24, 2751, (1983).
- [26] Boyer T.H., *Thermal effects of acceleration for a classical dipole oscillator in classical electromagnetic zero-point radiation*, Phys. Rev. D 29, 1089, (1984).
- [27] Cole D.C., *Thermal Effects of Acceleration for a Spatially Extended Electromagnetic System in Classical Electromagnetic Zero Point Radiation: Transversely Positioned Classical Oscillators*, Phys. Rev. D 35, 562-583, (1987).
- [28] Boyer T.H., *Classical and Quantum Interpretations Regarding Thermal Behavior in a Coordinate Frame Accelerating Through Zero-Point Radiation*, arXiv:1011.1426v1, (2010).
- [29] Puthoff H.E., *Ground state of hydrogen as a zero point fluctuation determined state*, Phys. Rev. D 35, 3266, (1987).
- [30] Cole and Zou Y., *Quantum mechanical ground state of hydrogen obtained from classical electrodynamics*, Physics Letters A, Vol. 317, No. 1-2, pp. 14-20, (2003).
- [31] Nieuwenhuizen Th.M., *On the Stability of Classical Orbits of the Hydrogen Ground State in Stochastic Electrodynamics*, Entropy, 18, 135, (2016).
- [32] Cetto A.M., de la Peña-Auerbach L. and Valdés-Hernández A., *Atomic radiative corrections without QED: role of the zero-point field*, Rev. Mex. Fis. vol. 59, no. 5, 433–443, (2013).

- [33] Simaciu I. and Ciubotariu C., *Classical model of Electron in Stochastic Electrodynamics*, Rev. Mex. Fis. 47(4), 392, (2001).
- [34] Simaciu I., Dumitrescu Gh. and Dinu M., *New Results in Stochastic Physics*, Romanian Reports in Physics 47, 537, (1995).
- [35] Barbat T., Ashgriz N. and Liu C.S., *Dynamics of two interacting bubbles in an acoustic field*, J. Fluid Mech. 389, 137, (1999).
- [36] Doinikov A.A., *Recent Research Developments in Acoustics*, Vol. 1, (Transworld Research Network, Trivandrum, Kerala, 2003), pp. 39-67.
- [37] Ainslie M.A. and Leighton T.G., *Review of scattering and extinction cross-sections, damping factors and resonance frequencies of a spherical gas bubble*, J. Acoust. Soc. Am. 130 (5), Pt. 2 (2011) Pub12667, <https://doi.org/10.1121/1.3628321>.
- [38] Jackson J.D., *Classical Electrodynamics*, 2nd ed., Wiley, New York, (1975).
- [39] Boyer T.H., *Connecting blackbody radiation, relativity, and discrete charge in classical electrodynamics*, Found. Phys. 37, p. 999, (2007), <https://doi.org/10.1007/s10701-007-9139-3>.
- [40] Feynman R.P., Leighton R.B. and Matthew S., *The Feynman Lectures on Physics*, Vol. 1, Massachusetts: Addison-Wesley, (1964).
- [41] Gradshteyn I.S. and Ryzhik I.M., *Table of Integrals, Series, and Products*, 7th Edition, Jeffrey, Alan and Zwillinger, Daniel (eds.), Academic Press, (2007).
- [42] Griffiths D., *Introduction to Elementary Particles*, Wiley-Vch., (2008).
- [43] Patrignani C. et al. (Particle Data Group), (2016), "*Quarks*", Chin. Phys. C. 40: 100001.
- [44] Simaciu I., Dumitrescu Gh. and Borsos Z., *Mach's Principle in the Acoustic World*, arXiv:1907.05713, (2019).
- [45] Țiplea G., Simaciu I., Milea P.L. and Șchiopu P., *Extraction of energy from the vacuum in SED: theoretical and technological models and limitations*, UPB Scientific Bulletin, Series A: Applied Mathematics and Physics 83(2), pp. 267-286, 2021.
- [46] Simaciu I., Borsos Z., Dumitrescu Gh., Silva G.T. and Bărbat T., *The acoustic force of electrostatic type*, Bul. Inst. Politeh. Iași, Sect. Mat., Mec. teor., Fiz. 65 (69), No 2, (2019) 17; arXiv:1711.03567v1.
- [47] Zavtrak S.T., *A classical treatment of the long-range radiative interaction of small particles*, Journal of Physics A, General Physics 23 (9), 1493, (1999).
- [48] Dinu I., *Fundamentals of a Theory of Aether – Part 2*, 2021.
- [49] Uwe R. Fischer, *Motion of Quantized Vortices as Elementary Objects*, Annals of Physics, Volume 278, Issue 1, pp. 62-85, (1999).
- [50] Allen J.J., Jouanne Y., Shashikanth B.N., *Vortex interaction with a moving sphere*, Journal of Fluid Mechanics, Cambridge, Vol. 587, pp. 337-346, 2007.
- [51] Chena L., Zhanga L., Shaoa X.M., *The motion of small bubble in the ideal vortex flow*, Procedia Engineering 126, pp. 228-231, 2015.
- [52] Cui B., Ni B. and Wu Q., *Bubble-bubble interaction effects on dynamics of multiple bubbles in a vortical flow field*, Advances in Mechanical Engineering, Vol. 8(2), pp. 1-12, 2016.
- [53] Butto N., *A New Theory on Electron Wave-Particle Duality*, Journal of High Energy Physics, Gravitation and Cosmology, 6, pp. 567-578, 2020.

- [54] Ruban V.P., *Bubbles with Attached Quantum Vortices in Trapped Binary Bose–Einstein Condensates*, Journal of Experimental and Theoretical Physics, Vol. 133, pp. 779–785, 2021.
- [55] de la Peña-Auerbach L. and Cetto A.M. and Valdés-Hernández A., *Proposed physical explanation for the electron spin and related antisymmetry*, Quantum Studies: Mathematics and Foundations volume 6, pp. 45–53, (2019).
- [56] Barut A.O. and Zanghi N., *Classical Model of the Dirac Electron*, Phys. Rev. Lett. 52, pp. 2009-2012, (1984).
- [57] Burinskii A., *What tells Gravity on the shape and size of an electron*, Physics of Particles and Nuclei 45(1), (2012).