FORMALIZED MATHEMATICS
Vol. 30, No. 1, Pages 1–12, 2022
ISSN: 1426–2630, e-ISSN: 1898–9934
DOI: 10.2478/forma-2022-0001

sciendo
sciendo.com/journal/forma

Intuitionistic Propositional Calculus in the Extended Framework with Modal Operator. Part II

Takao Inoué
Department of Medical Molecular Informatics
Meiji Pharmaceutical University
Tokyo, Japan
Graduate School of Science and Engineering
Hosei University, Tokyo, Japan
Department of Applied Informatics
Faculty of Science and Engineering
Hosei University, Tokyo, Japan

Riku Hanaoka
Keyaki-Sou 403
Midori-cho 5-17-27
Koganei-city
184-0003, Tokyo
Japan

Summary. This paper is a continuation of Inoué [5]. As already mentioned in the paper, a number of intuitionistic provable formulas are given with a Hilbert-style proof. For that, we make use of a family of intuitionistic deduction theorems, which are also presented in this paper by means of Mizar system [2], [1]. Our axiom system of intuitionistic propositional logic IPC is based on the propositional subsystem of H_1-**IQC** in Troelstra and van Dalen [6, p. 68]. We also owe Heyting [4] and van Dalen [7]. Our treatment of a set-theoretic intuitionistic deduction theorem is due to Agata Darmochwał's Mizar article "Calculus of Quantifiers. Deduction Theorem" [3].

MSC: 03B20 03F03 68V20

Keywords: intuitionistic logic; deduction theorem; consequence operator

MML identifier: INTPRO_2, version: 8.1.12 5.71.1431

1. The Notion of Proof in Intuitionistic Setting

From now on i, j, n, k, l denote natural numbers, T, S, X, Y, Z denote subsets of MC-w.f.f., p, q, r, t, F, H, G denote elements of MC-w.f.f., and s, U, V denote MC-formulas.

Let p, q be elements of MC-w.f.f.. The functor $p \Leftrightarrow q$ yielding an element of MC-w.f.f. is defined by the term

© 2022 The Author(s) / AMU
(Association of Mizar Users)
under CC BY-SA 3.0 license

(Def. 1) $(p \Rightarrow q) \wedge (q \Rightarrow p)$.

The functor Proof-Step-Kinds-IPC yielding a set is defined by the term

(Def. 2) $\{k : k \leqslant 10\}$.

Now we state the proposition:

(1) (i) $0 \in$ Proof-Step-Kinds-IPC and ... and

(ii) $10 \in$ Proof-Step-Kinds-IPC.

One can verify that Proof-Step-Kinds-IPC is non empty and Proof-Step-Kinds-IPC is finite.

From now on f, g denote finite sequences of elements of MC-w.f.f. $\times$ Proof-Step-Kinds-IPC. Now we state the proposition:

(2) Let us consider a natural number n. If $1 \leqslant n \leqslant \operatorname{len} f$, then $(f(n))_{\mathbf{2}} = 0$ or ... or $(f(n))_{\mathbf{2}} = 10$.

Let P_1 be a finite sequence of elements of MC-w.f.f. $\times$ Proof-Step-Kinds-IPC and n be a natural number. Let us consider X. We say that P_1 is a correct n-th step w.r.t. IPC (X) if and only if

(Def. 3) (i) $(P_1(n))_{\mathbf{1}} \in X$, **if** $(P_1(n))_{\mathbf{2}} = 0$,

(ii) there exists p and there exists q such that $(P_1(n))_{\mathbf{1}} = p \Rightarrow (q \Rightarrow p)$, **if** $(P_1(n))_{\mathbf{2}} = 1$,

(iii) there exists p and there exists q and there exists r such that $(P_1(n))_{\mathbf{1}} = p \Rightarrow (q \Rightarrow r) \Rightarrow (p \Rightarrow q \Rightarrow (p \Rightarrow r))$, **if** $(P_1(n))_{\mathbf{2}} = 2$,

(iv) there exists p and there exists q such that $(P_1(n))_{\mathbf{1}} = p \wedge q \Rightarrow p$, **if** $(P_1(n))_{\mathbf{2}} = 3$,

(v) there exists p and there exists q such that $(P_1(n))_{\mathbf{1}} = p \wedge q \Rightarrow q$, **if** $(P_1(n))_{\mathbf{2}} = 4$,

(vi) there exists p and there exists q such that $(P_1(n))_{\mathbf{1}} = p \Rightarrow (q \Rightarrow p \wedge q)$, **if** $(P_1(n))_{\mathbf{2}} = 5$,

(vii) there exists p and there exists q such that $(P_1(n))_{\mathbf{1}} = p \Rightarrow p \vee q$, **if** $(P_1(n))_{\mathbf{2}} = 6$,

(viii) there exists p and there exists q such that $(P_1(n))_{\mathbf{1}} = q \Rightarrow p \vee q$, **if** $(P_1(n))_{\mathbf{2}} = 7$,

(ix) there exists p and there exists q and there exists r such that $(P_1(n))_{\mathbf{1}} = p \Rightarrow r \Rightarrow (q \Rightarrow r \Rightarrow (p \vee q \Rightarrow r))$, **if** $(P_1(n))_{\mathbf{2}} = 8$,

(x) there exists p such that $(P_1(n))_{\mathbf{1}} = \text{FALSUM} \Rightarrow p$, **if** $(P_1(n))_{\mathbf{2}} = 9$,

(xi) there exists i and there exists j and there exists p and there exists q such that $1 \leqslant i < n$ and $1 \leqslant j < i$ and $p = (P_1(j))_{\mathbf{1}}$ and $q = (P_1(n))_{\mathbf{1}}$ and $(P_1(i))_{\mathbf{1}} = p \Rightarrow q$, **if** $(P_1(n))_{\mathbf{2}} = 10$.

Let us consider f. We say that f is a proof w.r.t. IPC (X) if and only if

(Def. 4) $f \neq \emptyset$ and for every n such that $1 \leqslant n \leqslant \operatorname{len} f$ holds f is a correct n-th step w.r.t. IPC (X).

Now we state the propositions:

(3) If f is a proof w.r.t. IPC (X), then $\operatorname{rng} f \neq \emptyset$.

(4) If f is a proof w.r.t. IPC (X), then $1 \leqslant \operatorname{len} f$.

(5) If f is a proof w.r.t. IPC (X), then $(f(1))_{\mathbf{2}} = 0$ or ... or $(f(1))_{\mathbf{2}} = 10$. The theorem is a consequence of (4) and (2).

(6) If $1 \leqslant n \leqslant \operatorname{len} f$, then f is a correct n-th step w.r.t. IPC (X) iff $f \frown g$ is a correct n-th step w.r.t. IPC (X).

PROOF: If f is a correct n-th step w.r.t. IPC (X), then $f \frown g$ is a correct n-th step w.r.t. IPC (X). $(f(n))_{\mathbf{2}} = 0$ or ... or $(f(n))_{\mathbf{2}} = 10$. □

(7) If $1 \leqslant n \leqslant \operatorname{len} g$ and g is a correct n-th step w.r.t. IPC (X), then $f \frown g$ is a correct $n + \operatorname{len} f$-th step w.r.t. IPC (X). The theorem is a consequence of (2).

(8) If f is a proof w.r.t. IPC (X) and g is a proof w.r.t. IPC (X), then $f \frown g$ is a proof w.r.t. IPC (X). The theorem is a consequence of (6) and (7).

(9) If f is a proof w.r.t. IPC (X) and $X \subseteq Y$, then f is a proof w.r.t. IPC (Y). The theorem is a consequence of (2).

(10) If f is a proof w.r.t. IPC (X) and $1 \leqslant l \leqslant \operatorname{len} f$, then $(f(l))_{\mathbf{1}} \in \operatorname{CnIPC}(X)$.

PROOF: For every n such that $1 \leqslant n \leqslant \operatorname{len} f$ holds $(f(n))_{\mathbf{1}} \in \operatorname{CnIPC}(X)$. □

Let us consider f. Assume $f \neq \emptyset$. The functor Effect-IPC(f) yielding an element of MC-w.f.f. is defined by the term

(Def. 5) $(f(\operatorname{len} f))_{\mathbf{1}}$.

Now we state the proposition:

(11) If f is a proof w.r.t. IPC (X), then Effect-IPC$(f) \in \operatorname{CnIPC}(X)$. The theorem is a consequence of (4) and (10).

2. A CONSEQUENCE AS A SET OF ALL INTUITIONISTIC PROVABLE FORMULAS

Now we state the proposition:

(12) $X \subseteq \{F :$ there exists f such that f is a proof w.r.t. IPC (X) and Effect-IPC$(f) = F\}$. The theorem is a consequence of (1).

Let us consider X. Now we state the propositions:

(13) Suppose $Y = \{p :$ there exists f such that f is a proof w.r.t. IPC (X) and Effect-IPC$(f) = p\}$. Then Y is IPC theory.

(14) $\{p :$ there exists f such that f is a proof w.r.t. IPC (X) and Effect-IPC $(f) = p\} = \mathrm{CnIPC}(X)$. The theorem is a consequence of (12) and (13).

(15) $p \in \mathrm{CnIPC}(X)$ if and only if there exists f such that f is a proof w.r.t. IPC (X) and Effect-IPC$(f) = p$. The theorem is a consequence of (14).

(16) If $p \in \mathrm{CnIPC}(X)$, then there exists Y such that $Y \subseteq X$ and Y is finite and $p \in \mathrm{CnIPC}(Y)$.

PROOF: Consider f such that f is a proof w.r.t. IPC (X) and Effect-IPC$(f) = p$. Consider A being a set such that A is finite and $A \subseteq$ MC-w.f.f. and $\operatorname{rng} f \subseteq A \times$ Proof-Step-Kinds-IPC. If $1 \leqslant n \leqslant \operatorname{len} f$, then f is a correct n-th step w.r.t. IPC (Y). □

3. The Intuitionistic Provable Relation

Let us consider X and s. We say that $X \vdash_{IPC} (s)$ if and only if

(Def. 6) $s \in \mathrm{CnIPC}(X)$.

We say that $\vdash_{IPC} s$ if and only if

(Def. 7) $\emptyset_{\text{MC-w.f.f.}} \vdash_{IPC} s$.

Now we state the propositions:

(17) $X \vdash_{IPC} (p \Rightarrow (q \Rightarrow p))$.

(18) $X \vdash_{IPC} (p \Rightarrow (q \Rightarrow r) \Rightarrow (p \Rightarrow q \Rightarrow (p \Rightarrow r)))$.

(19) $X \vdash_{IPC} (p \wedge q \Rightarrow p)$.

(20) $X \vdash_{IPC} (p \wedge q \Rightarrow q)$.

(21) $X \vdash_{IPC} (p \Rightarrow (q \Rightarrow p \wedge q))$.

(22) $X \vdash_{IPC} (p \Rightarrow p \vee q)$.

(23) $X \vdash_{IPC} (q \Rightarrow p \vee q)$.

(24) $X \vdash_{IPC} (p \Rightarrow r \Rightarrow (q \Rightarrow r \Rightarrow (p \vee q \Rightarrow r)))$.

(25) $X \vdash_{IPC} (\mathrm{FALSUM} \Rightarrow p)$.

(26) If $X \vdash_{IPC} p$ and $X \vdash_{IPC} (p \Rightarrow q)$, then $X \vdash_{IPC} (q)$.

(27) $\vdash_{IPC} p \Rightarrow (q \Rightarrow p)$.

(28) $\vdash_{IPC} p \Rightarrow (q \Rightarrow r) \Rightarrow (p \Rightarrow q \Rightarrow (p \Rightarrow r))$.

(29) $\vdash_{IPC} p \wedge q \Rightarrow p$.

(30) $\vdash_{IPC} p \wedge q \Rightarrow q$.

(31) $\vdash_{IPC} p \Rightarrow (q \Rightarrow p \wedge q)$.

(32) $\vdash_{IPC} p \Rightarrow p \vee q$.

(33) $\vdash_{IPC} q \Rightarrow p \vee q$.

(34) $\vdash_{IPC} p \Rightarrow r \Rightarrow (q \Rightarrow r \Rightarrow (p \vee q \Rightarrow r))$.

(35) $\vdash_{IPC} \mathrm{FALSUM} \Rightarrow p$.

(36) If $\vdash_{IPC} p$ and $\vdash_{IPC} p \Rightarrow q$, then $\vdash_{IPC} q$.

Let us consider s. We say that s is IPC-valid if and only if

(Def. 8) $\emptyset_{\text{MC-w.f.f.}} \vdash_{IPC} (s)$.

One can verify that s is IPC-valid if and only if the condition (Def. 9) is satisfied.

(Def. 9) $s \in$ IPC-Taut.

Now we state the propositions:

(37) If p is IPC-valid, then $X \vdash_{IPC} (p)$.

(38) $p \Rightarrow (q \Rightarrow p)$ is IPC-valid.

(39) $p \Rightarrow (q \Rightarrow r) \Rightarrow (p \Rightarrow q \Rightarrow (p \Rightarrow r))$ is IPC-valid.

(40) $p \wedge q \Rightarrow p$ is IPC-valid.

(41) $p \wedge q \Rightarrow q$ is IPC-valid.

(42) $p \Rightarrow (q \Rightarrow p \wedge q)$ is IPC-valid.

(43) $p \Rightarrow p \vee q$ is IPC-valid.

(44) $q \Rightarrow p \vee q$ is IPC-valid.

(45) $p \Rightarrow r \Rightarrow (q \Rightarrow r \Rightarrow (p \vee q \Rightarrow r))$ is IPC-valid.

(46) FALSUM $\Rightarrow p$ is IPC-valid.

(47) If p is IPC-valid and $p \Rightarrow q$ is IPC-valid, then q is IPC-valid.

In the sequel X, T denote subsets of MC-w.f.f., F, G, H, p, q, r, t denote elements of MC-w.f.f., s, h denote MC-formulas, f denotes a finite sequence of elements of MC-w.f.f. $\times$ Proof-Step-Kinds-IPC, and i, j denote elements of $\mathbb{N}$.

4. The First Deduction Theorem for IPC

Now we state the propositions:

(48) $X \vdash_{IPC} (p \Rightarrow p)$. The theorem is a consequence of (26).

(49) $X \vdash_{IPC}$ (IVERUM).

(50) If $X \vdash_{IPC} (p)$, then $X \vdash_{IPC} (q \Rightarrow p)$.

(51) If p is IPC-valid, then $X \vdash_{IPC} (p)$.

(52) If $X \cup \{F\} \vdash_{IPC} (G)$, then $X \vdash_{IPC} (F \Rightarrow G)$.

PROOF: Consider f such that f is a proof w.r.t. IPC $(X \cup \{F\})$ and Effect-IPC$(f) = G$. Define $\mathcal{P}$[natural number] $\equiv$ if $1 \leqslant \$_1 \leqslant \operatorname{len} f$, then for every H such that $H = (f(\$_1))_{\mathbf{1}}$ holds $X \vdash_{IPC} (F \Rightarrow H)$. For every

natural number n such that for every natural number k such that $k < n$ holds $\mathcal{P}[k]$ holds $\mathcal{P}[n]$. For every natural number n, $\mathcal{P}[n]$. $1 \leqslant \operatorname{len} f$. $\square$

5. A Family of Deduction Theorems for IPC

From now on F_1, F_2, F_3, F_4, F_5, F_6, F_7, F_8, F_9, F_{10}, G denote MC-formulas and x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10}, x denote elements of MC-w.f.f..

Let x_1, x_2, x_3 be elements of MC-w.f.f.. Let us observe that the functor $\{x_1, x_2, x_3\}$ yields a subset of MC-w.f.f.. Let x_1, x_2, x_3, x_4 be elements of MC-w.f.f.. One can check that the functor $\{x_1, x_2, x_3, x_4\}$ yields a subset of MC-w.f.f.. Let x_1, x_2, x_3, x_4, x_5 be elements of MC-w.f.f.. One can verify that the functor $\{x_1, x_2, x_3, x_4, x_5\}$ yields a subset of MC-w.f.f.. Let x_1, x_2, x_3, x_4, x_5, x_6 be elements of MC-w.f.f.. One can verify that the functor $\{x_1, x_2, x_3, x_4, x_5, x_6\}$ yields a subset of MC-w.f.f.. Let x_1, x_2, x_3, x_4, x_5, x_6, x_7 be elements of MC-w.f.f..

One can check that the functor $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7\}$ yields a subset of MC-w.f.f.. Let x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8 be elements of MC-w.f.f.. Let us note that the functor $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\}$ yields a subset of MC-w.f.f.. Let x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9 be elements of MC-w.f.f.. One can verify that the functor $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9\}$ yields a subset of MC-w.f.f.. Let x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10} be elements of MC-w.f.f.. Observe that the functor $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10}\}$ yields a subset of MC-w.f.f.. Now we state the propositions:

(53) If $\{F\} \vdash_{IPC} (G)$, then $\vdash_{IPC} F \Rightarrow G$. The theorem is a consequence of (52).

(54) If $\{F_1, F_2\} \vdash_{IPC} (G)$, then $\{F_2\} \vdash_{IPC} (F_1 \Rightarrow G)$. The theorem is a consequence of (52).

(55) If $\{F_1, F_2, F_3\} \vdash_{IPC} (G)$, then $\{F_2, F_3\} \vdash_{IPC} (F_1 \Rightarrow G)$. The theorem is a consequence of (52).

(56) If $\{F_1, F_2, F_3, F_4\} \vdash_{IPC} (G)$, then $\{F_2, F_3, F_4\} \vdash_{IPC} (F_1 \Rightarrow G)$. The theorem is a consequence of (52).

(57) If $\{F_1, F_2, F_3, F_4, F_5\} \vdash_{IPC} (G)$, then $\{F_2, F_3, F_4, F_5\} \vdash_{IPC} (F_1 \Rightarrow G)$. The theorem is a consequence of (52).

(58) If $\{F_1, F_2, F_3, F_4, F_5, F_6\} \vdash_{IPC} (G)$, then $\{F_2, F_3, F_4, F_5, F_6\} \vdash_{IPC} (F_1 \Rightarrow G)$. The theorem is a consequence of (52).

(59) Suppose $\{F_1, F_2, F_3, F_4, F_5, F_6, F_7\} \vdash_{IPC} (G)$. Then $\{F_2, F_3, F_4, F_5, F_6, F_7\} \vdash_{IPC} (F_1 \Rightarrow G)$. The theorem is a consequence of (52).

(60) Suppose $\{F_1, F_2, F_3, F_4, F_5, F_6, F_7, F_8\} \vdash_{IPC} (G)$. Then $\{F_2, F_3, F_4, F_5, F_6, F_7, F_8\} \vdash_{IPC} (F_1 \Rightarrow G)$. The theorem is a consequence of (52).

(61) Suppose $\{F_1, F_2, F_3, F_4, F_5, F_6, F_7, F_8, F_9\} \vdash_{IPC} (G)$. Then $\{F_2, F_3, F_4, F_5, F_6, F_7, F_8, F_9\} \vdash_{IPC} (F_1 \Rightarrow G)$. The theorem is a consequence of (52).

From now on x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10} denote objects.

Now we state the propositions:

(62) $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10}\} = \{x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10}\} \cup \{x_1\}$.

(63) Suppose $\{F_1, F_2, F_3, F_4, F_5, F_6, F_7, F_8, F_9, F_{10}\} \vdash_{IPC} (G)$. Then $\{F_2, F_3, F_4, F_5, F_6, F_7, F_8, F_9, F_{10}\} \vdash_{IPC} (F_1 \Rightarrow G)$. The theorem is a consequence of (62) and (52).

6. Intuitionistic Provable Formulas and Theorems

Now we state the propositions:

(64) $\{p\} \vdash_{IPC} (p)$.

(65) If $X \vdash_{IPC} (p)$ and $X \subseteq Y$, then $Y \vdash_{IPC} (p)$. The theorem is a consequence of (15) and (9).

(66) If $p \in X$, then $X \vdash_{IPC} (p)$. The theorem is a consequence of (64) and (65).

(67) If $p \in X$, then $p \in \mathrm{CnIPC}(X)$. The theorem is a consequence of (66).

(68) If $p \in$ IPC-Taut, then $\vdash_{IPC} p$.

(69) If $\vdash_{IPC} p$, then $p \in$ IPC-Taut.

(70) $p \in$ IPC-Taut if and only if $\vdash_{IPC} p$.

(71) $\vdash_{IPC} p \Rightarrow (p \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM})$. The theorem is a consequence of (66), (26), (54), and (53).

(72) $\{p \wedge q\} \vdash_{IPC} (p)$. The theorem is a consequence of (19), (64), and (26).

(73) $\{p \wedge q\} \vdash_{IPC} (q)$. The theorem is a consequence of (20), (64), and (26).

(74) $\vdash_{IPC} (p \Rightarrow q) \wedge (p \Rightarrow (q \Rightarrow \mathrm{FALSUM})) \Rightarrow (p \Rightarrow \mathrm{FALSUM})$. The theorem is a consequence of (66), (19), (26), (20), (54), and (53).

(75) $\vdash_{IPC} p \Rightarrow \mathrm{FALSUM} \Rightarrow (p \Rightarrow q)$. The theorem is a consequence of (68).

(76) $\vdash_{IPC} (p \Rightarrow r) \wedge (q \Rightarrow r) \Rightarrow (p \vee q \Rightarrow r)$. The theorem is a consequence of (72), (73), (24), (26), and (53).

(77) $\vdash_{IPC} p \wedge (p \Rightarrow q) \Rightarrow q$. The theorem is a consequence of (72), (73), (26), and (53).

(78) $\vdash_{IPC} p \Rightarrow (p \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM})$. The theorem is a consequence of (69), (71), and (68).

(79) $\vdash_{IPC} (p \Rightarrow \mathrm{FALSUM}) \vee q \Rightarrow (p \Rightarrow q)$. The theorem is a consequence of (69), (75), (76), and (68).

(80) $\vdash_{IPC} p \Rightarrow q \Rightarrow (q \Rightarrow \mathrm{FALSUM} \Rightarrow (p \Rightarrow \mathrm{FALSUM}))$.

(81) $\vdash_{IPC} (p \Rightarrow \mathrm{FALSUM}) \vee (q \Rightarrow \mathrm{FALSUM}) \Rightarrow (p \wedge q \Rightarrow \mathrm{FALSUM})$. The theorem is a consequence of (69), (76), (80), and (68).

(82) Let us consider MC-formulas p, q. If $\vdash_{IPC} p$ and $\vdash_{IPC} q$, then $\vdash_{IPC} p \wedge q$. The theorem is a consequence of (31) and (36).

(83) If $\vdash_{IPC} p \Rightarrow q$ and $\vdash_{IPC} q \Rightarrow p$, then $\vdash_{IPC} p \Leftrightarrow q$.

(84) $\vdash_{IPC} p \Rightarrow p$. The theorem is a consequence of (27), (28), and (26).

(85) $\vdash_{IPC} p \Leftrightarrow p$. The theorem is a consequence of (84) and (82).

(86) $\vdash_{IPC} p \wedge q \Rightarrow \mathrm{FALSUM} \Rightarrow (p \Rightarrow (q \Rightarrow \mathrm{FALSUM}))$. The theorem is a consequence of (66), (21), (26), (55), (54), and (53).

(87) $\vdash_{IPC} p \Rightarrow (q \Rightarrow \mathrm{FALSUM}) \Rightarrow (p \wedge q \Rightarrow \mathrm{FALSUM})$. The theorem is a consequence of (66), (19), (26), (20), (54), and (53).

(88) $\vdash_{IPC} (p \wedge q \Rightarrow \mathrm{FALSUM}) \Leftrightarrow (p \Rightarrow (q \Rightarrow \mathrm{FALSUM}))$. The theorem is a consequence of (86), (87), and (83).

(89) $\vdash_{IPC} p \wedge q \Rightarrow \mathrm{FALSUM} \Rightarrow (q \Rightarrow (p \Rightarrow \mathrm{FALSUM}))$. The theorem is a consequence of (66), (21), (26), (55), (54), and (53).

(90) $\vdash_{IPC} q \Rightarrow (p \Rightarrow \mathrm{FALSUM}) \Rightarrow (p \wedge q \Rightarrow \mathrm{FALSUM})$. The theorem is a consequence of (66), (19), (26), (20), (54), and (53).

(91) $\vdash_{IPC} (q \Rightarrow (p \Rightarrow \mathrm{FALSUM})) \Leftrightarrow (p \wedge q \Rightarrow \mathrm{FALSUM})$. The theorem is a consequence of (89), (90), and (83).

(92) $\vdash_{IPC} p \Rightarrow (q \Rightarrow (p \wedge q \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM}))$. The theorem is a consequence of (66), (21), (65), (26), (55), (54), and (53).

(93) $\vdash_{IPC} q \Rightarrow (p \Rightarrow (p \wedge q \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM}))$. The theorem is a consequence of (66), (21), (65), (26), (55), (54), and (53).

(94) $\vdash_{IPC} p \Rightarrow (p \wedge q \Rightarrow \mathrm{FALSUM} \Rightarrow (q \Rightarrow \mathrm{FALSUM}))$. The theorem is a consequence of (66), (21), (65), (26), (55), (54), and (53).

(95) $\vdash_{IPC} q \Rightarrow (p \wedge q \Rightarrow \mathrm{FALSUM} \Rightarrow (p \Rightarrow \mathrm{FALSUM}))$. The theorem is a consequence of (66), (21), (65), (26), (55), (54), and (53).

(96) $\vdash_{IPC} p \vee q \Rightarrow \mathrm{FALSUM} \Rightarrow (p \Rightarrow \mathrm{FALSUM}) \wedge (q \Rightarrow \mathrm{FALSUM})$. The theorem is a consequence of (68).

(97) $\vdash_{IPC} (p \Rightarrow \mathrm{FALSUM}) \wedge (q \Rightarrow \mathrm{FALSUM}) \Rightarrow (p \vee q \Rightarrow \mathrm{FALSUM})$.

(98) $\vdash_{IPC} (p \vee q \Rightarrow \mathrm{FALSUM}) \Leftrightarrow (p \Rightarrow \mathrm{FALSUM}) \wedge (q \Rightarrow \mathrm{FALSUM})$. The theorem is a consequence of (96), (97), and (83).

(99) $\vdash_{IPC} p \wedge (p \Rightarrow \mathrm{FALSUM}) \Rightarrow \mathrm{FALSUM}$.

(100) $\vdash_{IPC} \mathrm{FALSUM} \Leftrightarrow p \wedge (p \Rightarrow \mathrm{FALSUM})$. The theorem is a consequence of (35), (99), and (83).

(101) $\vdash_{IPC} p \Rightarrow \mathrm{FALSUM} \Rightarrow (p \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM})$.

(102) $\vdash_{IPC} p \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM} \Rightarrow (p \Rightarrow \text{FALSUM})$. The theorem is a consequence of (69), (71), and (68).

(103) $\vdash_{IPC} (p \Rightarrow \text{FALSUM}) \Leftrightarrow (p \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM})$. The theorem is a consequence of (101), (102), and (83).

(104) $\vdash_{IPC} p \Rightarrow \text{FALSUM} \Rightarrow q \Rightarrow (p \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM} \Rightarrow q)$. The theorem is a consequence of (66), (102), (65), (26), (54), and (53).

(105) $\vdash_{IPC} p \Rightarrow q \Rightarrow (p \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM} \Rightarrow (q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM}))$. The theorem is a consequence of (69), (80), and (68).

(106) $\vdash_{IPC} p \wedge (q \Rightarrow \text{FALSUM}) \Rightarrow (p \Rightarrow q \Rightarrow \text{FALSUM})$. The theorem is a consequence of (66), (19), (26), (20), (54), and (53).

(107) $\vdash_{IPC} p \Rightarrow q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM} \Rightarrow (p \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM} \Rightarrow (q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM}))$. The theorem is a consequence of (66), (21), (26), (106), (80), (36), (65), (56), (55), (54), and (53).

(108) $\vdash_{IPC} p \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM} \Rightarrow (q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM}) \Rightarrow (p \Rightarrow q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM})$. The theorem is a consequence of (66), (79), (80), (36), (65), (26), (96), (19), (20), (54), and (53).

(109) $\vdash_{IPC} (p \Rightarrow q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM}) \Leftrightarrow (p \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM} \Rightarrow (q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM}))$. The theorem is a consequence of (107), (108), and (83).

(110) $\vdash_{IPC} p \wedge q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM} \Rightarrow (p \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM}) \wedge (q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM})$. The theorem is a consequence of (29), (30), (80), (36), and (68).

(111) $\vdash_{IPC} (p \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM}) \wedge (q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM}) \Rightarrow (p \wedge q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM})$. The theorem is a consequence of (66), (21), (26), (56), (19), (55), (20), (54), and (53).

(112) $\vdash_{IPC} (p \wedge q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM}) \Leftrightarrow (p \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM}) \wedge (q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM})$. The theorem is a consequence of (110), (111), and (83).

(113) $\vdash_{IPC} p \Rightarrow q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM} \Rightarrow (p \Rightarrow (q \Rightarrow \text{FALSUM} \Rightarrow \text{FALSUM}))$. The theorem is a consequence of (66), (107), (65), (26), (71), (54), and (53).

(114) If $\vdash_{IPC} r$ and $\{r\} \vdash_{IPC} (q)$, then $\vdash_{IPC} q$. The theorem is a consequence of (53) and (36).

(115) If $X \vdash_{IPC} (r)$ and $X \cup \{r\} \vdash_{IPC} (q)$, then $X \vdash_{IPC} (q)$. The theorem is a consequence of (52) and (26).

(116) If $X \vdash_{IPC} (r)$ and $Y \cup \{r\} \vdash_{IPC} (q)$, then $X \cup Y \vdash_{IPC} (q)$. The theorem is a consequence of (52), (65), and (26).

(117) If $\vdash_{IPC} p$ and $\{r\} \vdash_{IPC} (q)$, then $\{p \Rightarrow r\} \vdash_{IPC} (q)$. The theorem is a consequence of (65), (64), (26), and (115).

(118) If $X \vdash_{IPC} (p)$ and $X \cup \{r\} \vdash_{IPC} (q)$, then $X \cup \{p \Rightarrow r\} \vdash_{IPC} (q)$. The theorem is a consequence of (65), (66), (26), and (115).

(119) $\{q\} \vdash_{IPC} (q \vee r)$. The theorem is a consequence of (64), (22), and (26).

(120) $\{r\} \vdash_{IPC} (q \vee r)$. The theorem is a consequence of (64), (23), and (26).

(121) If $\{p\} \vdash_{IPC} (r)$ and $\{q\} \vdash_{IPC} (r)$, then $\{p \vee q\} \vdash_{IPC} (r)$. The theorem is a consequence of (34), (53), (36), (65), (26), and (64).

(122) If $X \cup \{p\} \vdash_{IPC} (r)$ and $X \cup \{q\} \vdash_{IPC} (r)$, then $X \cup \{p \vee q\} \vdash_{IPC} (r)$. The theorem is a consequence of (52), (24), (26), (64), and (65).

(123) If $X \cup \{p\} \vdash_{IPC} (r)$ and $Y \cup \{q\} \vdash_{IPC} (r)$, then $(X \cup Y) \cup \{p \vee q\} \vdash_{IPC} (r)$. The theorem is a consequence of (52), (65), (24), (26), and (64).

(124) $\vdash_{IPC} p \Rightarrow q \vee (p \Rightarrow r) \Rightarrow (p \Rightarrow q \vee r)$. The theorem is a consequence of (120), (65), (64), (118), (119), (122), (52), and (53).

(125) $\vdash_{IPC} p \Rightarrow (p \Rightarrow \mathrm{FALSUM} \Rightarrow q)$. The theorem is a consequence of (66), (26), (25), (54), and (53).

(126) $\vdash_{IPC} p \Rightarrow q \Rightarrow (q \wedge r \Rightarrow \mathrm{FALSUM} \Rightarrow (p \wedge r \Rightarrow \mathrm{FALSUM}))$. The theorem is a consequence of (66), (20), (26), (19), (21), (55), (54), and (53).

(127) $\vdash_{IPC} p \Rightarrow q \Rightarrow (q \vee r \Rightarrow \mathrm{FALSUM} \Rightarrow (p \vee r \Rightarrow \mathrm{FALSUM}))$. The theorem is a consequence of (66), (68), (65), (26), (55), (54), and (53).

Let p be an element of MC-w.f.f.. Note that the functor $\mathrm{neg}(p)$ yields an element of MC-w.f.f. and is defined by the term

(Def. 10) $p \Rightarrow \mathrm{FALSUM}$.

The functor $\mathrm{neg}^2(p)$ yielding an element of MC-w.f.f. is defined by the term

(Def. 11) $p \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM}$.

The functor $\mathrm{neg}^3(p)$ yielding an element of MC-w.f.f. is defined by the term

(Def. 12) $p \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM}$.

The functor $\mathrm{neg}^4(p)$ yielding an element of MC-w.f.f. is defined by the term

(Def. 13) $p \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM}$.

The functor $\mathrm{neg}^5(p)$ yielding an element of MC-w.f.f. is defined by the term

(Def. 14) $p \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM} \Rightarrow \mathrm{FALSUM}$.

Now we state the propositions:

(128) $\vdash_{IPC} p \Rightarrow \mathrm{neg}(\mathrm{neg}(p))$.

(129) $\vdash_{IPC} p \Rightarrow \mathrm{neg}^2(p)$.

(130) $\vdash_{IPC} (p \Rightarrow q) \wedge (p \Rightarrow \mathrm{neg}(q)) \Rightarrow \mathrm{neg}(p)$.

(131) $\vdash_{IPC} \mathrm{neg}(p) \Rightarrow (p \Rightarrow q)$.

(132) $\vdash_{IPC} p \Rightarrow \mathrm{neg}(\mathrm{neg}(\mathrm{neg}(\mathrm{neg}(p))))$.
(133) $\vdash_{IPC} \mathrm{neg}(p) \vee q \Rightarrow (p \Rightarrow q)$.
(134) $\vdash_{IPC} p \Rightarrow q \Rightarrow (\mathrm{neg}(q) \Rightarrow \mathrm{neg}(p))$.
(135) $\vdash_{IPC} \mathrm{neg}(p) \vee \mathrm{neg}(q) \Rightarrow \mathrm{neg}(p \wedge q)$.
(136) $\vdash_{IPC} \mathrm{neg}(p \wedge q) \Rightarrow (p \Rightarrow \mathrm{neg}(q))$.
(137) $\vdash_{IPC} p \Rightarrow \mathrm{neg}(q) \Rightarrow \mathrm{neg}(p \wedge q)$.
(138) $\vdash_{IPC} \mathrm{neg}(p \wedge q) \Leftrightarrow (p \Rightarrow \mathrm{neg}(q))$.
(139) $\vdash_{IPC} \mathrm{neg}(p \wedge q) \Rightarrow (q \Rightarrow \mathrm{neg}(p))$.
(140) $\vdash_{IPC} q \Rightarrow \mathrm{neg}(p) \Rightarrow \mathrm{neg}(p \wedge q)$.
(141) $\vdash_{IPC} (q \Rightarrow \mathrm{neg}(p)) \Leftrightarrow \mathrm{neg}(p \wedge q)$.
(142) $\vdash_{IPC} p \Rightarrow (q \Rightarrow \mathrm{neg}(\mathrm{neg}(p \wedge q)))$.
(143) $\vdash_{IPC} q \Rightarrow (p \Rightarrow \mathrm{neg}(\mathrm{neg}(p \wedge q)))$.
(144) $\vdash_{IPC} p \Rightarrow (\mathrm{neg}(p \wedge q) \Rightarrow \mathrm{neg}(q))$.
(145) $\vdash_{IPC} q \Rightarrow (\mathrm{neg}(p \wedge q) \Rightarrow \mathrm{neg}(p))$.
(146) $\vdash_{IPC} \mathrm{neg}(p \vee q) \Rightarrow \mathrm{neg}(p) \wedge \mathrm{neg}(q)$.
(147) $\vdash_{IPC} \mathrm{neg}(p) \wedge \mathrm{neg}(q) \Rightarrow \mathrm{neg}(p \vee q)$.
(148) $\vdash_{IPC} \mathrm{neg}(p \vee q) \Leftrightarrow \mathrm{neg}(p) \wedge \mathrm{neg}(q)$.
(149) $\vdash_{IPC} p \wedge \mathrm{neg}(p) \Rightarrow \mathrm{FALSUM}$.
(150) $\vdash_{IPC} \mathrm{FALSUM} \Leftrightarrow p \wedge \mathrm{neg}(p)$.
(151) $\vdash_{IPC} \mathrm{neg}(p) \Rightarrow \mathrm{neg}(\mathrm{neg}(\mathrm{neg}(p)))$.
(152) $\vdash_{IPC} \mathrm{neg}(\mathrm{neg}(\mathrm{neg}(p))) \Rightarrow \mathrm{neg}(p)$.
(153) $\vdash_{IPC} \mathrm{neg}(p) \Leftrightarrow \mathrm{neg}(\mathrm{neg}(\mathrm{neg}(p)))$.
(154) $\vdash_{IPC} \mathrm{neg}(p) \Rightarrow q \Rightarrow (\mathrm{neg}(\mathrm{neg}(\mathrm{neg}(p))) \Rightarrow q)$.
(155) $\vdash_{IPC} p \Rightarrow q \Rightarrow (\mathrm{neg}(\mathrm{neg}(p)) \Rightarrow \mathrm{neg}(\mathrm{neg}(q)))$.
(156) $\vdash_{IPC} p \wedge \mathrm{neg}(q) \Rightarrow \mathrm{neg}(p \Rightarrow q)$.
(157) $\vdash_{IPC} \mathrm{neg}(\mathrm{neg}(p \Rightarrow q)) \Rightarrow (\mathrm{neg}(\mathrm{neg}(p)) \Rightarrow \mathrm{neg}(\mathrm{neg}(q)))$.
(158) $\vdash_{IPC} \mathrm{neg}(\mathrm{neg}(p)) \Rightarrow \mathrm{neg}(\mathrm{neg}(q)) \Rightarrow \mathrm{neg}(\mathrm{neg}(p \Rightarrow q))$.
(159) $\vdash_{IPC} \mathrm{neg}(\mathrm{neg}(p \Rightarrow q)) \Leftrightarrow (\mathrm{neg}(\mathrm{neg}(p)) \Rightarrow \mathrm{neg}(\mathrm{neg}(q)))$.
(160) $\vdash_{IPC} \mathrm{neg}(\mathrm{neg}(p \wedge q)) \Rightarrow \mathrm{neg}(\mathrm{neg}(p)) \wedge \mathrm{neg}(\mathrm{neg}(q))$.
(161) $\vdash_{IPC} \mathrm{neg}(\mathrm{neg}(p)) \wedge \mathrm{neg}(\mathrm{neg}(q)) \Rightarrow \mathrm{neg}(\mathrm{neg}(p \wedge q))$.
(162) $\vdash_{IPC} \mathrm{neg}(\mathrm{neg}(p \wedge q)) \Leftrightarrow \mathrm{neg}(\mathrm{neg}(p)) \wedge \mathrm{neg}(\mathrm{neg}(q))$.
(163) $\vdash_{IPC} \mathrm{neg}(\mathrm{neg}(p \Rightarrow q)) \Rightarrow (p \Rightarrow \mathrm{neg}(\mathrm{neg}(q)))$.
(164) $\vdash_{IPC} p \Rightarrow (\mathrm{neg}(p) \Rightarrow q)$.
(165) $\vdash_{IPC} p \Rightarrow q \Rightarrow (\mathrm{neg}(q \wedge r) \Rightarrow \mathrm{neg}(p \wedge r))$.
(166) $\vdash_{IPC} p \Rightarrow q \Rightarrow (\mathrm{neg}(q \vee r) \Rightarrow \mathrm{neg}(p \vee r))$.

References

[1] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Matuszewski, Adam Naumowicz, Karol Pąk, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8_17.

[2] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Matuszewski, Adam Naumowicz, and Karol Pąk. The role of the Mizar Mathematical Library for interactive proof development in Mizar. *Journal of Automated Reasoning*, 61(1):9–32, 2018. doi:10.1007/s10817-017-9440-6.

[3] Agata Darmochwał. Calculus of quantifiers. Deduction theorem. *Formalized Mathematics*, 2(**2**):309–312, 1991.

[4] Arend Heyting. *Intuitionism. An introduction.* Elsevier, Amsterdam, 3rd revised ed., 1971.

[5] Takao Inoué. Intuitionistic propositional calculus in the extended framework with modal operator. Part I. *Formalized Mathematics*, 11(**3**):259–266, 2003.

[6] Anne Sjerp Troelstra and Dirk van Dalen. *Constructivism in mathematics. An introduction. Volume I*, volume 121 of *Studies in Logic and the Foundations of Mathematics*. Amsterdam etc.: North-Holland, 1988. ISBN 0-444-70506-6.

[7] Dirk van Dalen. *Logic and Structure.* London: Springer, 2013. ISBN 978-1-4471-4557-8; 978-1-4471-4558-5. doi:10.1007/978-1-4471-4558-5.

Accepted April 30, 2022