General Mathematics Vol. 29, No. 1 (2021), 103–119
DOI:10.2478/gm-2021-0009

sciendo

Horadam polynomials and their applications to a certain family of bi-univalent functions defined by Wanas operator [1]

Abbas Kareem Wanas, Timilehin Gideon Shaba

Abstract

In this article, by making use of Horadam polynomials, we introduce and investigate a certain family $\mathfrak{T}_{\Sigma}(\lambda, \alpha, \beta, k, \gamma; x)$ of analytic and bi-univalent functions associated with Wanas operator which defined in the open unit disk $\mathbb{U}$. We establish upper bounds for the initial Taylor-Maclaurin coefficients and obtain the Fekete-Szegö inequality of functions belonging to this family. We also point out several certain special cases for our results.

2010 Mathematics Subject Classification: Primary 30C45; Secondary 30C50.

Key words and phrases: Bi-univalent function, Horadam polynomials, Upper Bounds, Fekete-Szegö problem, Wanas operator.

[1] *Received 17 April, 2021*
Accepted for publication (in revised form) 13 June, 2021

1 Introduction

Let $\mathcal{A}$ be the collection of analytic functions in the open unit disk $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ that have the following normalized form:

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n. \tag{1}$$

Further, let S indicate the subclass of $\mathcal{A}$ consisting of functions which are univalent in $\mathbb{U}$. According to the Koebe one-quarter theorem [12] every function $f \in S$ has an inverse f^{-1} defined by $f^{-1}(f(z)) = z$, $(z \in \mathbb{U})$ and $f(f^{-1}(w)) = w$, $(|w| < r_0(f), r_0(f) \geqq \frac{1}{4})$, where

$$g(w) = f^{-1}(w) = w - a_2 w^2 + \left(2a_2^2 - a_3\right) w^3 - \left(5a_2^3 - 5a_2 a_3 + a_4\right) w^4 + \cdots . \tag{2}$$

A function $f \in \mathcal{A}$ is said to be bi-univalent in $\mathbb{U}$ if both f and f^{-1} are univalent in $\mathbb{U}$. Let Σ stands for the class of bi-univalent functions in $\mathbb{U}$ given by (1). In fact, Srivastava et al. [38] have actually revived the study of analytic and bi-univalent functions in recent years, it was followed by such works as those by Caglar et al. [10], Bulut [9], Adegani et al. [2] and others (see, for example [6, 7, 26, 27, 28, 29, 30, 33, 34, 35, 42, 45, 48, 50, 52]). From the work of Srivastava et al. [38], we choose to recall the following examples of functions in the class Σ :

$$\frac{z}{1-z}, \quad -\log(1-z) \qquad \text{and} \qquad \frac{1}{2}\log\left(\frac{1+z}{1-z}\right).$$

We notice that the class Σ is not empty. However, the Koebe function is not a member of Σ.

The problem to find the general coefficient bounds on the Taylor-Maclaurin coefficients $|a_n|$ $(n \in \mathbb{N};\ n \geqq 3)$ for functions $f \in \Sigma$ is still not completely addressed for many of the subclasses of the bi-univalent function class Σ (see, for example, [33, 39, 41]).

The Fekete-Szegö functional $\left|a_3 - \mu a_2^2\right|$ for $f \in S$ is well known for its rich history in the field of Geometric Function Theory. Its origin was in the disproof by Fekete and Szegö [13] of the Littlewood-Paley conjecture that the coefficients of odd univalent functions are bounded by unity. The functional

has since received great attention, particularly in the study of many subclasses of the family of univalent functions. This topic has become of considerable interest among researchers in Geometric Function Theory (see, for example, [21, 24, 36, 37, 40]).

With a view to recalling the principle of subordination between analytic functions, let the functions f and g be analytic in $\mathbb{U}$. We say that the function f is subordinate to g, if there exists a Schwarz function ω, which is analytic in $\mathbb{U}$ with $\omega(0) = 0 \,\text{and}\, |\omega(z)| < 1\,(z \in \mathbb{U})$, such that $f(z) = g\big(\omega(z)\big)$. This subordination is denoted by $f \prec g \,\text{or}\, f(z) \prec g(z) \quad (z \in \mathbb{U})$. It is well known that (see [23]), if the function g is univalent in $\mathbb{U}$, then

$$f \prec g \quad (z \in \mathbb{U}) \iff f(0) = g(0) \quad \text{and} \quad f(\mathbb{U}) \subseteq g(\mathbb{U}).$$

Recently, Horzum and Koçer [17] considered the Horadam polynomials $h_n(x)$, which are given by the following recurrence relation (see also Horadam and Mahon [16]):

$$h_n(x) = pxh_{n-1}(x) + qh_{n-2}(x) \quad (x \in \mathbb{R};\ n \in \mathbb{N} - \{1, 2\}), \tag{3}$$

with $h_1(x) = a \quad \text{and} \quad h_2(x) = bx$, for some real constant a, b, p and q. The characteristic equation of repetition relation (3) is $t^2 - pxt - q = 0$. This equation has two real roots $x_1 = \frac{px+\sqrt{p^2x^2+4q}}{2}$ and $x_2 = \frac{px-\sqrt{p^2x^2+4q}}{2}$.

Remark 1 *By selecting the particular values of a, b, p and q, the Horadam polynomial $h_n(x)$ reduces to several polynomials. Some of these special cases are recorded below.*

1. Taking $a = b = p = q = 1$, we obtain the Fibonacci polynomials $F_n(x)$.

2. Taking $a = 2$ and $b = p = q = 1$, we attain the Lucas polynomials $L_n(x)$.

3. Taking $a = q = 1$ and $b = p = 2$, we have the Pell polynomials $P_n(x)$.

4. Taking $a = b = p = 2$ and $q = 1$, we get the Pell-Lucas polynomials $Q_n(x)$.

5. Taking $a = b = 1$, $p = 2$ and $q = -1$, we obtain the Chebyshev polynomials $T_n(x)$ of the first kind.

6. Taking $a = 1$, $b = p = 2$ and $q = -1$, we have the Chebyshev polynomials $U_n(x)$ of the second kind.

These polynomials, the families of orthogonal polynomials and other special polynomials as well as their generalizations are potentially important in a variety of disciplines in many of sciences, specially in the mathematics, statistics and physics. For more information associated with these polynomials see [14, 15, 16, 19, 20].

The generating function of the Horadam polynomials $h_n(x)$ (see [17]) is given by

$$\Pi(x,z) = \sum_{n=1}^{\infty} h_n(x) z^{n-1} = \frac{a + (b - ap)xz}{1 - pxz - qz^2}. \tag{4}$$

In fact, Srivastava et al. [31] have already used Horadam polynomials in a similar context involving analytic and bi-univalent functions in recent years, it was followed by such works as those by Magesh et al. [22], Al-Amoush [3], Wanas and Alina [49], Abirami et al. [1] and Srivastava et al. [43].

Recently, Wanas [47] introduced the following operator (so-called Wanas operator) $\mathcal{W}_{\alpha,\beta}^{k,\gamma} : \mathcal{A} \to \mathcal{A}$ defined by

$$\mathcal{W}_{\alpha,\beta}^{k,\gamma} f(z) = z + \sum_{n=2}^{\infty} \left[\Phi_n(k,\alpha,\beta)\right]^{\gamma} a_n z^n, \tag{5}$$

where

$$\Phi_n(k,\alpha,\beta) = \sum_{m=1}^{k} \binom{k}{m} (-1)^{m+1} \left(\frac{\alpha^m + n\beta^m}{\alpha^m + \beta^m}\right) \tag{6}$$

$$\left(\alpha \in \mathbb{R};\ \beta \geqq 0 \ with\ \alpha + \beta > 0;\ m, \gamma \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}\right).$$

Special cases of this operator can be found in [4, 5, 8, 11, 18, 25, 32, 44, 46]. For more details see [51].

2 Main Results

We begin this section by defining the family $\mathfrak{T}_{\Sigma}(\lambda, \alpha, \beta, k, \gamma; x)$ as follows:

Definition 1 *For $0 \leqq \lambda \leqq 1$ and $x \in \mathbb{R}$, a function $f \in \Sigma$ is said to be in the family $\mathfrak{T}_{\Sigma}(\lambda, \alpha, \beta, k, \gamma; x)$ if it fulfills the subordinations:*

$$1 + \frac{z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma} f(z)\right)'}{\mathcal{W}_{\alpha,\beta}^{k,\gamma} f(z)} + \frac{z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma} f(z)\right)''}{\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma} f(z)\right)'} - \frac{\lambda z^2\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma} f(z)\right)'' + z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma} f(z)\right)'}{\lambda z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma} f(z)\right)' + (1-\lambda)\mathcal{W}_{\alpha,\beta}^{k,\gamma} f(z)}$$
$$\prec \Pi(x, z) + 1 - a$$

and

$$1 + \frac{w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma} g(w)\right)'}{\mathcal{W}_{\alpha,\beta}^{k,\gamma} g(w)} + \frac{w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma} g(w)\right)''}{\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma} g(w)\right)'} - \frac{\lambda w^2\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma} g(w)\right)'' + w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma} g(w)\right)'}{\lambda w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma} g(w)\right)' + (1-\lambda)\mathcal{W}_{\alpha,\beta}^{k,\gamma} g(w)}$$
$$\prec \Pi(x, w) + 1 - a,$$

where a is real constant and the function $g = f^{-1}$ is given by (2).

Remark 2 *For $\lambda = \gamma = 0$, a function $f \in \Sigma$ is in the family*

$$\mathfrak{T}_{\Sigma}(0, \alpha, \beta, k, 0; x) =: \mathcal{K}_{\Sigma}(x)$$

which was considered recently by Magesh et al. in [22], if the following conditions are satisfied:

$$1 + \frac{zf''(z)}{f'(z)} \prec \Pi(x, z) + 1 - a$$

and

$$1 + \frac{wg''(w)}{g'(w)} \prec \Pi(x, w) + 1 - a,$$

where $z, w \in \mathbb{U}$ and the function g is described in (2).

Remark 3 *For $\lambda = 1$ and $\gamma = 0$, a function $f \in \Sigma$ is in the family*

$$\mathfrak{T}_{\Sigma}(1, \alpha, \beta, k, 0; x) =: \mathcal{W}_{\Sigma}(x)$$

which was considered recently by Srivastava et al. in [31], the following conditions are satisfied:

$$\frac{zf'(z)}{f(z)} \prec \Pi(x, z) + 1 - a \quad \text{and} \quad \frac{wg'(w)}{g(w)} \prec \Pi(x, w) + 1 - a,$$

where $z, w \in \mathbb{U}$ and the function g is described in (2).

Theorem 1 *For* $0 \leqq \lambda \leqq 1$ *and* $x \in \mathbb{R}$, *let* $f \in \mathcal{A}$ *be in the family* $\mathfrak{T}_{\Sigma}(\lambda, \alpha, \beta, k, \gamma; x)$. *Then*

$$|a_2| \leqq \frac{|bx|\sqrt{|bx|}}{\sqrt{\left|\left[\vartheta(\lambda,\gamma,k,\alpha,\beta)b - p\,(2-\lambda)^2\,\Phi_2^{2\gamma}(k,\alpha,\beta)\right]bx^2 - qa\,(2-\lambda)^2\,\Phi_2^{2\gamma}(k,\alpha,\beta)\right|}}$$

and

$$|a_3| \leqq \frac{|bx|}{(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)} + \frac{b^2x^2}{(2-\lambda)^2\,\Phi_2^{2\gamma}(k,\alpha,\beta)},$$

where

$$\vartheta(\lambda,\gamma,k,\alpha,\beta) = 2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta) - \left(5-(\lambda+1)^2\right)\Phi_2^{2\gamma}(k,\alpha,\beta) \tag{7}$$

Proof. Let $f \in \mathfrak{T}_{\Sigma}(\lambda, \alpha, \beta, k, \gamma; x)$. Then there are two analytic functions $u, v : \mathbb{U} \longrightarrow \mathbb{U}$ given by

$$u(z) = u_1 z + u_2 z^2 + u_3 z^3 + \cdots \quad (z \in \mathbb{U}) \tag{8}$$

and

$$v(w) = v_1 w + v_2 w^2 + v_3 w^3 + \cdots \quad (w \in \mathbb{U}), \tag{9}$$

with $u(0) = v(0) = 0$ and $\max\{|u(z)|, |v(w)|\} < 1$, $z, w \in \mathbb{U}$ such that

$$1 + \frac{z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)'}{\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)} + \frac{z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)''}{\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)'} - \frac{\lambda z^2\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)'' + z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)'}{\lambda z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)' + (1-\lambda)\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)} = \Pi(x,u(z)) + 1 - a$$

and

$$1 + \frac{w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)'}{\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)} + \frac{w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)''}{\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)'} - \frac{\lambda w^2\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)'' + w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)'}{\lambda w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)' + (1-\lambda)\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)} = \Pi(x,v(w)) + 1 - a.$$

Or, equivalently

$$1 + \frac{z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)'}{\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)} + \frac{z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)''}{\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)'} - \frac{\lambda z^2\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)'' + z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)'}{\lambda z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)' + (1-\lambda)\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)} = 1 + h_1(x) + h_2(x)u(z) + h_3(x)u^2(z) + \cdots \tag{10}$$

and

(11)
$$1+\frac{w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)'}{\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)}+\frac{w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)''}{\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)'}-\frac{\lambda w^2\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)''+w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)'}{\lambda w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)'+(1-\lambda)\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)}$$
$$=1+h_1(x)+h_2(x)v(w)+h_3(x)v^2(w)+\cdots.$$

Combining (8), (9), (10) and (11) yields

$$1+\frac{z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)'}{\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)}+\frac{z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)''}{\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)'}-\frac{\lambda z^2\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)''+z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)'}{\lambda z\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)\right)'+(1-\lambda)\mathcal{W}_{\alpha,\beta}^{k,\gamma}f(z)}$$
$$=1+h_2(x)u_1z+\left[h_2(x)u_2+h_3(x)u_1^2\right]z^2+\cdots \tag{12}$$

and

(13)
$$1+\frac{w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)'}{\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)}+\frac{w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)''}{\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)'}-\frac{\lambda w^2\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)''+w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)'}{\lambda w\left(\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)\right)'+(1-\lambda)\mathcal{W}_{\alpha,\beta}^{k,\gamma}g(w)}$$
$$=1+h_2(x)v_1w+\left[h_2(x)v_2+h_3(x)v_1^2\right]w^2+\cdots.$$

It is quite well-known that if $\max\{|u(z)|,|v(w)|\}<1$, $z,w\in\mathbb{U}$, then

$$|u_i|\leqq 1 \qquad \text{and} \qquad |v_i|\leqq 1 \qquad (\forall\, i\in\mathbb{N}). \tag{14}$$

Comparing the corresponding coefficients in (12) and (13), after simplifying, we have

$$(2-\lambda)\Phi_2^{\gamma}(k,\alpha,\beta)a_2=h_2(x)u_1, \tag{15}$$

$$2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)a_3-\left(5-(\lambda+1)^2\right)\Phi_2^{2\gamma}(k,\alpha,\beta)a_2^2$$
$$=h_2(x)u_2+h_3(x)u_1^2, \tag{16}$$

$$-(2-\lambda)\Phi_2^{\gamma}(k,\alpha,\beta)a_2=h_2(x)v_1 \tag{17}$$

and

$$2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)(2a_2^2-a_3)-\left(5-(\lambda+1)^2\right)\Phi_2^{2\gamma}(k,\alpha,\beta)a_2^2 = h_2(x)v_2+h_3(x)v_1^2. \tag{18}$$

It follows from (15) and (17) that

$$u_1=-v_1 \tag{19}$$

and

$$2\left(2-\lambda\right)^2\Phi_2^{2\gamma}(k,\alpha,\beta)a_2^2=h_2^2(x)(u_1^2+v_1^2). \tag{20}$$

If we add (16) to (18), we find that

$$2\vartheta(\lambda,\gamma,k,\alpha,\beta)a_2^2=h_2(x)(u_2+v_2)+h_3(x)(u_1^2+v_1^2), \tag{21}$$

where $\vartheta(\lambda,\gamma,k,\alpha,\beta)$ is given by (7).
Substituting the value of $u_1^2+v_1^2$ from (20) in the right hand side of (21), we deduce that

$$a_2^2=\frac{h_2^3(x)(u_2+v_2)}{2\left[h_2^2(x)\vartheta(\lambda,\gamma,k,\alpha,\beta)-h_3(x)\left(2-\lambda\right)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right]}. \tag{22}$$

Further computations using (3), (14) and (22), we obtain

$$|a_2|\leqq\frac{|bx|\sqrt{|bx|}}{\sqrt{\left|\left[\vartheta(\lambda,\gamma,k,\alpha,\beta)b-p\left(2-\lambda\right)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right]bx^2-qa\left(2-\lambda\right)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right|}}.$$

Next, if we subtract (18) from (16), we can easily see that

$$4(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)(a_3-a_2^2)=h_2(x)(u_2-v_2)+h_3(x)(u_1^2-v_1^2). \tag{23}$$

In view of (19) and (20), we get from (23)

$$a_3=\frac{h_2(x)(u_2-v_2)}{4(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}+\frac{h_2^2(x)(u_1^2+v_1^2)}{2\left(2-\lambda\right)^2\Phi_2^{2\gamma}(k,\alpha,\beta)}.$$

Thus applying (3), we obtain

$$|a_3|\leqq\frac{|bx|}{(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}+\frac{b^2x^2}{\left(2-\lambda\right)^2\Phi_2^{2\gamma}(k,\alpha,\beta)}.$$

This completes the proof of Theorem 1.

By taking $\lambda=\gamma=0$ in Theorem 1, we state

Corollary 1 *[22] Let f given by 1 be in the family $\mathcal{K}_{\Sigma}(x)$. Then*

$$\begin{aligned} |a_2| &\leqq \frac{|bx|\sqrt{|bx|}}{\sqrt{|(2b-4p)bx^2-4qa|}}, \\ |a_3| &\leqq \frac{|bx|}{6}+\frac{b^2x^2}{4}. \end{aligned}$$

By taking $\lambda = 1$ and $\gamma = 0$ in Theorem 1, we state

Corollary 2 *[31] Let f given by 1 be in the family $\mathcal{W}_{\Sigma}(x)$. Then*

$$\begin{aligned} |a_2| &\leqq \frac{|bx|\sqrt{|bx|}}{\sqrt{|(b-p)bx^2-qa|}}, \\ |a_3| &\leqq \frac{|bx|}{2}+b^2x^2. \end{aligned}$$

In the next theorem, we present the Fekete-Szegö inequality for the family $\mathfrak{T}_{\Sigma}(\lambda,\alpha,\beta,k,\gamma;x)$.

Theorem 2 *For $0 \leqq \lambda \leqq 1$ and $x,\mu \in \mathbb{R}$, let $f \in \mathcal{A}$ be in the family $\mathfrak{T}_{\Sigma}(\lambda,\alpha,\beta,k,\gamma;x)$. Then*

$$\left|a_3-\mu a_2^2\right| \leqq \begin{cases} \frac{|bx|}{2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}; \\ for\ \ |\mu-1| \leqq \frac{\left|\left[\vartheta(\lambda,\gamma,k,\alpha,\beta)b-p(2-\lambda)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right]bx^2-qa(2-\lambda)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right|}{2b^2x^2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}, \\ \\ \frac{|bx|^3|\mu-1|}{\left|\left[\vartheta(\lambda,\gamma,k,\alpha,\beta)b-p(2-\lambda)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right]bx^2-qa(2-\lambda)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right|}; \\ \\ for\ \ |\mu-1| \geqq \frac{\left|\left[\vartheta(\lambda,\gamma,k,\alpha,\beta)b-p(2-\lambda)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right]bx^2-qa(2-\lambda)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right|}{2b^2x^2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}. \end{cases}$$

Proof. It follows from (22) and (23) that

$$\begin{aligned} a_3-\mu a_2^2 &= \frac{h_2(x)(u_2-v_2)}{4(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}+(1-\mu)\,a_2^2 \\ &= \frac{h_2(x)(u_2-v_2)}{4(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)} \\ &+ \frac{h_2^3(x)(u_2+v_2)\,(1-\mu)}{2\left[h_2^2(x)\vartheta(\lambda,\gamma,k,\alpha,\beta)-h_3(x)\,(2-\lambda)^2\,\Phi_2^{2\gamma}(k,\alpha,\beta)\right]} \\ &= \frac{h_2(x)}{2}\left[\left(\psi(\mu,x)+\frac{1}{2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}\right)u_2\right. \\ &+ \left.\left(\psi(\mu,x)-\frac{1}{2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}\right)v_2\right], \end{aligned}$$

where

$$\psi(\mu,x)=\frac{h_2^2(x)\,(1-\mu)}{h_2^2(x)\vartheta(\lambda,\gamma,k,\alpha,\beta)-h_3(x)\,(2-\lambda)^2\,\Phi_2^{2\gamma}(k,\alpha,\beta)}.$$

According to (3), we find that

$$\left|a_3-\mu a_2^2\right| \leqq \begin{cases} \frac{|bx|}{2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}, & 0 \leqq |\psi(\mu,x)| \leqq \frac{1}{2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}, \\ |bx|\,|\psi(\mu,x)|\,, & |\psi(\mu,x)| \geq \frac{1}{2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}. \end{cases}$$

After some computations, we obtain

$$\left|a_3-\mu a_2^2\right| \leqq \begin{cases} \frac{|bx|}{2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}; \\ for\ \ |\mu-1| \leqq \frac{\left|\left[\vartheta(\lambda,\gamma,k,\alpha,\beta)b-p(2-\lambda)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right]bx^2-qa(2-\lambda)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right|}{2b^2x^2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}, \\ \frac{|bx|^3|\mu-1|}{\left|\left[\vartheta(\lambda,\gamma,k,\alpha,\beta)b-p(2-\lambda)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right]bx^2-qa(2-\lambda)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right|}; \\ for\ \ |\mu-1| \geqq \frac{\left|\left[\vartheta(\lambda,\gamma,k,\alpha,\beta)b-p(2-\lambda)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right]bx^2-qa(2-\lambda)^2\Phi_2^{2\gamma}(k,\alpha,\beta)\right|}{2b^2x^2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}. \end{cases}$$

Putting $\mu=1$ in Theorem 2, we obtain the following result:

Corollary 3 *For* $0 \leqq \lambda \leqq 1$ *and* $x \in \mathbb{R}$, *let* $f \in \mathcal{A}$ *be in the family* $\mathfrak{T}_{\Sigma}(\lambda,\alpha,\beta,k,\gamma;x)$. *Then*

$$\left|a_3-a_2^2\right| \leqq \frac{|bx|}{2(3-2\lambda)\Phi_3^{\gamma}(k,\alpha,\beta)}.$$

Corollary 4 *[22] Let f given by 1 be in the family $\mathcal{K}_{\Sigma}(x)$. Then*

$$\left|a_3-\mu a_2^2\right| \leqq \begin{cases} \frac{|bx|}{6} & ;\ |\mu-1| \leqq \frac{\left|(2b-4p)bx^2-4qa\right|}{6b^2x^2}, \\ \\ \frac{|bx|^3|\mu-1|}{\left|(2b-4p)bx^2-4qa\right|} & ;\ |\mu-1| \geq \frac{\left|(2b-4p)bx^2-4qa\right|}{6b^2x^2}. \end{cases}$$

Corollary 5 *[31] Let f given by 1 be in the family $\mathcal{W}_{\Sigma}(x)$. Then*

$$\left|a_3-\mu a_2^2\right| \leqq \begin{cases} \frac{|bx|}{2} & ;\ |\mu-1| \leqq \frac{\left|(b-p)bx^2-qa\right|}{b^2x^2}, \\ \\ \frac{|bx|^3|\mu-1|}{\left|(b-p)bx^2-qa\right|} & ;\ |\mu-1| \geq \frac{\left|(b-p)bx^2-qa\right|}{b^2x^2}. \end{cases}$$

3 Conclusion

The fact that we can find many unique and effective uses of a large variety of interesting functions and specific polynomial in Geometric Function Theory provided the primary inspiration for our analysis in this article. The primary objective was to create a new subclass $\mathfrak{T}_{\Sigma}(\lambda,\alpha,\beta,k,\gamma;x)$ of the bi-univalent function class Σ defined by Wanas operator and also using the Horadam polynomial $h_n(x)$, which are given by the recurrence relation (3) and generating function $\Pi(x,z)$ in (4).

We generate Taylor-Maclaurin coefficient inequalities for functions belonging to this newly introduced bi-univalent function class $\mathfrak{T}_{\Sigma}(\lambda,\alpha,\beta,k,\gamma;x)$ and viewed the famous Fekete-Szegö problem. We've also made important correlations between our findings and those considered in previous studies.

References

[1] C. Abirami, N. Magesh, J. Yamini, *Initial bounds for certain classes of bi-univalent functions defined by Horadam polynomials*, Abstr. Appl. Anal., Art. ID 7391058, 2020, 1-8.

[2] E. A. Adegani, S. Bulut, A. A. Zireh, *Coefficient estimates for a subclass of analytic bi-univalent functions*, Bull. Korean Math. Soc., vol. 55, no. 2, 2018, 405-413.

[3] A. G. Al-Amoush, *Certain subclasses of bi-univalent functions involving the Poisson distribution associated with Horadam polynomials*, Malaya J. Mat., vol. 7, 2019, 618-624.

[4] J. W. Alexander, *Functions which map the interior of the unit circle upon simple region*, Annals Math., vol. 17, 1915, 12-22.

[5] F. M. Al-Oboudi, *On univalent functions defined by a generalized Sălăgean operator*, Int. J. Math. Math. Sci., vol. 27, 2004, 1429-1436.

[6] Ş. Altınkaya, *Bounds for a new subclass of bi-univalent functions subordinate to the Fibonacci numbers*, Turk. J. Math., vol. 44, no. 2, 2020, 553-560.

[7] Ş. Altınkaya, S. Yalçin, *Faber polynomial coefficient bounds for a subclass of bi-univalent functions*, Comptes Rendus Mathematique, vol. 353, no. 12, 2015, 1075-1080.

[8] S. D. Bernardi, *Convex and starlike univalent functions*, Transactions of the Amer. Math. Society, vol. 135, 1969, 429-446.

[9] S. Bulut, *Coefficient estimates for general subclasses of m-fold symmetric analytic bi-univalent functions*, Turk. J. Math., vol. 40, 2016, 1386-1397.

[10] M. Caglar, H. Orhan, N. Yagmur, *Coefficient bounds for new subclasses of bi-univalent functions*, Filomat, vol. 27, 2013, 1165-1171.

[11] N. E. Cho, H. M. Srivastava, *Argument estimates of certain analytic functions defined by a class of multiplier transformations*, Math. Comput. Model., vol. 37, 2003, 39-49.

[12] P. L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften, Band 259, Springer Verlag, New York, Berlin, Heidelberg and Tokyo, 1983.

[13] M. Fekete, G. Szegö, *Eine bemerkung uber ungerade schlichte funktionen*, J. London Math. Soc., vol. 2, 1933, 85-89.

[14] H. O. Güney, G. Murugusundaramoorthy, J. Sokół, *Subclasses of bi-univalent functions related to shell-like curves connected with Fibonacci numbers*, Acta Univ. Sapientiae, Mathematica, vol. 10, no. 1, 2018, 70-84.

[15] A. F. Horadam, *Jacobsthal Representation Polynomials*, The Fibonacci Quarterly, vol. 35, no. 2, 1997, 137-148.

[16] A. F. Horadam, J. M. Mahon, *Pell and Pell-Lucas polynomials*, The Fibonacci Quarterly, **23**(1) (1985), 7-20.

[17] T. Horzum, E. G. Kocer, *On some properties of Horadam polynomials*, Int. Math. Forum, vol. 4, 2009, 1243-1252.

[18] I. B. Jung, Y. C. Kim, H. M. Srivastava, *The Hardy space of analytic functions associated with certain one-parameter families of integral operators*, J. Math. Anal. Appl., vol. 176, 1993, 138-147.

[19] T. Koshy, *Fibonacci and Lucas Numbers with Applications*, A Wiley- Interscience Publication, New York, 2001.

[20] A. Lupas, *A Guide of Fibonacci and Lucas Polynomials*, Octagon Mathematics Magazine, vol. 7, no. 1, 1999, 2-12.

[21] N. Magesh, J. Yamini, *Fekete-Szegö problem and second Hankel determinant for a class of bi-univalent functions*, Tbilisi Math. J., vol. 11, no. 1, 2018, 141-157.

[22] N. Magesh, J. Yamini, C. Abirami, *Initial bounds for certain classes of bi-univalent functions defined by Horadam Polynomials*, arXiv:1812.04464v1, 2018, 1-14.

[23] S. S. Miller, P. T. Mocanu, *Differential Subordinations: Theory and Applications*, Series on Monographs and Textbooks in Pure and Applied Mathematics, vol. 225, Marcel Dekker Inc., New York and Basel, 2000.

[24] R. K. Raina, J. Sokół, *Fekete-Szegö problem for some starlike functions related to shell-like curves*, Math. Slovaca, vol. 66, 2016, 135-140.

[25] G. St. Sălăgean, *Subclasses of univalent functions*, Lecture Notes in Mathematics, Springer Verlag, Berlin, vol. 1013, 1983, 362-372.

[26] T. G. Shaba, *On some new subclass of bi-univalent functions associated with the Opoola differential operator*, Open J. Math. Anal., vol. 4, no. 2, 2020, 74-79.

[27] T. G. Shaba, *Subclass of bi-univalent functions satisfying subordinate conditions defined by Frasin differential operator*, Turkish Journal of Inequalities, vol.4, no. 2, 2020, 50-58.

[28] T. G. Shaba, *On some subclasses of bi-pseudo-starlike functions defined by Salagean differential operator*, Asia Pac. J. Math., vol. 8, no. 6, 2021, 1-11.

[29] T. G. Shaba, A. B. Patil, *Coefficient estimates for certain subclasses of m-fold symmetric bi-univalent functions associated with pseudo-starlike functions*, Earthline Journal of Mathematical Sciences, vol. 6, no. 2, 2021, 2581-8147.

[30] H. M. Srivastava, *Operators of basic (or q-) calculus and fractional q-calculus and their applications in geometric function theory of complex analysis*, Iran. J. Sci. Technol. Trans. A: Sci., vol. 44, 2020, 327-344.

[31] H. M. Srivastava, Ş. Altınkaya, S. Yalçin, *Certain subclasses of bi-univalent functions associated with the Horadam polynomials*, Iran. J. Sci. Technol. Trans. A Sci., vol. 43, 2019, 1873-1879.

[32] H. M. Srivastava, A. A. Attiya, *An integral operator associated with the Hurwitz-Lerch Zeta function and differential subordination*, Integral Transforms and Special Functions, vol. 18, 2007, 207-216.

[33] H. M. Srivastava, S. S. Eker, S. G. Hamidi, J. M. Jahangiri, *Faber polynomial coefficient estimates for bi-univalent functions defined by the Tremblay fractional derivative operator*, Bull. Iranian Math. Soc., vol. 44, no. 1, 2018, 149-157.

[34] H. M. Srivastava, S. Gaboury, F. Ghanim, *Coefficient estimates for some general subclasses of analytic and bi-univalent functions*, Africa Mat., vol. 28, 2017, 693-706.

[35] H. M. Srivastava, S. Gaboury, F. Ghanim, *Coefficient estimates for a general subclass of analytic and bi-univalent functions of the Ma-Minda type*, Rev. Real Acad. Cienc. Exactas Fís. Natur. Ser. A Mat. (RACSAM), vol. 112, 2018, 1157-1168.

[36] H. M. Srivastava, S. Hussain, A. Raziq, M. Raza, *The Fekete-Szegö functional for a subclass of analytic functions associated with quasi-subordination*, Carpathian J. Math., vol. 34, 2018, 103-113.

[37] H. M. Srivastava, A. K. Mishra, M. K. Das, *The Fekete-Szegö problem for a subclass of close-to-convex functions*, Complex Variables Theory Appl., vol. 44, 2001, 145-163.

[38] H. M. Srivastava, A. K. Mishra, P. Gochhayat, *Certain subclasses of analytic and bi-univalent functions*, Appl. Math. Lett., vol. 23, 2010, 1188-1192.

[39] H. M. Srivastava, A. Motamednezhad, E. A. Adegani, *Faber polynomial coefficient estimates for bi-univalent functions defined by using differential subordination and a certain fractional derivative operator*, Mathematics, Art. ID 172, vol. 8, 2020, 1-12.

[40] H. M. Srivastava, N. Raza, E. S. A. AbuJarad, G. Srivastava, M. H. AbuJarad, *Fekete-Szegö inequality for classes of* (p, q)*-starlike and* (p, q)*-convex functions*, Rev. Real Acad. Cienc. Exactas Fís. Natur. Ser. A Mat. (RACSAM), vol. 113, 2019, 3563-3584.

[41] H. M. Srivastava, F. M. Sakar, H. Ö. Güney, *Some general coefficient estimates for a new class of analytic and bi-univalent functions defined by a linear combination*, Filomat, vol. 32, 2018, 1313-1322.

[42] H. M. Srivastava, A. K. Wanas, *Initial Maclaurin coefficient bounds for new subclasses of analytic and m-fold symmetric bi-univalent functions*

defined by a linear combination, Kyungpook Math. J., vol. 59, no. 3, 2019, 493-503.

[43] H. M. Srivastava, A. K. Wanas, G. Murugusundaramoorthy, *Certain family of bi-univalent functions associated with Pascal distribution series based on Horadam polynomials*, Surveys in Mathematics and its Applications, vol. 16, 2021, 193-205.

[44] S. R. Swamy, *Inclusion properties of certain subclasses of analytic functions*, Int. Math. Forum, vol. 7, 2012, 1751-1760.

[45] S. R. Swamy, A. K. Wanas, Y. Sailaja, *Some special families of holomorphic and Sălăgean type bi-univalent functions associated with (m,n)-Lucas polynomials*, Communications in Mathematics and Applications, vol. 11, no. 4, 2020, 563-574.

[46] B. A. Uralegaddi, C. Somanatha, *Certain classes of univalent functions, in Current Topics in Analytic Function Theory*, (Edited by H. M. Srivastava and S. Own), 371-374, World Scientific, Singapore, 1992.

[47] A. K. Wanas, *New differential operator for holomorphic functions*, Earthline J. Math. Sci., vol. 2, 2019, 527-537.

[48] A. K. Wanas, *Applications of (M,N)-Lucas polynomials for holomorphic and bi-univalent functions*, Filomat, vol. 34, 2020, 3361-3368.

[49] A. K. Wanas, A. L. Alina, *Applications of Horadam polynomials on Bazilevič bi-univalent function satisfying subordinate conditions*, Journal of Physics: Conf. Series, vol. 1294, 2019, 1-6.

[50] A. K. Wanas, A. H. Majeed, *On subclasses of analytic and m-fold symmetric bi-univalent functions*, Iranian Journal of Mathematical Sciences and Informatics, vol. 15, no. 2, 2020, 51-60.

[51] A. K. Wanas, G. Murugusundaramoorthy, *Differential sandwich results for Wanas operator of analytic functions*, Mathematica Moravica, vol. 24, 2020, 17-28.

[52] A. K. Wanas, S. Yalçin, *Initial coefficient estimates for a new subclasses of analytic and m-fold symmetric bi-univalent functions*, Malaya J. Mat., vol. 7, 2019, 472-476.

Abbas Kareem Wanas
University of Al-Qadisiyah
College of Science
Department of Mathematics
Al Diwaniyah, Al-Qadisiyah, Iraq
e-mail: abbas.kareem.w@qu.edu.iq

Timilehin Gideon Shaba
University of Ilorin
Department of Mathematics
P. M. B. 1515, Ilorin, Nigeria
e-mail: shabatimilehin@gmail.com