



Best proximity points of Kannan type cyclic weak ϕ -contractions in ordered metric spaces

Erdal Karapınar

Abstract

In this manuscript, the existence of the best proximity of Kannan Type cyclic weak ϕ -contraction in ordered metric spaces is investigated. Some results of Rezapour-Derafshpour-Shahzad [22] are generalized.

1 Introduction and Preliminaries

In 1922, Banach [3] stated that every contraction on a complete metric space has a unique fixed point. This theorem is known as Banach contraction mapping principle or Banach fixed point theorem. Banach's theorem preserves its importance in fixed point theory and has applications not only in many branches of mathematics but also in economics. In particular, in micro economics, for the existence of Nash equilibria, fixed point theorems are used (See e.g. [19, 4]).

In 1969, Boyd and Wong [5] gave the definition of Φ -contraction: A self-mapping T on a metric space X is called Φ -contraction if there exists an upper semi-continuous function $\Phi : [0, \infty) \rightarrow [0, \infty)$ such that

$$d(Tx, Ty) \leq \Phi(d(x, y)) \quad \text{for all } x, y \in X.$$

Later, in 1997, Alber and Guerre-Delabriere [1], introduced the definition of weak ϕ -contraction: a self-mapping T on a metric space X is called weak ϕ -contraction if for each $x, y \in X$, there exists a function $\phi : [0, \infty) \rightarrow [0, \infty)$ such that

$$d(Tx, Ty) \leq d(x, y) - \phi(d(x, y)).$$

Key Words: ϕ -contraction, best proximity, ordered metric spaces

2010 Mathematics Subject Classification: 47H10, 54H25

Received: May, 2011.

Revised: November, 2011.

Accepted: January, 2012.

In addition, Alber and Guerre-Delabriere defined weak ϕ -contraction on Hilbert spaces and proved the existence of fixed points in Hilbert spaces. Rhoades [24] showed that most of the results in [1] are also valid for arbitrary Banach spaces.

Notice that if ϕ is a lower semi-continuous mapping, then $\Phi(u) = u - \phi(u)$ becomes Φ -contraction [5]. The notions Φ -contraction and weak ϕ -contraction have been studied by many authors, (e.g., [11, 25, 26, 27, 13, 14, 15].)

Next, we give some preliminaries and basic definitions which are used throughout the paper. Let X and Y be nonempty sets and $T : X \rightarrow X$ and $S : Y \rightarrow Y$. Define the set of all invariant non-empty subsets of X under T as follows:

$$I_T(X) := \{Z \in P_0(X) : T(Z) \subset Z\} \quad (1.1)$$

where $P_0(X)$ denotes the set of all non-empty subsets of X . Moreover, a map $(T \times S) : X \times Y \rightarrow X \times Y$ is defined as

$$[T \times S](x, y) = (Tx, Sy). \quad (1.2)$$

For a partially ordered set $(X, \leq)$, we denote the set of all comparable pair in the following way:

$$CP(X) := \{(x, y) \in X \times X : \text{either } x \leq y \text{ or } y \leq x\}. \quad (1.3)$$

Let $(X, d, \leq)$ be an ordered metric space and $T : X \rightarrow X$ be a self-mapping on X . For each x^* and non-empty subset Z of X , we define

$$Z_T(x^*) := \{x \in Z : \lim_{n \rightarrow \infty} T^{2n}x = x^*\}. \quad (1.4)$$

Cyclic maps were defined by Kirk-Srinivasan-Veeramani in 2003. They stated the following theorem (see [16], Theorem 1.1).

Theorem 1. *Let A and B be non-empty closed subsets of a complete metric space (X, d) . Suppose that $T : A \cup B \rightarrow A \cup B$ is a map satisfying $T(A) \subset B$ and $T(B) \subset A$ and there exists $k \in (0, 1)$ such that $d(Tx, Ty) \leq kd(x, y)$ for all $x \in A$ and $y \in B$. Then, T has a unique fixed point in $A \cap B$.*

Definition 2. (See [2]) *Let A and B be non-empty closed subsets of a metric space (X, d) and $\phi : [0, \infty) \rightarrow [0, \infty)$ be a strictly increasing map. A map $T : A \cup B \rightarrow A \cup B$ is called a cyclic weak ϕ -contraction if $T(A) \subset B$ and $T(B) \subset A$ and*

$$d(Tx, Ty) \leq d(x, y) - \phi(d(x, y)) + \phi(d(A, B)) \quad (1.5)$$

for all $x \in A$ and $y \in B$ where $d(A, B) = \inf\{d(a, b) : a \in A, b \in B\}$.

A point $x \in A \cup B$ is called a best proximity point if $d(x, Tx) = d(A, B)$. Further, if $\alpha \in (0, 1)$ and $\phi(t) = (1 - \alpha)t$, then T is called cyclic contraction (See [9]).

Very recently, Rezapour-Derafshpour-Shahzad (see[22],also [23]) stated the following theorem:

Theorem 3. *Let $(X, d, \leq)$ be an ordered metric space, A and B be non-empty subsets of X and $T : A \cup B \rightarrow A \cup B$ be decreasing, cyclic weak ϕ -contraction, that is, T satisfies (1.5). Suppose there exists $x_0 \in A$ such that $x_0 \leq T^2 x_0 \leq Tx_0$. Define $x_{n+1} = Tx_n$ and $d_n := d(x_{n+1}, x_n)$ for all $n \in \mathbb{N}$. Then $d_n \rightarrow d(A, B)$.*

In this manuscript, Theorem 3 and some other results of [22, 13] are generalized.

2 Main Results

In this section we define Kannan type cyclic weak ϕ -contraction and state our main results.

Definition 4. *Let A and B be non-empty subsets of a metric space (X, d) and $\phi : [0, \infty) \rightarrow [0, \infty)$ be a strictly increasing map. A map $T : A \cup B \rightarrow A \cup B$ is said to be Kannan type cyclic weak ϕ -contraction if $T(A) \subset B$ and $T(B) \subset A$ and*

$$d(Tx, Ty) \leq u(x, y) - \phi(u(x, y)) + \phi(d(A, B)) \quad (2.1)$$

for all $x \in A$ and $y \in B$ where $u(x, y) = \frac{1}{2}[d(x, Tx) + d(y, Ty)]$ and $d(A, B) = \inf\{d(a, b) : a \in A, b \in B\}$.

Example 5. *Let $X := \mathbb{R}$ with the usual metric. For $A = [0, 1]$, $B = [-1, 0]$, define $T : A \cup B \rightarrow A \cup B$ by $Tx = \frac{-3x}{4}$ for all $x \in A \cup B$ and $\phi(t) = \frac{t}{7}$. Then T satisfies (2.1), i.e., T is a Kannan type cyclic weak ϕ -contraction.*

Example 6. *Consider the Euclidean ordered space $X = \mathbb{R}$ with the usual metric. Let $A = B = [0, 1]$ and $T : A \cup B \rightarrow A \cup B$ be defined by*

$$Tx = \begin{cases} \frac{1}{4} & \text{if } x = 1, \\ \frac{1}{2} & \text{if } x \in [0, 1). \end{cases}$$

If $\phi : [0, \infty) \rightarrow [0, \infty)$ is defined by $\phi(t) = \frac{t}{8}$ then T is a Kannan type cyclic weak ϕ -contraction but not a cyclic weak ϕ -contraction. Indeed, for $x = 7/8$ and $y = 1$ we have

$$\begin{aligned} d(Tx, Ty) &\leq d(x, y) - \phi[d(x, y)] + d(A, B) \\ |T\frac{7}{8} - T1| &\leq |\frac{7}{8} - 1| - \frac{1}{8}|\frac{7}{8} - 1| \\ \frac{1}{4} &\leq \frac{7}{64} \end{aligned}$$

which shows that T is not a cyclic weak ϕ -contraction.

Let us check whether T is a Kannan type cyclic weak ϕ -contraction. Notice that if both x and y are in $[0, 1)$, T obviously satisfies (2.1). If $x = y = 1$ and T again satisfies (2.1). Now, consider the case $y \in [0, 1)$ and $x = 1$. Then

$$\begin{aligned} d(Tx, Ty) &\leq \frac{1}{2} [d(Tx, x) + d(Ty, y)] - \phi\left[\frac{1}{2} [d(Tx, x) + d(Ty, y)]\right] + d(A, B) \\ |T1 - Ty| &\leq \frac{1}{2} \left[\frac{3}{4} + |y - \frac{1}{2}|\right] - \frac{1}{8} \left(\frac{1}{2} \left[\frac{3}{4} + |y - \frac{1}{2}|\right]\right) \\ \frac{1}{4} &\leq \frac{21}{64} + \frac{7}{16} |y - \frac{1}{2}| \end{aligned}$$

which holds for every $y \in [0, 1)$. Thus, T is a Kannan type cyclic weak ϕ -contraction.

Theorem 7. Let $(X, d, \leq)$ be an ordered metric space, A and B be non-empty subsets of X . Suppose $T : A \cup B \rightarrow A \cup B$ is decreasing, Kannan type cyclic weak ϕ -contraction, that is, T satisfies (2.1). Assume that there exists $x_0 \in A$ such that $x_0 \leq T^2 x_0 \leq Tx_0$. Define $x_{n+1} = Tx_n$ and $d_n := d(x_{n+1}, x_n)$ for all $n \in \mathbb{N}$. Then $d_n \rightarrow d(A, B)$.

Proof. By the assumption, one can easily observe that

$$x_0 \leq x_2 \leq \cdots \leq x_{2n} \leq x_{2n+1} \leq \cdots \leq x_3 \leq x_1, \text{ for all } n \in \mathbb{N}.$$

Due to (2.1), we have

$$\begin{aligned} d_{n+1} &= d(x_{n+2}, x_{n+1}) = d(Tx_{n+1}, Tx_n) \\ &\leq \frac{1}{2} [d(x_{n+1}, Tx_{n+1}) + d(x_n, Tx_n)] - \phi\left(\frac{1}{2} [d(x_{n+1}, Tx_{n+1}) + d(x_n, Tx_n)]\right) + \phi(d(A, B)) \\ &\leq \frac{1}{2} [d_{n+1} + d_n] - \phi\left(\frac{1}{2} [d_{n+1} + d_n]\right) + \phi(d(A, B)). \end{aligned} \quad (2.2)$$

Notice that $d(A, B) \leq \frac{1}{2} [d(x_{n+1}, x_{n+2}) + d(x_n, x_{n+1})]$. Since ϕ is a strictly increasing map, we have

$$\phi(d(A, B)) \leq \phi\left(\frac{1}{2} [d(x_{n+1}, x_{n+2}) + d(x_n, x_{n+1})]\right). \quad (2.3)$$

Thus, regarding $0 \leq d_{n+1}$, the expression (2.2) turns into

$$0 \leq d_{n+1} \leq \frac{1}{2} [d_{n+1} + d_n] - \phi\left(\frac{1}{2} [d_{n+1} + d_n]\right) + \phi(d(A, B)) \quad (2.4)$$

which implies $0 \leq \frac{1}{2} d_{n+1} \leq \frac{1}{2} d_n$. Hence, the sequence $\{d_n\}$ is non-increasing and bounded below. If $d_{n_0} = 0$ for some $n_0 \in \mathbb{N}$, then clearly $d_n \rightarrow 0$, $d(A, B) = 0$ and so, $d_n \rightarrow d(A, B)$. Suppose $d_n \neq 0$ (that is, $d_n > 0$) for all $n \in \mathbb{N}$ and $d_n \rightarrow L$ for some $L \geq d(A, B)$. From (2.4), we obtain that

$$\phi\left(\frac{1}{2} [d_{n+1} + d_n]\right) \leq \frac{1}{2} [d_{n+1} + d_n] + \phi(d(A, B)) - d_{n+1}. \quad (2.5)$$

If we combine (2.3) and (2.5) we get

$$\phi(d(A, B)) \leq \phi\left(\frac{1}{2}[d_{n+1} + d_n]\right) \leq \frac{1}{2}[d_{n+1} + d_n] - d_{n+1} + \phi(d(A, B)) \quad (2.6)$$

which yields that $\phi(L) = \phi(d(A, B))$. Since ϕ is strictly increasing, then $L = d(A, B)$ which completes the proof. $\square$

The following result is a consequence of Theorem 7 and Theorem 2.2 of [22]. Regarding analogy, we omit the proof.

Corollary 8. *Let $(X, d, \leq)$ be an ordered metric space, A and B be non-empty subsets of X and $T : A \cup B \rightarrow A \cup B$ be decreasing map satisfying*

$$d(Tx, Ty) \leq m(x, y) - \phi(m(x, y)) + \phi(d(A, B)) \quad (2.7)$$

for all $x \in A$ and $y \in B$ where $m(x, y) = \max\{d(x, y), \frac{1}{2}[d(x, Tx) + d(y, Ty)]\}$. Suppose that there exists $x_0 \in A$ such that $x_0 \leq T^2x_0 \leq Tx_0$. Define $x_{n+1} = Tx_n$ and $d_n := d(x_{n+1}, x_n)$ for all $n \in \mathbb{N}$. Then $d_n \rightarrow d(A, B)$.

Our next results are concerned with orbital continuous maps and topological spaces with C -condition.

Definition 9. (See [22]) *A topological space X is said to have condition (C) if for each convergent monotone sequence $\{x_n\}$ (that is, $x_n \rightarrow x$, $x \in X$), there exists a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that every element of $\{x_{n_k}\}$ is comparable with the limit x . Further, X is called regular if every bounded monotone sequence in X is convergent.*

Definition 10. (See [6, 7], see also [12]) *A mapping T on metric space (X, d) is said to be orbitally continuous if $\lim_{i \rightarrow \infty} T^{n_i}(x) = z$ implies $\lim_{i \rightarrow \infty} T(T^{n_i}(x)) = Tz$.*

Remark 11. *It is clear that orbital continuity of T implies orbital continuity of T^m for any $m \in \mathbb{N}$.*

Theorem 12. *Let $(X, d, \leq)$ be a regular ordered metric space, A and B be non-empty subsets of X where A is closed. Assume $T : A \cup B \rightarrow A \cup B$ is a decreasing map satisfying Kannan type cyclic weak ϕ -contraction, that is, T satisfies (2.1). Suppose that there exists $x_0 \in A$ such that $x_0 \leq T^2x_0 \leq Tx_0$. Define $x_{n+1} = Tx_n$ and $d_n := d(x_{n+1}, x_n)$ for all $n \in \mathbb{N}$. If T is orbitally continuous or X has the property (C), then there exists $x \in A$ such that $d(x, Tx) = d(A, B)$.*

Proof. By the assumption, one can easily see that

$$x_0 \leq x_2 \leq \cdots \leq x_{2n} \leq x_1, \quad \text{for all } n \in \mathbb{N}.$$

Regarding that X is regular and A is closed, the sequence $\{x_{2n}\}$ converges to a point, say $x \in A$. Observe that

$$d(A, B) \leq d(x_{2n}, Tx) = d(Tx_{2n-1}, Tx) \leq d(Tx_{2n-1}, Tx_{2n}) + d(Tx_{2n}, Tx), \quad (2.8)$$

for all $n \in \mathbb{N}$. If T is orbitally continuous, then $d(Tx_{2n}, Tx) \rightarrow 0$. Regarding Theorem 7, we have $d(Tx_{2n-1}, Tx_{2n}) \rightarrow d(A, B)$ and hence

$$d(x, Tx) = d(A, B)$$

due to (2.8). Suppose now that X has the property (C). Taking the fact that the sequence $\{x_{2n}\}$ is bounded and increasing into account, one can find a subsequence $\{x_{2n_k}\}$ of $\{x_{2n}\}$ such that

$$x_{2n_1} \leq x_{2n_2} \leq \cdots \leq x_{2n_k} \leq \cdots \leq x.$$

Hence,

$$\begin{aligned} d(A, B) &\leq d(x_{2n_k}, Tx) = d(x_{2n_k-1}, Tx) \\ &\leq d(x_{2n_k-1}, Tx_{2n_k}) + d(Tx_{2n_k}, Tx) \\ &\leq d(x_{2n_k-1}, Tx_{2n_k}) + d(x_{2n_k}, Tx) \end{aligned} \quad (2.9)$$

which implies that $d(x, Tx) = d(A, B)$. $\square$

The following corollary results from Theorem 7 and Theorem 2.3 of [22]. The proof is omitted because of analogy.

Corollary 13. *Let $(X, d, \leq)$ be a regular ordered metric space, A and B be non-empty subsets of X where A is closed. Assume that $T : A \cup B \rightarrow A \cup B$ is a decreasing map satisfying*

$$d(Tx, Ty) \leq m(x, y) - \phi(m(x, y)) + \phi(d(A, B)) \quad (2.10)$$

for all $x \in A$ and $y \in B$ where $m(x, y) = \max\{d(x, y), \frac{1}{2}[d(x, Tx) + d(y, Ty)]\}$. Suppose that there exists $x_0 \in A$ such that $x_0 \leq T^2x_0 \leq Tx_0$. Define $x_{n+1} = Tx_n$ and $d_n := d(x_{n+1}, x_n)$ for all $n \in \mathbb{N}$. If T is orbitally continuous or X has the property (C), then there exists $x \in A$ such that $d(x, Tx) = d(A, B)$.

Corollary 14. *Let $(X, d, \leq)$ be an ordered metric space, A and B be non-empty subsets of X . Define $T : A \cup B \rightarrow A \cup B$ such that $T(A) \subset B$, $T(B) \subset A$*

and $(A \times B) \cap CP(X) \in I_T(X)$. Assume that there exists $x_0 \in A$ such that $(x_0, Tx_0) \in CP(X)$ and

$$d(Tx, Ty) \leq \frac{1}{2}[d(x, Tx) + d(Ty, y)] - \phi\left(\frac{1}{2}[d(x, Tx) + d(Ty, y)]\right) + \phi(d(A, B)), \quad (2.11)$$

for all $x \in A$ and $y \in B$ with $(x, y) \in CP(X)$, where $\phi : [0, \infty) \rightarrow [0, \infty)$ is a strictly increasing map. Define $x_{n+1} = Tx_n$ and $d_n := d(x_{n+1}, x_n)$ for all $n \in \mathbb{N}$. If T is orbitally continuous or X has the property (C), then $d_n \rightarrow d(A, B)$.

Proof. By the assumption, it is easy to see that,

$$\begin{aligned} d(T^{2n+1}x_0, T^{2n}x_0) &\leq \frac{1}{2}[d(T^{2n}x_0, T^{2n+1}x_0) + d(T^{2n}x_0, T^{2n-1}x_0)] \\ &\quad - \phi\left(\frac{1}{2}[d(T^{2n}x_0, T^{2n+1}x_0) + d(T^{2n}x_0, T^{2n-1}x_0)]\right) + \phi(d(A, B)), \end{aligned} \quad (2.12)$$

for all $n \in \mathbb{N}$. Clearly, $d(A, B) \leq \frac{1}{2}[d(T^{2n}x_0, T^{2n+1}x_0) + d(T^{2n}x_0, T^{2n-1}x_0)]$. Since ϕ is a strictly increasing map, then

$$\phi(d(A, B)) \leq \phi\left(\frac{1}{2}[d(T^{2n}x_0, T^{2n+1}x_0) + d(T^{2n}x_0, T^{2n-1}x_0)]\right).$$

Regarding $0 \leq d_{n+1}$, the expression (2.12) becomes

$$0 \leq d_{n+1} \leq \frac{1}{2}[d_{n+1} + d_n] - \phi\left(\frac{1}{2}[d_{n+1} + d_n]\right) + \phi(d(A, B)) \quad (2.13)$$

which implies that $0 \leq \frac{1}{2}d_{n+1} \leq \frac{1}{2}d_n$. Hence, $\{d_n\}$ is decreasing and bounded below. If $d_{n_0} = 0$ for some $n_0 \in \mathbb{N}$, then clearly $d_n \rightarrow d(A, B) = 0$. Consider the other case $d_n > 0$ for all $n \in \mathbb{N}$. Let $d_n \rightarrow L$ for some $L \geq d(A, B)$. By the assumptions, we have

$$\phi(d(A, B)) \leq \phi\left(\frac{1}{2}[d_{n+1} + d_n]\right) \leq \frac{1}{2}[d_{n+1} + d_n] - d_{n+1} + \phi(d(A, B)) \quad (2.14)$$

which yields $\phi(d_n) \rightarrow \phi(d(A, B))$, and therefore $\phi(L) = \phi(d(A, B))$. Since ϕ is strictly increasing, $L = d(A, B)$. $\square$

The following result is a consequence of Theorem 14 and Theorem 2.3 of [22]. Regarding analogy, we omit the proof.

Corollary 15. Let $(X, d, \leq)$ be an ordered metric space, A and B be non-empty subsets of X . Define $T : A \cup B \rightarrow A \cup B$ such that $T(A) \subset B$, $T(B) \subset A$ and $(A \times B) \cap CP(X) \in I_T(X)$. Assume that there exists $x_0 \in A$ such that $(x_0, Tx_0) \in CP(X)$ and

$$d(Tx, Ty) \leq m(x, y) - \phi(m(x, y)) + \phi(d(A, B)) \quad (2.15)$$

for all $x \in A$ and $y \in B$ with $(x, y) \in CP(X)$ where

$$m(x, y) = \max\{d(x, y), \frac{1}{2}[d(x, Tx) + d(y, Ty)]\}$$

and $\phi : [0, \infty) \rightarrow [0, \infty)$ is a strictly increasing map. Define $x_{n+1} = Tx_n$ and $d_n := d(x_{n+1}, x_n)$ for all $n \in \mathbb{N}$. If T is orbitally continuous or X has the property (C), then $d_n \rightarrow d(A, B)$.

Theorem 16. Let $(X, d, \leq)$ be an ordered metric space, A and B be non-empty subsets of X . Define $T : A \cup B \rightarrow A \cup B$ such that $T(A) = B$, $T(B) \subset A$ and $(A \times B) \cap CP(X) \in I_T(X)$. Assume that there exist $x, y \in A$ such that $(x, z), (z, y) \in CP(X)$. Suppose also that there exist $x_0, u \in A$ such that $x_0 \in A_T(u)$, $(x_0, Tx_0) \in CP(X)$ and

$$d(Tx, Ty) \leq \frac{1}{2}[d(x, Tx) + d(Ty, y)] - \phi(\frac{1}{2}[d(x, Tx) + d(Ty, y)]) + \phi(d(A, B)) \quad (2.16)$$

for all $x \in A$ and $y \in B$ with $(x, y) \in CP(X)$ where $\phi : [0, \infty) \rightarrow [0, \infty)$ is a strictly increasing map. Moreover, $y \in A$, $(x, y) \in CP(X)$ and $x \in A_T(u)$ implies that $y \in A_T(u)$. Then, $A_T(u) = A$ and the following holds:

$$B_T(Tu) = B \text{ and } d(u, Tu) = d(A, B) \Leftrightarrow T \text{ is orbitally continuous.} \quad (2.17)$$

Proof. First, we prove that $A_T(u) = A$. Take $x \in A$. If $(x, x_0) \in CP(X)$, then $x \in A_T(u)$ and so $A_T(u) = A$. If $(x, x_0) \notin CP(X)$, then by assumption, there exists $z \in A$ such that $(x, z), (x_0, z) \in CP(X)$. Therefore, $x \in A_T(u)$. Hence, in any case, $A_T(u) = A$.

Now we show (2.17). Assume that T is orbitally continuous. Fix $y \in B$. Choose $\tilde{x} \in A$ such that $T\tilde{x} = y$. Since $A_T(u) = A$ and $T^{2n}\tilde{x} \rightarrow u$, then $T^{2n+1}\tilde{x} \rightarrow Tu$. That is, $T^{2n}y \rightarrow Tu$ and thus $B_T(Tu) = B$. If $d(A, B) = d(u, Tu)$ then the proof is done. Assume that $d(A, B) \neq d(u, Tu)$. Since $d(x_0, Tx_0) \in CP(X)$ and by (2.16), the sequence $\{d(T^{2n+1}x_0, T^{2n}x_0)\}$ is decreasing. Hence, Theorem 14 implies that $d(T^{2n+1}x_0, T^{2n}x_0) \downarrow d(A, B)$. Choose $n \in \mathbb{N}$ in a way that

$$d(A, B) \leq d(T^{2n+1}x_0, T^{2n}x_0) < d(u, Tu). \quad (2.18)$$

Substitute $r = T^{2n+1}x_0$ and $s = T^{2n}x_0$. Since $(r, s) \in CP(X)$ then $(Tr, Ts) \in CP(X)$ and thus $\{d(T^{2n}r, T^{2n}s)\}$ is a decreasing sequence.

Hence $d(T^{2n}r, T^{2n}s) \downarrow d(u, Tu)$ and so $d(u, Tu) \leq d(r, s) = d(T^{2n+1}x_0, T^{2n}x_0) < d(u, Tu)$ which contradicts (2.18). Thus, $d(A, B) = d(u, Tu)$.

To prove the inverse inclusion of (2.17), assume that $B_T(Tu) = B$ and $d(A, B) = d(u, Tu)$. Let $x \in A \cup B$ and $T^{n(i)}x \rightarrow w$ for some $w \in A \cup B$. To complete the proof, it is sufficient to show that T is orbitally continuous, that is, $T^{n(i)+1}x \rightarrow Tw$. Define the subsets $S_A = A \cap \{T^{n(i)}x\}$ and $S_B = B \cap \{T^{n(i)}x\}$. Observe that S_A and S_B are subsequences of $\{T^{n(i)}x\}$, that is, $S_A = A \cap \{T^{n(i)}x\} = \{T^{n_1(i)}x\}$ and $S_B = B \cap \{T^{n(i)}x\} = \{T^{n_2(i)}x\}$. We consider the following two cases:

The first case, $d(u, Tu) = d(A, B) = 0$, in other words $Tu = u$. On account of Theorem 14 and using the fact that $\{T^{n_1(i)}x\}$ is a subsequence of $\{T^{n(i)}x\}$ and the assumption $T^{n(i)}x \rightarrow w$, we get $T^{n_1(i)}x \rightarrow w$. On the other hand, $\{T^{n_1(i)}x\}$ is also a subsequence of $\{T^{2n}x\}$ and thus, $T^{n_1(i)}x \rightarrow u$. Hence, we conclude that $u = w$, and further $w = u = Tu = Tw$. Then we have

$$\begin{aligned} d(T(T^{n(i)}x), Tw) &\leq \frac{1}{2}[d(T^{n(i)}x, T(T^{n(i)}x)) + d(Tw, w)] \\ &\quad - \phi(\frac{1}{2}[d(T^{n(i)}x, T(T^{n(i)}x)) + d(Tw, w)]) + \phi(d(A, B)) \\ &\leq \frac{1}{2}[d(T^{n(i)}x, T(T^{n(i)}x)) + d(Tw, w)] \\ &\leq \frac{1}{2}[d(T^{n(i)}x, T(T^{n(i)}x))] \\ &\leq \frac{1}{2}[d(T^{n(i)}x, Tw) + d(T^{n(i)}x, w)] \quad (\text{since } Tw = w) \end{aligned} \tag{2.19}$$

which implies that $\frac{1}{2}d(T(T^{n(i)}x), Tw) \leq \frac{1}{2}d(T^{n(i)}x, w)$, or equivalently, $T(T^{n(i)}x) \rightarrow Tw$. Hence, T is orbitally continuous.

Consider the second case, $d(u, Tu) = d(A, B) > 0$. We assert that both S_A or S_B are finite. Indeed, if both S_A and S_B are infinite, then as in the first case, $T^{n_1(i)+1}x \rightarrow u$ and $T^{n_2(i)+1}x \rightarrow Tu$. Since both $\{T^{n_1(i)}x\}$ and $\{T^{n_2(i)}x\}$ are subsets of $\{T^{n(i)}x\}$ and $T^{n(i)}x \rightarrow w$, then both subsequences converge to w and thus $w = Tu = u$. So $d(A, B) = d(u, Tu) = 0$ which is a contradiction. Therefore, either S_A or S_B is finite. Assume that $S_B = \{b_1, b_2, \dots, b_m\}$ is finite. As in the first case, we get that $T^{n_1(i)}x \rightarrow u$, $T^{n_1(i)+1}x \rightarrow Tu$ and $w = u$. Then, $\{T^{n(i)+1}x\} = \{T^{n_1(i)}x\} \cup \{Tb_1, \dots, Tb_m\}$ and $T^{n(i)+1}x \rightarrow Tw$. Suppose now that $S_A = \{a_1, a_2, \dots, a_j\}$ is finite and recall that $\{T^{n_2(i)}x\}$ is a subsequence of $\{T^{n(i)}x\}$. Then, $T^{n_2(i)}x \rightarrow w$. Moreover, $\{T^{n_2(i)}x\}$ is also a subsequence of $\{T^{2n+1}x\} = T^{2n}(Tx)$ and $Tx \in B = B_T(Tu)$, so $T^{n_2(i)}x \rightarrow Tu$. Hence, $w = Tu$. Due to the fact that $\{T^{n_2(i)}x\}$ is a subset of $\{T^{2n+2}x\} = \{T^{2n}(T^2x)\}$ and $T^2x \in A = A_T(u)$, we have

$$T^{n_2(i)+1}x \rightarrow u. \tag{2.20}$$

We claim that $Tw = u$. Since $(u, u) \in CP(X)$, by triangle inequality we have $d(u, T^2u) \leq d(u, T^{2n}u) + d(T^{2n}u, T^2u)$. By assumptions of the theorem, we have $d(T^{2n}u, T^2u) \leq d(T^{2n-2}u, u)$ which implies that

$$d(u, T^2u) \leq d(u, T^{2n}u) + d(T^{2n-2}u, u).$$

Taking into account that $u \in A = A_T(u)$, we have $T^{2n}u \rightarrow u$ and $T^{2n-2}u \rightarrow u$. Hence, $u = T^2u$. Since $w = Tu$, then $Tw = T^2u = u$. So, the equation (2.20) becomes $T^{n_2(i)+1} \rightarrow u = Tw$. Thus, $T^{n(i)+1}x \rightarrow Tw$ where $\{T^{n(i)+1}x\} = \{T^{n_2(i)}x\} \cup \{Ta_1, \dots, a_j\}$ which completes the proof. $\square$

Notice that the condition

$$d(Tx, Ty) \leq \frac{1}{2}[d(Tx, x) + d(Ty, y)] - \phi\left(\frac{1}{2}[d(Tx, x) + d(Ty, y)]\right) + d(A, B),$$

for all $x \in A, y \in B$ with $(x, y) \in CP(X)$, does not imply

$$y \in A, (x, y) \in CP(X), x \in A_T(u) \Rightarrow y \in A_T(u).$$

The following example is given to support our assertion:

Example 17. Let $X := \mathbb{R}^2$ with the usual metric and the following order:

$$(a, b) \leq (c, d) \Leftrightarrow a \leq c \text{ and } c \leq d.$$

For $A = \{x_1 = (5, 2), x_2 = (1, 2)\}, B = \{y_1 = (3, 0), y_2 = (0, 4)\}$, define $T : A \cup B \rightarrow A \cup B$ by $Tx_1 = y_2, Tx_2 = y_1, Ty_1 = x_2, Ty_2 = x_1$. Here, $x_2 \leq x_1$ and $y_1 \leq x_1$ and the others are not comparable. Let $\phi(t) = \frac{t}{3}$. Observe that $d(Tx_1, Ty_1) = d(x_2, y_2) = d(A, B) = \sqrt{5}$ and $\frac{1}{2}[d(Tx_1, x_1) + d(Ty_1, y_1)] = \frac{1}{2}[\sqrt{8} + \sqrt{29}]$. Then we have

$$d(Tx_1, Ty_1) \leq \frac{1}{2}[d(Tx_1, x_1) + d(Ty_1, y_1)] - \phi\left(\frac{1}{2}[d(Tx_1, x_1) + d(Ty_1, y_1)]\right) + d(A, B),$$

while $T^{2n}x_1 \rightarrow x_1$ and $T^{2n}x_2 \rightarrow x_2$.

Last we give a consequence of Theorem 7 and Theorem 2.4 of [22].

Corollary 18. Let $(X, d, \leq)$ be an ordered metric space, A and B be non-empty subsets of X . Define $T : A \cup B \rightarrow A \cup B$ such that $T(A) = B, T(B) \subset A$ and $(A \times B) \cap CP(X) \in I_T(X)$. Assume that there exist $x, y \in A$ such that $(x, z), (z, y) \in CP(X)$. Suppose that there exist $x_0, u \in A$ such that $x_0 \in A_T(u), (x_0, Tx_0) \in CP(X)$ and

$$d(Tx, Ty) \leq m(x, y) - \phi(m(x, y)) + \phi(d(A, B)) \quad (2.21)$$

for all $x \in A$ and $y \in B$ with $(x, y) \in CP(X)$ where

$$m(x, y) = \max\left\{d(x, y), \frac{1}{2}[d(x, Tx) + d(y, Ty)]\right\}$$

and $\phi : [0, \infty) \rightarrow [0, \infty)$ is a strictly increasing map. Moreover, $y \in A$, $(x, y) \in CP(X)$ and $x \in A_T(u)$ implies that $y \in A_T(u)$. Then, $A_T(u) = A$ and the following hold:

$$B_T(Tu) = B \quad \text{and} \quad d(u, Tu) = d(A, B) \Leftrightarrow T \text{ is orbitally continuous.} \quad (2.22)$$

Regarding analogy, we omit the proof.

References

- [1] Ya. I. Alber, S. Guerre-Delabriere, Principle of weakly contractive maps in Hilbert space In: *New Results in Operator Theory, Advances and Appl.* (Eds: I. Gohberg and Yu. Lyubich,) 722, Birkhäuser, Basel, Oper. Theory Adv. Appl., **98**, (1997).
- [2] M.A. Al-Thafai, N. Shahzad, Convergence and existence for best proximity points, *Nonlinear Analysis*, **70**(2009) 3665–3671 .
- [3] S. Banach, Sur les operations dans les ensembles abstraits et leur application aux equations itegrales, *Fund. Math.*, **3**(1922)133–181 .
- [4] K.C.Border, Fixed Point Theorems with Applications to Economics and Game Theory , Cambridge University Press, Newyork,1985.
- [5] D.W. Boyd, and S.W. Wong, On nonlinear contractions, *Proc. Amer. Math. Soc.* **20** (1969) 458-464.
- [6] Lj. B. Ćirić, On contraction type mappings *Math. Balkanica*, **1**(1971) 52–57 .
- [7] Lj. B. Ćirić, On some maps with a nonunique fixed point. *Publ. Inst. Math.*, **17**(1974) 52–58 .
- [8] H.S.Ding,L. Li, and S. Radenovic, Bhaskar-Lakshmikantham-Ćirić-type theorems in partially ordered metric space, preprint.
- [9] A.A.Eldered, P.Veeramani, Proximal pointwise contraction. *Topology and its Applications*, **156** (2009) 2942–2948.
- [10] A.A.Eldered, P.Veeramani, Convergence and existence for best proximity points. *J. Math. Anal. Appl.*, **323** (2006)1001–1006.

- [11] N.Hussain, G. Jungck, Common fixed point and invariant approximation results for noncommuting generalized (f, g) -nonexpansive maps, *J. Math. Anal. Appl.*, **321**(2006) 851-861.
- [12] E.Karapınar, Some Nonunique Fixed Point Theorems of Ćirić type on Cone Metric Spaces *Abst. Appl. Anal.*, **2010**, Article id 123094, 14 pages (2010), doi:10.1155/2010/123094 .
- [13] E.Karapınar, Fixed point theory for cyclic weak ϕ -contraction, *Appl.Math.Lett.*, **24**(2011),no : 6, 822-825.
- [14] E.Karapınar, Best proximity points of cyclic mappings, *Appl.Math.Lett.*, Volume 335, Issue 1, 1 November 2007, Pages 79-92.
- [15] E.Karapınar, İ.M. Erhan and A.Y.Ulus.: Fixed Point Theorem for Cyclic Maps on Partial Metric Spaces, *Appl. Math. Inf. Sci.*, **6** (2012), no:1, 239-244.
- [16] W.A. Kirk, P.S. Srinivasan, and P. Veeramani, Fixed Points for mapping satisfying cyclical contractive conditions *Fixed Point Theory*, **4** (2003)79–89.
- [17] Mitrovic, Z. D. : A coupled best approximations theorem in normed spaces, *Nonlinear Anal.* 72 (2010), 4049–4052.
- [18] J.J.Nieto, R.L. and Pouso, R. Rodriguez-Lopez, Fixed point theorems in ordered abstract spaces, *Proc. Amer. Math. Soc.*, **137** (2007) 2505–2517.
- [19] E.A.Ok, Real Analysis with Economic Applications, Princeton University Press, 2007.
- [20] G.Petruşel, Cyclic representations and periodic points, *Studia Univ. Babeş-Bolyai Math.*, **50** (2005) 107–112.
- [21] G. Petruşel and I.A. Rus, Fixed point theorems in ordered L-spaces *Proc. Amer. Math. Soc.*, **134**(2006) 411–418.
- [22] Sh.Rezapour, M. Derafshpour, N. Shahzad, Best proximity point of cyclic φ -contractions in ordered metric spaces, *Topological Methods in Nonlinear Analysis*, **37**(2011)193-202.
- [23] Sh.Rezapour, M. Derafshpour, N. Shahzad, Best Proximity Points of Cyclic φ -Contractions on Reflexive Banach Spaces, *Fixed Point Theory and Applications*, **2010**, Article id 946178, 7 pages,(2010).

- [24] B. E. Rhoades, Some theorems on weakly contractive maps. *Nonlinear Anal.*, **47** (2001), no:4, 2683–2693.
- [25] Y. Song, Coincidence points for noncommuting f-weakly contractive mappings, *Int. J. Comput. Appl. Math.*, **2**(2007),no:1, 51-57 .
- [26] Y.Song, S. Xu, A note on common fixed-points for Banach operator pairs, *Int. J. Contemp. Math. Sci.*, **2**(2007) 1163-1166.
- [27] Q. Zhang, and Y. Song, Fixed point theory for generalized φ -weak contractions, *Appl. Math. Lett.*, **22**(2009),no:1, 7578.

Department of Mathematics,
Atilim University
06836, Incek, Ankara, Turkey
e-mail: erdalkarapinar@yahoo.com

